

Adherence to Mediterranean diet and risk of developing diabetes: prospective cohort study

M Á Martínez-González, professor of epidemiology and chair,¹ C de la Fuente-Arrillaga, research assistant,¹ J M Nunez-Cordoba, research fellow,^{1,2} F J Basterra-Gortari, research fellow,^{1,3} J J Beunza, assistant professor,¹ Z Vazquez, research assistant,¹ S Benito, research assistant,¹ A Tortosa, research fellow,¹ M Bes-Rastrollo, assistant professor¹

¹Department of Preventive Medicine and Public Health, Medical School-Clinica Universitaria, University of Navarra, Pamplona, Spain

²Department of Preventive Medicine and Quality Management, Hospital Virgen del Camino, Pamplona, Spain

³Department of Endocrinology, Hospital of Navarra, Pamplona, Spain

Correspondence to: M Á Martínez-González mamartinez@unav.es

doi:10.1136/bmj.39561.501007.BE

ABSTRACT

Objective To assess the relation between adherence to a Mediterranean diet and the incidence of diabetes among initially healthy participants.

Design Prospective cohort study with estimates of relative risk adjusted for sex, age, years of university education, total energy intake, body mass index, physical activity, sedentary habits, smoking, family history of diabetes, and personal history of hypertension.

Setting Spanish university department.

Participants 13 380 Spanish university graduates without diabetes at baseline followed up for a median of 4.4 years.

Main outcome measures Dietary habits assessed at baseline with a validated 136 item food frequency questionnaire and scored on a nine point index. New cases of diabetes confirmed through medical reports and an additional detailed questionnaire posted to those who self reported a new diagnosis of diabetes by a doctor during follow-up. Confirmed cases of type 2 diabetes.

Results Participants who adhered closely to a Mediterranean diet had a lower risk of diabetes. The incidence rate ratios adjusted for sex and age were 0.41 (95% confidence interval 0.19 to 0.87) for those with moderate adherence (score 3-6) and 0.17 (0.04 to 0.75) for those with the highest adherence (score 7-9) compared with those with low adherence (score <3). In the fully adjusted analyses the results were similar. A two point increase in the score was associated with a 35% relative reduction in the risk of diabetes (incidence rate ratio 0.65, 0.44 to 0.95), with a significant inverse linear trend (P=0.04) in the multivariate analysis.

Conclusion Adherence to a Mediterranean diet is associated with a reduced risk of diabetes.

INTRODUCTION

Diabetes mellitus is an increasingly important global public health problem that threatens to reach pandemic levels by 2030.^{1,2} As some randomised trials have consistently shown, increased physical activity and weight loss are efficient approaches for the control and prevention of type 2 diabetes.^{3,4} Diet has also long been believed to be an important risk factor for

diabetes. Many studies have shown that the Mediterranean food pattern has a role in prevention of cardiovascular disease.⁵⁻⁹ The similarity of some risk factors and some empirical and mechanistic evidence suggest that the Mediterranean diet can also protect against diabetes. The major protective characteristics include a high intake of fibre, a high intake of vegetable fat, a low intake of *trans* fatty acids, and a moderate intake of alcohol.^{4,10,11} Moreover, a particular feature of the diet is the abundant use of virgin olive oil for cooking, frying, spreading on bread, or dressing salads.¹² This leads to a high ratio of monounsaturated fatty acids to saturated fatty acids. This ratio can be used to score adherence to a Mediterranean diet⁸ as the traditional diet. Despite having a relatively high total fat content, this food pattern is rich in monounsaturated fatty acids (from olive oil) and poor in saturated fatty acids. Diets rich in monounsaturated fatty acids improve lipid profiles and glycaemic control in people with diabetes, suggesting that a high intake improves insulin sensitivity.¹³⁻¹⁶ Together these associations suggest the hypothesis that following an overall pattern of Mediterranean diet can protect against diabetes. In addition to having a long tradition of use without evidence of harm, a Mediterranean diet is highly palatable, and people are likely to comply with it.¹⁷

Few prospective studies have evaluated the specific role of the Mediterranean diet on the risk of developing diabetes in initially healthy Mediterranean populations. A prospective follow-up study recently reported an inverse association between adherence to the diet and the incidence of diabetes.¹⁸ All members of that study, however, had survived a previous myocardial infarction and the tool to assess dietary habits had inherent limitations because it included only a short list of items, and the authors did not attempt to measure the entire diet.

We evaluated the association between adherence to a Mediterranean diet and the incidence of diabetes using a full validated food frequency questionnaire to measure the entire diet.

METHODS

Study population

The objectives, design, and methods of the SUN project ("Seguimiento Universidad de Navarra" (University of Navarra follow-up) have been described elsewhere.^{19,20} Briefly, we invited all former students of the University of Navarra, registered nurses from Spanish provinces, and university graduates from other associations to take part in the project. We posted them a questionnaire and an invitation letter to participate in a long term follow-up study. The first participants were recruited in December 1999. Using successive waves of invitation to different groups of university graduates, we have increased the sample size of the cohort. The design is that of a dynamic cohort and the recruitment is permanently open. We considered a response to the initial questionnaire as informed consent to participate in the study.

After the baseline assessment, every two years we send participants follow-up questionnaires on diet, lifestyle, risk factors, and medical conditions.²⁰ Participants who do not reply to the follow-up questionnaires are sent up to five additional mailings. Participants are aged 20-90 years at baseline.

Up to November 2007, the project had enrolled 18 700 participants. From these, 15 970 were eligible for follow-up because they had taken part for more than two years and had completed the follow-up process (entering the cohort up to March 2005), of whom 13 753 (86.1%) had completed at least their two year follow-up questionnaire. Among them, 9298 participants who had taken part for more than four years had completed at least the four year follow-up questionnaire and 4993 had completed the six year follow-up questionnaire. For this analysis, we excluded participants who had diabetes at baseline and those who reported a baseline history of cardiovascular disease. This left 13 380 for analyses.

Dietary assessment

We assessed dietary habits at baseline using a semi-quantitative food frequency questionnaire with 136 items, previously validated in Spain.²¹ In the validation study, the correlation coefficients between the questionnaire and diet records ranged from 0.45 for vitamin A to 0.91 for alcohol.²¹ With regard to gross misclassification, on average 3% of participants classified in the highest or lowest fifth of the distribution by food record were assigned to the lowest or highest fifth by the food frequency questionnaire.²¹ The questionnaire was based on typical portion sizes and had nine options for the frequency of intake in the previous year for each food item (ranging from never or almost never to six or more times a day). We used the sum of the consumption of each of several food items to estimate the overall consumption of the food group to which each item belonged (vegetables, fruits, cereals, legumes, fish, meats, fast food, dairy products). We applied the specified portion sizes for each item and summed up across all foods of that group. Then we

computed the mean for all the participants in each category of the three categories of the score to create an estimated daily intake.

Nutrient intake scores were computed with a specifically developed computer program. A trained dietitian updated the nutrient data bank using the latest available information included in food composition tables for Spain. The questionnaire included questions on use of fats and oils, cooking methods, and dietary supplements. For example, we calculated total consumption of olive oil (a major source of monounsaturated fat in this population) from the nutrient score of olive oil consumption, taking into account the amount and type of olive oil used for cooking or frying and the use of olive oil for salad dressing or as a spread on bread or other foods.

Adherence to the Mediterranean diet was appraised according to the score created by Trichopoulou et al,²² which is extensively used because it is simple and has variants created to evaluate multiple diet-health relations.²³ Originally, this index included only eight components to define the Mediterranean diet²²: high ratio of monounsaturated:saturated fatty acids, moderate intake of alcohol, high intake of legumes, high intake of grains, high intake of fruit and nuts, high intake of vegetables, low intake of meat and meat products, and moderate intake of milk and dairy products. The same authors subsequently added high intake of fish.⁸ The index assigns a score of 0 or 1 according to the daily intake of each of the nine components. With the exception of alcohol, the sex specific medians of the sample were used as cut-off points and the unit of measurement was grams per day.^{8,22-24} For each of the six protective components (fatty acid ratio, legumes, grains, fruits, vegetables, or fish) participants received one point if their intake was over the sample median. Participants received one point if the intake was below the median for the two non-protective components (dairy products or meat). For alcohol one point was scored if consumption was 10-50 g/day for men or 5-25 g/day for women. If participants met all the characteristics of the Mediterranean diet, their score was the highest (nine points), reflecting maximum adherence. If they met none of the characteristics the score was minimum (zero), reflecting no adherence at all.

Assessment of other covariates

The baseline questionnaire requested information about the main known risk factors for diabetes. We also collected information on sociodemographic factors (sex, age, university degree, and marital and employment status), anthropometric measurements (weight, height), health related habits (smoking status, physical activity, sedentary lifestyle), and clinical variables (use of medication, personal and family history of diabetes and hypertension, coronary heart disease, and other cardiovascular disease). The validity of self reported weight and body mass index (BMI) has been previously documented in a subsample of this

cohort.²⁵ We assessed the participation and time spent in 17 different activities, using a physical activity questionnaire previously validated in Spain.²⁶ The time spent at each activity in hours each week was multiplied by its typical energy expenditure, expressed in metabolic equivalent tasks (METs), then summed over all activities to yield a METs-hour/week score for each participant.²⁷

Ascertainment of diabetes

Participants reported any medical diagnosis of diabetes at baseline and in each of the follow-up questionnaires. The baseline questionnaire also collected data on use of insulin or oral antidiabetic agents. Follow-up questionnaires recorded the date of any new diagnosis of diabetes (if applicable).

Participants were considered to have diabetes at baseline if they reported a medical diagnosis of diabetes or were receiving insulin or oral antidiabetic agents. Probable cases of new onset diabetes were defined as those participants who reported a diagnosis of diabetes made by a doctor in the follow-up questionnaire and did not have diabetes at baseline. We sent additional questionnaires to participants who reported probable new onset diabetes, requesting confirmation of the diagnosis, the type of diabetes, the date of the diagnosis, whether the diabetes was diagnosed during pregnancy, the highest fasting glucose value that they ever had, whether they had ever undergone an oral glucose tolerance test and its results, their current use of insulin or oral antidiabetic agents, and the incidence of some complications. We also asked them to send us the medical report detailing the diagnosis. A panel of physicians, blinded to the information about dietary habits, used the information provided by participants (additional questionnaires and medical reports) to classify the diagnosis as incident type 2 diabetes or not. The diagnosis criteria for type 2 diabetes were those of the American Diabetes Association (symptoms of diabetes plus casual plasma glucose concentration ≥ 11.1 mmol/l or fasting plasma glucose ≥ 7.0 mmol/l, or two hour postload glucose ≥ 11.1 mmol/l during an oral glucose tolerance test; in the absence of unequivocal hyperglycaemia, confirmation by repeat testing on a different day is requested). Participants were classified as having confirmed incident type 2 diabetes only if they sent us written confirmation that the diagnosis had been received at least six months after they completed the baseline questionnaire and the diagnosis was not made during pregnancy (that is, we excluded cases of gestational diabetes).

Statistical analysis

To estimate incidence rate ratios and their 95% confidence intervals, we ran Poisson regression models with robust standard errors, controlling simultaneously for known risk factors for diabetes: sex, age, years of university education, total energy intake, body mass index, physical activity, hours sitting down a

week, family history of diabetes, baseline hypertension, and smoking. For all analyses, we considered participants with the lowest level of adherence to the Mediterranean diet (score <3) at baseline as the reference category.

When we designed the study, taking into account the planned age and sex distribution of cohort members and the available data about incidence of diabetes in Spain, we expected an overall rate of new onset type 2 diabetes of three cases for 1000 person years of follow-up. Thus, in the follow-up of 12 000 cohort members for an average 4.4 years we would have expected 158 new cases of type 2 diabetes and the statistical power would have been $>90\%$ for the comparison of the two extreme quarters, assuming a relative risk of 0.5 and a two tailed α error of 0.05. The statistical power for the trend test would be higher because both intermediate quarters would also be taken into account.

Because of the surprisingly low incidence of confirmed cases of diabetes, instead of using quarters, we grouped the exposure variable into only three categories. In addition to the reference category (lowest adherence, score <3), we built two other categories: moderate adherence (score 3-6) and high adherence (score 7-9). We did not use any imputation method for the missing data of participants lost to follow-up. Missing data for baseline leisure time physical activity (5.2%), smoking (3.6%), height (1.0%), or weight (0.2%) were imputed by using regression models. We treated participants with missing data for hours spent sitting down a week (20.9%) as a separate category for this variable. We conducted tests of linear trend (likelihood ratio test) across increasing categories of the score by using the score as a continuous variable and adjusting for the potential confounders. All P values were two tailed, and significance was set at $P < 0.05$.

RESULTS

During the follow-up period (median 4.4 years), 103 participants initially free of diabetes (according to their baseline questionnaire) self reported a new diagnosis of diabetes. After we excluded 39 women who developed gestational diabetes, six cases of impaired glucose tolerance mistakenly reported by participants as diabetes, three participants with prevalent type 2 diabetes at baseline who did not report it in the baseline questionnaire, one case of incident type 1 diabetes, and 21 participants who did not send us any additional medical report, we identified 33 cases of new onset confirmed type 2 diabetes among 58 918 person years of follow-up.

Table 1 shows the characteristics of the participants according to their level of adherence to the Mediterranean diet. Though participants with the highest adherence to the diet (score >6) had a higher level of leisure time physical activity, they also exhibited a higher baseline prevalence of most risk factors for diabetes because they were older, had a higher BMI, a higher total energy intake, were more likely to have

Table 1 | Baseline characteristics of 13 380 participants according to adherence (Trichopoulou's score²²) to Mediterranean food pattern at baseline. Figures are means (SD) unless stated otherwise

	Low (score 0-2; n=2253)	Moderate (score 3- 6; n=9604)	High (score 7-9, n=1523)
Sex (% men)	39.7	39.0	44.1
Age (years)	33.8 (9.9)	37.9 (11.7)	42.9 (12.4)
Years of university education	5.0 (1.6)	5.0 (1.6)	5.0 (1.6)
Body mass index (kg/m ²)	23.0 (3.4)	23.4 (3.4)	23.8 (3.4)
Weight (kg)	66.2 (14.4)	66.9 (13.9)	67.8 (13.3)
Mean yearly weight increase (kg)	0.4 (4.1)	0.3 (3.3)	0.4 (4.3)
Physical activity (METs-h/week)	24.1 (23.4)	29.3 (26.2)	36.4 (29.2)
Hours/week sitting down	39.0 (15.5)	37.2 (15.6)	35.6 (15.0)
Family history of diabetes (%)	11.9	13.8	16.7
Former smokers (%)	22.1	28.8	37.5
Current smokers (%)	24.2	24.2	22.7
Hypertension at baseline (%)*	6.2	9.0	13.1
Total energy intake (MJ/day)†	10 (4)	11 (4)	12 (4)
Fatty acids (% of total caloric intake)‡:			
Saturated	15 (3)	12 (3)	10 (2)
Monounsaturated	16 (3)	16 (4)	15 (4)
Dietary intake (g/day)‡:			
Alcohol	4 (9)	7 (11)	10 (11)
Total dietary fibre	18 (8)	29 (15)	41 (18)
Cereal fibre	2 (2)	4 (3)	5 (4)
Olive oil	12 (11)	19 (17)	26 (18)
Vegetables	308 (189)	547 (388)	773 (430)
Fruits	176 (153)	371 (347)	559 (408)
Cereals	77 (69)	113 (89)	150 (98)
Legumes	17 (14)	24 (24)	31 (38)
Fish	63 (41)	103 (79)	140 (75)
Meats/meat products	199 (100)	187 (108)	154 (93)
Fast food‡	27 (25)	22 (28)	17 (34)
Dairy products	338 (266)	226 (233)	109 (133)

METs-h/week=metabolic equivalents of weekly leisure time physical activities.
 *Medical diagnosis required.
 †Estimated values, see text.
 ‡Hamburger, sausages, and pizza.

high blood pressure or a family history of diabetes, and more likely to be former smokers.

When we assessed the risk of diabetes according to levels of adherence to the diet we found a significant inverse association after adjustment for age and sex (table 2). Further multivariate adjustment for total years of university education (three categories), BMI (continuous), family history of diabetes (two categories), hypertension at baseline (two categories), physical activity (three categories), hours sitting down a week (five categories), smoking (three categories), and total energy intake (continuous) did not lead to substantial changes in the estimates, age being the major confounding factor. High adherence to the diet (score >6) was associated with an 83% relative reduction in the risk of developing diabetes (incidence rate ratio 0.17, 95% confidence interval 0.04 to 0.75). When we introduced the score as a continuous variable we found that an increase of two points in the score was associated with an incidence rate ratio of 0.65 (0.44 to 0.95) in the fully adjusted model, indicating that a two

point increase in the score resulted in a 35% relative reduction in the risk of developing diabetes. The multivariate adjusted inverse linear trend was significant ($P=0.04$).

DISCUSSION

This large prospective study shows that a traditional Mediterranean food pattern is associated with a significant reduction in the risk of developing type 2 diabetes.

Interestingly, among participants with the highest adherence to the diet, there was a high prevalence of important risk factors for diabetes, such as older age, higher BMI, family history of diabetes, and personal history of hypertension and a higher proportion of ex-smokers. Therefore, we would have expected a higher incidence of diabetes among these participants. These higher risk participants with better adherence to the diet, however, had a lower risk of diabetes, suggesting that the diet might have a substantial potential for prevention. This finding is consistent with our previous report of an inverse association between a Mediterranean diet and the metabolic syndrome.²⁴ The metabolic syndrome is closely associated with a generalised metabolic disorder of insulin resistance, which is one of the underlying causes of type 2 diabetes. Therefore a high biological plausibility exists to support the causality of our findings. In addition, a previous cohort study of survivors of myocardial infarction also reported that a higher adherence to a Mediterranean diet was associated with a reduction in the risk of type 2 diabetes,¹⁸ despite use of a relatively inaccurate tool for the dietary assessment. The inverse graded dose-response pattern and the significant inverse trend that we observed also support a causal relation.

Diet and disease

Experimental evidence suggesting that a diet similar to the Mediterranean diet and rich in plant based foods might protect against diabetes can also be found in the Indian diabetes prevention programme. That trial promoted a plant based diet (together with increases in physical activity) and found a significant reduction in the incidence of diabetes,²⁸ although there are clear differences between the Mediterranean and Indian food patterns.

There is an analogy between coronary heart disease and diabetes because patients with type 2 diabetes and no coronary heart disease have a risk of coronary heart disease similar to patients without diabetes but with prior coronary heart disease.²⁹ There is evidence that a Mediterranean diet protects against coronary heart disease, and the analogy between coronary heart disease and diabetes suggests that this diet might also prevent diabetes. It has been shown to provide protection against coronary mortality^{8,9,30,31} and the incidence of non-fatal myocardial infarction.³² In addition, both cohort studies³³⁻³⁵ and randomised trials³⁶ have found that adherence to a Mediterranean

diet protects against mortality in patients who already have established coronary heart disease.

The potential mechanisms explaining the protective effect of a Mediterranean diet on diabetes have been reviewed elsewhere.³⁷⁻⁴⁰ Two trials have shown that virgin olive oil protects against insulin resistance and the metabolic syndrome.^{6,7} Indexes of insulin resistance were significantly improved among participants allocated to a Mediterranean diet rich in virgin olive oil. Olive oil is rich in monounsaturated fatty acids, and a diet rich in monounsaturated fatty acids is beneficial among those with diabetes and might lead to improved insulin sensitivity and better lipid profiles than diets rich in carbohydrate.^{13,15,39} People allocated to a Mediterranean diet secrete more glucagon-like peptide-1,¹⁶ a finding also seen in animals.¹⁴ The non-fat minor components of virgin olive oil also exert a beneficial effect on pro-inflammatory cytokines.⁴⁰⁻⁴²

Apart from olive oil, adherence to an overall Mediterranean-type food pattern is related to lower plasma concentrations of inflammatory markers and markers of endothelial dysfunction.^{43,44} These biomarkers are predictive of the future occurrence of type 2 diabetes.^{45,46} In addition, a large cross sectional study nested in the nurses' health study found that increased adherence to a Mediterranean diet was associated with higher levels of adiponectin,⁴⁷ and higher levels of adiponectin are associated with a reduced risk of diabetes.

Limitations

The number of new cases of diabetes was small, despite the follow-up of several thousand people for over four years. This small number of incident cases is a major drawback and can compromise the statistical power of our study. Nevertheless, our participants had high absolute levels of consumption of the typical food items of the Mediterranean diet, even among those participants classified as poorly compliant (score <3). For example, among those in the lowest category of adherence to the diet, the estimated mean daily absolute consumption (g/day) of olive oil (12), vegetables (308), fruits (176), cereals (77), and legumes (17) can be considered as healthy for the standards of nutritional epidemiological studies. This high

consumption of plant based foods in our cohort could be because our participants were from a Mediterranean country and were highly educated and health conscious. The small number of new cases observed in this study should therefore not be surprising. If the Mediterranean diet is actually protecting against diabetes, we would expect a low incidence in a young cohort (mean age is 37.8 years) with these characteristics. The low number of observed cases precluded assessment of the specific role of single dietary factors because we would have needed multiple adjustments for other dietary factors (in addition to the confounders already included in the model) and the statistical power to adjust for so many factors would have been low. The Mediterranean food pattern has the potential to minimise confounding by including nutritional confounders in the score and capturing effect modification among the nutritional variables.⁵

Diabetes might have been under-reported in our participants, despite their high educational level and easy access to medical care—half of them are health professionals. The proportions of participants aged over 65 years across increasing categories of adherence to the diet were 0.6% (score <3), 1.7% (3-6), and 3.5% (>6). Therefore, older participants were more compliant with the diet. If selective under-reporting of diabetes was present among the older participants, this would provide an alternative non-causal explanation for our findings. We acknowledge this possibility, but the proportion of selective underdiagnosis of diabetes among older people is likely to be lower as they receive closer medical attention and routine medical care in Spain includes assessment of fasting glucose.

All our participants are university graduates and the generalisability of our findings to other groups with less education should be assumed only on biological grounds but not at all on “representativeness” of the general population. Also, the building of the diet score was based on sample specific median cut-off points, and our participants had high absolute levels of consumption of favourable foods and low absolute intakes of detrimental foods. Therefore, it will be difficult to compare our results with those of non-Mediterranean countries where levels of consumption of favourable foods in the general population are much lower. This is an additional limitation for external validity, but we selected the score proposed by Trichopoulos et al for the sake of comparability with previous studies.^{8,22-24,30-34}

A potential caveat might be the quality of our nutritional assessment. Food frequency questionnaires are known to contain a certain degree of measurement error, which might affect results that depend on such questionnaires to assess diet and risk of chronic disease. Total energy intake was included as a covariate in the model to achieve the equivalent of an isocaloric diet and to reduce measurement error in the score. Measurement error, however, would probably have introduced non-differential misclassification, and the

Table 2 | Incidence and relative risk of type 2 diabetes (confirmed cases) during follow-up according to adherence (Trichopoulos's score²²) to Mediterranean food pattern at baseline

	No in group	Unadjusted cumulative incidence of type 2 diabetes (%)	Incidence rate ratio* adjusted for age and sex (95% CI)	Multivariate adjusted incidence rate ratio (95% CI)†
Low (score 0-2)	2253	0.40	1 (reference)	1 (reference)
Moderate (score 3-6)	9604	0.23	0.41 (0.19 to 0.87)	0.40 (0.18 to 0.90)
High (score 7-9)	1523	0.13	0.17 (0.04 to 0.75)	0.17 (0.04 to 0.72)

*Poisson regression model with robust standard errors.

†Adjusted for sex, age, years of university education (three categories), body mass index (continuous), family history of diabetes (two categories), hypertension at baseline (two categories), physical activity (three categories), hours/week sitting down (five categories), smoking (three categories), total energy intake (continuous). P=0.04 for trend from likelihood ratio test when Trichopoulos's score was introduced as continuous variable in fully multivariate adjusted model.

WHAT IS ALREADY KNOWN ON THIS TOPIC

A Mediterranean diet protects against coronary mortality and is inversely associated with metabolic syndrome and inflammatory markers

The diet was found to be inversely associated with the incidence of diabetes among patients who survived myocardial infarction

WHAT THIS STUDY ADDS

Higher adherence to a Mediterranean diet rich in olive oil, plant based foods (fruits, vegetables, and legumes), and fibre but low in meats was inversely associated with incidence of type 2 diabetes among initially healthy participants

implications for the results of this error would have been to bias the estimates towards the null.

A potential limitation, inherent to every observational design, is the possibility of residual confounding by unmeasured or unrecorded factors. Our major confounders, however, were sex and age. Additional adjustment for other factors made only negligible changes in the estimates, suggesting that residual confounding is unlikely.

Conclusion

Our prospective cohort study suggests that substantial protection against diabetes can be obtained with the traditional Mediterranean diet, rich in olive oil, vegetables, fruits, nuts, cereals, legumes, and fish but relatively low in meat and dairy products. The limited number of cases of diabetes and the possibility of under-reporting, however, requires that further larger cohorts and trials are needed to confirm our findings.

We are indebted to the participants of the SUN study for their continued cooperation and participation. We also thank others members of the SUN study group including: M Seguí-Gomez, J de Irala, J A Martínez, A Martí, M Serrano-Martínez, F Guillen-Grima, E Toledo, R M Pajares, University of Navarra; M Delgado-Rodríguez, University of Jaen; J Llorca, University of Cantabria; E Smyth, University of Santiago de Compostela; A Sanchez-Villegas, University of Las Palmas; I Marques-Lopes, University of Saragossa; A Alonso, University of Minnesota; and C N Lopez, Harvard University. We thank the members of the Department of Nutrition of the Harvard School of Public Health (A Ascherio, F B Hu, W C Willett) who helped us to design the SUN study.

Contributors: MAM-G and MB-R were responsible for study design, recruitment of participants, and statistical analyses and wrote the first draft of the manuscript; CF-A, ZV, SB, and AT were involved in data collection, technical support, and data management; JMN-C, FJB-G, JJB, and MAM-G applied the diagnosis criteria and comprised the medical panel for the adjudication of events. MAM-G obtained funding and is guarantor. All authors provided suggestions during the different phases of the preparation of the manuscript and approved the final version submitted for publication.

Funding: Spanish Ministry of Health (Instituto de Salud Carlos III, Fondo de Investigaciones Sanitarias projects PI030678, PI040233, PI042241, PI050514, PI050976, PI070240; RD 06/0045) and the Navarra Regional Government (Department of Health, PI41/2005).

Competing interests: None declared.

Ethical approval: Institutional Review Board of the University of Navarra.

Provenance and peer review: Not commissioned; externally peer reviewed.

- Hossain P, Kavar B, El Nahas M. Obesity and diabetes in the developing world—a growing challenge. *N Engl J Med* 2007;356:213-5.
- International Diabetes Federation. *Diabetes atlas*. 3rd ed. Brussels: International Diabetes Federation, 2006.

- Alberti KGMM, Zimmet P, Straw J. International diabetes federation: a consensus on type 2 diabetes prevention. *Diabet Med* 2007;24:451-63.
- Hu FB, Manson JE, Stampfer MJ, Colditz G, Liu S, Solomon CG, Willett WC. Diet, lifestyle, and the risk of type 2 diabetes mellitus in women. *N Engl J Med* 2001;345:790-7.
- Martínez-González MA, Sánchez-Villegas A. The emerging role of Mediterranean diets in cardiovascular epidemiology: MUFA, olive oil, red wine or the whole pattern? *Eur J Epidemiol* 2004;19:9-13.
- Estruch R, Martínez-González MA, Corella D, Salas-Salvadó J, Ruiz-Gutiérrez V, Covas MI, et al. Effects of a Mediterranean-style diet on cardiovascular risk factors: a randomized trial. *Ann Intern Med* 2006;145:1-11.
- Esposito K, Marfella R, Ciotola M, Di Palo C, Guigliano F, Giugliano G, et al. Effect of a Mediterranean-style diet on endothelial dysfunction and markers of vascular inflammation in the metabolic syndrome. *JAMA* 2004;292:1440-6.
- Trichopoulos A, Costacou T, Bamia C, Trichopoulos D. Adherence to a Mediterranean diet and survival in a Greek population. *N Engl J Med* 2003;348:2599-608.
- Knoops KT, de Groot LC, Kromhout D, Perrin AE, Moreiras-Varela O, Menotti A, et al. Mediterranean diet, lifestyle factors, and 10-year mortality in elderly European men and women. The HALE project. *JAMA* 2004;292:1433-9.
- Salmeron J, Hu FB, Manson JE, Stampfer MJ, Colditz GA, Rimm EB, et al. Dietary fat intake and the risk of type 2 diabetes in women. *Am J Clin Nutr* 2001;73:1019-26.
- Schulze MB, Schulz M, Heidemann C, Schienkiewitz A, Hoffmann K, Boeing H. Fiber and magnesium intake and incidence of type 2 diabetes: a prospective study and meta-analysis. *Arch Intern Med* 2007;167:956-65.
- Perez-Jimenez F, Alvarez de Cienfuegos G, Badimon L, Barja G, Battino M, Blanco A, et al. International conference on the healthy effect of virgin olive oil. *Eur J Clin Invest* 2005;35:421-4.
- Garg A. High monounsaturated-fat diets for patients with diabetes mellitus: a meta-analysis. *Am J Clin Nutr* 1998;67(suppl):577-82S.
- Rocca AS, LaGreca J, Kalitsky J, Brubaker PL. Monounsaturated fatty acid diets improve glycemic tolerance through increased secretion of glucagon-like peptide-1. *Endocrinology* 2001;142:1148-55.
- Ros E. Dietary cis-monounsaturated fatty acids and metabolic control in type 2 diabetes. *Am J Clin Nutr* 2003;78(suppl):617-25S.
- Paniagua JA, de la Sacristana AG, Sánchez E, Romero I, Vidal-Puig A, Berral F, et al. MUFA-rich diet improves postprandial glucose, lipid and GLP-1 responses in insulin-resistant subjects. *J Am Coll Nutr* 2007;26:434-44.
- Willett WC. The Mediterranean diet: science and practice. *Public Health Nutr* 2006;9:105-10.
- Mozaffarian D, Marfisi R, Levantesi G, Silletta MG, Tavazzi L, Tognoni G, et al. Incidence of new-onset diabetes and impaired fasting glucose in patients with recent myocardial infarction and the effect of clinical and lifestyle risk factors. *Lancet* 2007;370:667-75.
- Martínez-González MA, Sánchez-Villegas A, de Irala J, Martí A, Martínez JA. Mediterranean diet and stroke: objectives and design of the SUN project. *Nutr Neurosci* 2002;5:65-73.
- Seguí-Gomez M, de la Fuente C, Vazquez Z, de Irala J, Martínez-González MA. Cohort profile: the "Seguimiento Universidad de Navarra" (SUN) study. *Int J Epidemiol* 2006;35:1417-22.
- Martin-Moreno JM, Boyle P, Gorgojo L, Maisonneuve P, Fernandez-Rodríguez JC, Salvini S, et al. Development and validation of a food frequency questionnaire in Spain. *Int J Epidemiol* 1993;22:512-9.
- Trichopoulos A, Kouris-Blazos A, Wahlquist M, Gnardellis D, Lagiou P, Polychronopoulos E, et al. Diet and overall survival in elderly people. *BMJ* 1995;311:1457-60.
- Bach A, Serra-Majem L, Carrasco JL, Roman B, Ngo J, Bertomeu I, et al. The use of indexes evaluating the adherence to the Mediterranean diet in epidemiological studies: a review. *Public Health Nutr* 2006;9:132-46.
- Tortosa A, Bes-Rastrollo M, Sanchez-Villegas A, Basterra-Gortari FJ, Nuñez-Córdoba JM, Martínez-González MA. Mediterranean diet inversely associated with the incidence of metabolic syndrome: the SUN prospective cohort. *Diabetes Care* 2007;30:2957-9.
- Bes-Rastrollo M, Perez Valdivieso JR, Sanchez-Villegas A, Alonso A, Martínez-González MA. Validación del peso e índice de masa corporal auto-declarados de los participantes de una cohorte de graduados universitarios. *Rev Esp Obes* 2005;3:183-9 [in Spanish].
- Martínez-González MA, Lopez-Fontana C, Varo JJ, Sánchez-Villegas A, Martínez JA. Validation of the Spanish version of the physical activity questionnaire used in the nurses' health study and health professionals' follow-up study. *Public Health Nutr* 2005;8:920-7.
- Ainsworth BE, Haskell WL, Whitt MC, Irwin ML, Swartz AM, Strath SJ, et al. Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc* 2000;32:498-504.
- Ramachandran A, Snehalatha C, Mary S, Mukesh B, Bhaskar AD, Vijay V. The Indian diabetes prevention programme shows that

- lifestyle modification and metformin prevent type 2 diabetes in Asian Indian subjects with impaired glucose tolerance (IDPP-1). *Diabetologia* 2006;49:289-97.
- 29 González-Clemente JM, Palma S, Arroyo J, Vilardell C, Caixàs A, Giménez-Palop O, et al. Is diabetes mellitus a coronary heart disease equivalent? Results of a meta-analysis of prospective studies. *Rev Esp Cardiol* 2007;60:1167-76.
 - 30 Trichopoulou A, Orfanos P, Norat T, Bueno-de-Mesquita B, Ocké MC, Peeters PH, et al. Modified Mediterranean diet and survival: EPIC-elderly prospective cohort study. *BMJ* 2005;330:991.
 - 31 Mitrou PN, Kipnis V, Thiébaud AC, Reedy J, Subar AF, Wirfält E, et al. Mediterranean dietary pattern and prediction of all-cause mortality in a US population: results from the NIH-AARP diet and health study. *Arch Intern Med* 2007;167:2461-8.
 - 32 Martínez-González MA, Fernández-Jame E, Serrano-Martínez M, Martí A, Martínez JA, Martín-Moreno JM. Mediterranean diet and reduction in the risk of a first acute myocardial infarction: an operational healthy dietary score. *Eur J Nutr* 2002;41:153-60.
 - 33 Trichopoulou A, Bamia C, Trichopoulos D. Mediterranean diet and survival among patients with coronary heart disease in Greece. *Arch Intern Med* 2005;165:929-35.
 - 34 Trichopoulou A, Bamia C, Norat T, Overvad K, Schmidt EB, Tjønneland A, et al. Modified Mediterranean diet and survival after myocardial infarction: the EPIC-Elderly study. *Eur J Epidemiol* 2007;22:871-81.
 - 35 Iestra J, Knoop K, Kromhout D, de Groot L, Grobbee D, van Staveren W. Lifestyle, Mediterranean diet and survival in European post-myocardial infarction patients. *Eur J Cardiovasc Prev Rehabil* 2006;13:894-900.
 - 36 De Lorgeril M, Salen P, Martin JL, Monjaud I, Delaye J, Mamelle N. Mediterranean diet, traditional risk factors, and the rate of cardiovascular complications after myocardial infarction: final report of the Lyon diet heart study. *Circulation* 1999;99:779-85.
 - 37 Schröder H. Protective mechanisms of the Mediterranean diet in obesity and type 2 diabetes. *J Nutr Biochem* 2007;18:149-60.
 - 38 Biesalski HK. Diabetes preventive components in the Mediterranean diet. *Eur J Nutr* 2004;43(suppl 1):126-30.
 - 39 Pérez-Jiménez F, López-Miranda J, Mata P. Protective effect of dietary monounsaturated fat on arteriosclerosis: beyond cholesterol. *Atherosclerosis* 2002;163:385-98.
 - 40 Perez-Martinez P, Lopez-Miranda J, Blanco-Colio L, Bellido C, Jimenez Y, Moreno JA, et al. The chronic intake of a Mediterranean diet enriched in virgin olive oil, decreases nuclear transcription factor kappaB activation in peripheral blood mononuclear cells from healthy men. *Atherosclerosis* 2007;194:e141-6.
 - 41 Salas-Salvado J, Garcia-Arellano A, Estruch R, Marquez-Sandoval F, Corella D, Fiol M, et al. Components of the mediterranean-type food pattern and serum inflammatory markers among patients at high risk for cardiovascular disease. *Eur J Clin Nutr* 2007 Apr 18 [Epub ahead of print].
 - 42 Fitó M, Cladellas M, de la Torre R, Martí J, Muñoz D, Schröder H, et al. Anti-inflammatory effect of virgin olive oil in stable coronary disease patients: a randomized, crossover, controlled trial. *Eur J Clin Nutr* 2007;62:570-4.
 - 43 Fung TT, McCullough ML, Newby PK, Manson JE, Meigs JB, Rifai N, et al. Diet-quality scores and plasma concentrations of markers of inflammation and endothelial dysfunction. *Am J Clin Nutr* 2005;82:163-73.
 - 44 Lopez-Garcia E, Schulze MB, Fung TT, Meigs JB, Rifai N, Manson JE, et al. Major dietary patterns are related to plasma concentrations of markers of inflammation and endothelial dysfunction. *Am J Clin Nutr* 2004;80:1029-35.
 - 45 Meigs JB, Hu FB, Rifai N, Manson JE. Biomarkers of endothelial dysfunction and risk of type 2 diabetes mellitus. *JAMA* 2004;291:1978-86.
 - 46 Pradhan AD, Manson JE, Rifai N, Buring JE, Ridker PM. C-reactive protein, interleukin-6, and risk of developing type-2 diabetes. *JAMA* 2001;286:327-34.
 - 47 Mantzoros CS, Williams CJ, Manson JE, Meigs JB, Hu FB. Adherence to the Mediterranean dietary pattern is positively associated with plasma adiponectin concentrations in diabetic women. *Am J Clin Nutr* 2006;84:328-35.

Accepted: 5 April 2008