

gross atheroma is rare in patients with chronic alcoholism, a condition commonly associated with severe protein deficiency. It would seem, therefore, that the degree of protein consumption is yet another factor which will require investigation during the course of further research designed to establish the aetiology of coronary artery disease.—I am, etc.,

Medical Research Laboratories,
Charing Cross Hospital,
London W.C.2.

M. A. PEYMAN.

REFERENCES

- 1 Mills, C. A., and Necheles, H., *Proc. Soc. exp. Biol. (N.Y.)*, 1927, **25**, 195.
- 2 ——— *Chin. J. Physiol.*, 1928, **2**, 19.
- 3 Necheles, H., and Mills, C. A., *ibid.*, 1928, **2**, 25.
- 4 Mills, C. A., Necheles, H., and Mao-Keng Chu, *ibid.*, 1928, **2**, 219.
- 5 Ahrens, E. H., jun., *Amer. J. Med.*, 1957, **23**, 928.

Post-mortem Serum Cholesterol

SIR,—Dr. J. N. Glanville's data (December 24, 1960, p. 1852) are a useful contribution in a field that has been little explored. In fact, its contours are extremely complex. In 1936 Lande and Sperry,¹ at Schonheimer's suggestion, made successful cholesterol analyses on cadaver sera, and Faber² considered that any differences encountered in serial estimations were within the limits of experimental error. More recently Paterson and Dyer³ have compared ante- and post-mortem blood-cholesterol levels in a closed population and have concluded that they were comparable provided death was sudden.

I have been collecting data from cadaver sera for some years now, and I believe that all these authors have in fact missed a small autolytic change. Many serum constituents, particularly enzymes, increase very rapidly in the serum for a few hours after death, then slowly for 48 hours before falling off again. In very old bodies there is often a further final rise.⁴ These changes are much less striking in non-protein substances and in them the initial sharp rise is not seen. The effect of this rather complex change is that in most cases in which analysis is likely to be contemplated—that is, those of simpler constituents on sera obtained within two days of death—a direct linear increase with time can be expected. My data are based on analyses now of nearly 1,000 cadaver sera, and the curves have been constructed by simply relating time after death to result of analysis. The values obtained in several hundred cholesterol analyses give means in successive 12-hour periods of 195, 207, 213, 232, followed by a fall to 200 in the 49- to 60-hour period. There is thus an hourly increment of about 0.5 mg. per 100 ml. The conventional Bloor method was used.

Such a correction does indeed give discrepancies well within the experimental limits but quite large enough to have a 20:1 significance in large groups with differing mean times after death. I have, however, quite insufficient data on pairs of ante- and post-mortem analyses to plot the point corresponding to the 12 hours preceding death. If Dr. Glanville would do this, the information would be most valuable and would, I think, be unique.—I am, etc.,

East Ham Memorial Hospital,
London E.7.

J. B. ENTICKNAP.

REFERENCES

- 1 Sperry, W. M., in *Chemistry of Lipids as Related to Atherosclerosis*, 1958, ed. I. H. Page. Thomas, Springfield, Illinois.
- 2 Faber, M., *Acta med. scand.*, 1946, **125**, 418.
- 3 Paterson, J. C., and Dyer, L., *Circulation*, 1959, **20**, 989.
- 4 Enticknap, J. B., *J. forensic Med.*, 1960, **7**, 135.

Antibody to Penicillin

SIR,—We read with great interest your leading article on this matter (August 6, p. 444), as it seems you were not aware that the findings of Ley *et al.*¹ have been confirmed and extended by us in a Spanish paper with an English summary.² The work of Watson *et al.*³ was published after this. We found antibodies to penicillin-treated red cells in the sera of four out of nine patients with penicillin shock, and Epp⁵ reported similar results. However, in such haemagglutinating sera we did not detect precipitins, and tests of passive transfer of sensitivity with these sera were negative. Thus, your conclusion that "the basis of sensitivity to penicillin must be sought elsewhere" and not in the haemagglutinin has been anticipated. There is yet another point. What seems to be antibody to penicillin may act in fact on an accompanying substance.—We are, etc.,

Faculdade de Medicina,
São Paulo, Brazil.

F. OTTENSOOSER.
E. MENDES.

REFERENCES

- 1 Ley, A. B., *et al.*, *Science*, 1958, **127**, 1118.
- 2 Ottensooser, F., and Mendes, E., *Sangre*, 1958, **3**, 417.
- 3 Watson, K. C., Joubert, S. M., and Bennett, M. A. E., *Nature (Lond.)*, 1959, **183**, 468.
- 4 ——— *J. Immunology*, 1960, **3**, 1.
- 5 Epp, M., *Science*, 1959, **130**, 1472.

Arteritis

SIR,—In connexion with the article by Drs. J. W. Paulley and J. P. Hughes (November 26, p. 1562), the possibility of an immunological phenomenon being concerned with the causation of temporal arteritis was mentioned in correspondence by Dr. G. D. Kersley (December 31, p. 1953). Together with my laboratory colleagues I have been working on these lines for the past two years. Since much of the inflammatory activity appears histologically to be centred around damaged elastic tissue, we have worked principally with elastin and its soluble derivatives. Experimental work, to be published elsewhere, has not so far afforded any evidence for the hypothesis that temporal arteritis is a disease associated with an auto-immune reaction involving elastin or its soluble derivatives.—I am, etc.,

The Royal Infirmary,
Sheffield.

G. V. BALMFORTH.

Pathology of Chronic Bronchitis

SIR,—Your annotation (December 31, 1960, p. 1946) on the pathology of chronic bronchitis raises several interesting points. The histological changes in chronic bronchitis consist not only in the increase in size and number of deep mucous glands but in the more or less complete loss of the normal serous glands and of the demilunes of serous cells normally found in some mucous glands. Whether the serous cells atrophy or are converted to mucus-secreting cells is unknown.

In a recent study of bronchial biopsies from patients with chronic bronchitis and asthma, Dr. Michaels and I found¹ that the mucoid change was almost entirely confined to patients with chronic bronchitis—that is, those who had had persistent or recurrent cough and sputum for at least two years. Cases of pure asthma—that is, with no coincident or superimposed chronic bronchitis—with few exceptions had the normal predominantly serous pattern of deep glands even after many years. It is interesting that in the series of deaths in status asthmaticus recorded by Houston, De Navasquez, and Trounce² the bronchi were:

completely obstructed by mucus, yet the deep glands were histologically normal.

Within the bronchitic group of biopsies, the degree of mucous change was not related to the use of tobacco or the presence of infection but depended on the duration of the disease. Anything more than slight change was only found in bronchitics with at least ten years' history, while the most severe changes occurred in patients with over twenty years' history.

Goblet cells are more easily studied in biopsy that in necropsy specimens. In the series referred to above, goblet cell hyperplasia was equally common in cases of bronchitis, asthma, and bronchitis with asthma. It bore no relation to the degree of mucoid change in the deep glands but was significantly more frequent where there was infection with *H. influenzae*.—I am, etc.,

St. Mary's Hospital Medical School, A. A. GLYNN.
London W.2.

REFERENCES

- ¹ Glynn, A. A., and Michaels, L., *Thorax*, 1960, **15**, 142.
² Houston, J. C., De Navasquez, S., and Trounce, J. R., *ibid.*, 1953, **8**, 207.

Megaloblastic Madness

SIR,—We would like to congratulate Dr. A. D. M. Smith (December 24, 1960, p. 1840) on his important paper. Dr. Smith refers to the results of our study of 14 male patients with tobacco amblyopia seen at the Bristol Eye Hospital which showed that an essential feature was a low vitamin-B₁₂ concentration in the serum.¹ However, his statement that "unfortunately bone-marrow studies were not performed in this series" is not strictly accurate. It was, in fact, decided that biopsy of the sternal marrow should be undertaken in every case of tobacco amblyopia with histamine-fast achlorhydria and evidence of neurological involvement, irrespective of whether or not anaemia was present. Using this method of selection we found two patients had a megaloblastic bone-marrow and one patient a normoblastic marrow.

We certainly did not expect that low or very low serum vitamin-B₁₂ levels would also be found in patients with free acid on gastric analysis, and in this clinical trial we did not feel justified in withholding treatment with cyanocobalamin till the results of microbiological assay were available in view of the frequency of severe and progressive deterioration of vision. Now, being wise after the event, we entirely agree with Dr. Smith that it would have been of considerable interest to know if biopsy of the marrow showed normoblastic or megaloblastic erythropoiesis in our remaining cases.

In this connexion, we would like to refer briefly to one of our cases included in our series. The presentation of tobacco amblyopia and mental changes, consisting of acute depression, irritability, lack of concentration, and failing memory, in a non-anaemic man, aged 52, is of particular interest. Although free acid was present in the gastric juice, the serum vitamin-B₁₂ level was exceptionally low at 15 µg./ml. We therefore suggested that in any case of unexplained peripheral neuropathy, myelopathy, or encephalopathy the concentration of vitamin B₁₂ in the serum should be measured, *even if histamine-fast achlorhydria is not present*. The interesting findings of Dr. Smith add further support to this view.—We are, etc.,

Princess Margaret Hospital, A. G. FREEMAN.
Swindon, Wiltshire.
Institute of Ophthalmology, J. M. HEATON.
London W.C.1.

REFERENCE

- ¹ Heaton, J. M., McCormick, A. J. A., and Freeman, A. G., *Lancet*, 1958, **2**, 286.

SIR,—May I plead that your contributors cease using the word "madness" when describing a new, or apparently new, diagnostic category? Apart from the fact that this word carries prejudices in both medical and lay circles, it can mislead: madness implies a pretty severe form of mental disorder whereas, as Dr. A. D. M. Smith points out at the beginning of his excellent paper (December 24, 1960, p. 1840), the mental symptoms in "megaloblastic madness" vary from the mildest to the most severe; and a mild neurotic reaction or mood swing is not "madness."

Nearly thirty years ago, when I reported a relevant case,¹ I was content to use the phrase "mental disease." It would be interesting to know whether there is a valid medical or social reason for the increasing use of the word "madness." If one of your readers can supply this, I shall resign myself to a series of articles, not only on "megaloblastic madness," but also on "myxoedematous madness," "diabetic madness," "syphilitic madness," "influenzal madness," and so forth.—I am, etc.,

Park Prewett Hospital,
Basingstoke.

I. ATKIN.

REFERENCE

- ¹ Atkin, I., *Lancet*, 1932, **2**, 569.

Familial Pseudocholinesterase Deficiency

SIR,—Dr. F. D. Adrianvala (December 31, 1960, p. 1955) finds that none of the members of the family of a case of suxamethonium apnoea with low pseudocholinesterase was similarly affected. He is correct in concluding that "these results lend no support to the suggestion of any genetic behaviour on the part of pseudocholinesterase deficiency." We can furnish Dr. Adrianvala with hundreds of examples of patients with low pseudocholinesterase level where there is no evidence of inheritance, but equally also with dozens where family study has borne out a "genetic behaviour on the part of pseudocholinesterase deficiency." When the genetic deficiency was first postulated¹ it was made clear that pseudocholinesterase is low in the majority of cases because of liver disease, extreme malnutrition, etc., but that there had been instances where the clinical findings did not correlate with the laboratory results. When the families of such individuals with low pseudocholinesterase (usually discovered because of their sensitivity to suxamethonium) were investigated the picture of a familial deficiency emerged.²⁻⁵

Fortunately it is now possible to determine *in vitro* whether a low pseudocholinesterase level is of the inherited or the acquired type. Kalow and his collaborators found in Canada⁶⁻⁸ that, in the familial type, normal enzyme is replaced by another which is not only less active but also shows a diminished sensitivity to inhibition by dibucaine (cinchocaine). The percentage inhibition is constant for any one individual and is independent of variations in enzyme level. Kalow has devised a test system in which plasmas from normal homozygotes lose 80% of their activity in the presence of dibucaine, and those of abnormal homozygotes are inhibited to only about 16%. Furthermore, whereas the range of pseudocholinesterase levels of heterozygotes overlaps at both ends with those found in abnormal and normal homozygotes, there is no such overlap in the range of pseudocholinesterase inhibition by dibucaine, which for heterozygotes lies midway between the 80% and 16% inhibition of normal and abnormal homozygotes respectively at about 60%.

In collaboration with Professor H. Harris and Dr. Mary Whittaker we have recently compared the pseudocholinesterase activity and the sensitivity towards dibucaine in