

SERUM VITAMIN B₁₂ IN VEGETARIANS

BY

D. K. BANERJEE, M.Sc.
Research Assistant

AND

J. B. CHATTERJEA, M.D., F.N.I.
Professor of Haematology

From the Haematological Unit, Indian Council of Medical Research, and the Department of Haematology, School of Tropical Medicine, Calcutta, India

The vegetables normally consumed by man in general have a negligible amount of vitamin B₁₂ (Lewis *et al.*, 1949; Peeler *et al.*, 1951). The vitamin B₁₂ content of milk is much lower than that of liver, meat, fish, and egg. Thus an average vegetarian diet may not have enough vitamin B₁₂ to ensure optimum human nutrition. The vitamin B₁₂ content of serum in vegetarians has been reported to be low (Wokes *et al.*, 1955; Dhopeswarkar *et al.*, 1956; Chatterjea *et al.*, 1959). In nutritional macrocytic anaemia (N.M.A.) a low concentration of serum vitamin B₁₂ has been found in approximately 50% of cases (Das Gupta *et al.*, 1955; Chatterjea, 1958). The main cause of vitamin-B₁₂ deficiency in N.M.A. is held to be inadequacy of the diet (Das Gupta *et al.*, 1953; Chatterjea, 1958); the possibility of an intestinal malabsorptive factor has also been raised (Baker, 1958). A significant proportion of the population of India may be categorized as vegetarian, due to social custom, to religious tenets, or to poor economic status. The average red-cell and haemoglobin levels of vegetarian Indians cannot, however, be regarded as significantly different from those of non-vegetarians in India or in other parts of the world (Sokhey *et al.*, 1937, 1938; Das Gupta, 1949).

In view of the above considerations, it was thought worth while to investigate the serum vitamin B₁₂ in two groups of subjects—non-vegetarian and vegetarian—as seen in the general population and in those with haematological disorders.

Material and Method

The present communication deals with observations on 232 subjects, consisting of 61 normal persons, 67 with tropical eosinophilia, 39 with N.M.A., 35 with aplastic anaemia, and 30 with chronic myeloid leukaemia. The distribution of non-vegetarians and vegetarians is shown in the Tables.

Subjects belonging to the vegetarian group were taking rice, wheat, vegetables, pulses, and milk, but no fish, meat, or egg. Each one of them had been taking the vegetarian diet for a period varying from 5 to 25 years. The daily consumption of milk was in general small, varying from 6 to 12 oz. (170 to 340 ml.) per head.

The serum vitamin B₁₂ was estimated by the method of Ross (1952), using *Euglena gracilis* var. *bacillaris* as the test organism. The total vitamin B₁₂ content was expressed in micromicrograms per millilitre.

Results


The concentration of total serum vitamin B₁₂ in each individual subject belonging to the two groups is represented in the Chart.

The mean value and range of variation of total vitamin B₁₂ of serum in the different subjects belonging to the two groups are recorded in Tables I and II.

From the Chart it is seen that six of the normal subjects belonging to the vegetarian group had serum vitamin B₁₂ values lower than the lowest value in the non-vegetarian group—that is, 98 μg./ml. In tropical eosinophilia a low level of serum vitamin B₁₂ was found in 8 out of 16 cases. In nutritional macrocytic anaemia,

TABLE I.—Non-vegetarian

Subjects	No. of Cases	Total Vitamin B ₁₂ in μg./ml. of Serum	
		Range	Mean ± S.E.
Normal	51	98 to 940	281.9 ± 22.7
Tropical eosinophilia	51	100 to 670	254.8 ± 16.1
Nutritional macrocytic anaemia	26	0 to 330	136.9 ± 20.7
Aplastic anaemia	33	144 to 1,800	621.9 ± 76.6
Chronic myeloid leukaemia	25	960 to 7,200	2,754.0 ± 317.2



Scattergram showing distribution of serum vitamin B₁₂ in vegetarians and non-vegetarians.

TABLE II.—*Vegetarian*

	No. of Cases	Total Vitamin B ₁₂ in $\mu\text{g.}/\text{ml.}$ of Serum		Cases with $<98 \mu\text{g.}/\text{ml.}$ Vitamin B ₁₂
		Range	Mean \pm S.E.	
Normal ..	10	36 to 290	92.3 \pm 24.3	60%
Tropical eosinophilia ..	16	20 to 490	139.5 \pm 30.6	50%
Nutritional macrocytic anaemia ..	13	20 to 196	47.5 \pm 12.7	92.3%
Aplastic anaemia ..	2	40 and 80	—	Low in both the cases
Chronic myeloid leukaemia ..	5	1,190 to 3,100	2,219 \pm 318.8	0

a low serum vitamin B₁₂ value was found in all the vegetarians except one. Of the aplastic anaemia patients only two were vegetarians, and in both of them the serum vitamin B₁₂ was low, being 40 and 80 $\mu\text{g.}/\text{ml.}$ respectively. No differential pattern was seen in chronic myeloid leukaemia, all the subjects showing elevated level.

A comparative study of Tables I and II shows that in the vegetarians the mean values were significantly lower than those in non-vegetarians in all groups except chronic myeloid leukaemia. In the vegetarians the percentage of cases having a serum vitamin B₁₂ level lower than 98 $\mu\text{g.}/\text{ml.}$ was 92.3% in N.M.A., 60% in normals, and 50% in tropical eosinophilia.

Discussion

Results of the present investigation indicate that the vegetarians have a serum vitamin B₁₂ concentration which is significantly lower than that in the non-vegetarians. On the basis of clinical evaluation and analysis of dietary habits, vitamin B₁₂ deficiency in the vegetarians appears to be dietary in origin. In a parallel study with ⁶⁰Co-labelled vitamin B₁₂ for investigating the pathogenesis of vitamin B₁₂ deficiency in N.M.A., dietary inadequacy appeared to be more important than the factor of intestinal malabsorption (Banerjee *et al.*, 1959).

In normal subjects and the cases of tropical eosinophilia there was no significant anaemia or macrocytosis. The blood picture in vegetarians and non-vegetarians was more or less similar, though their serum vitamin B₁₂ levels were different. Some of the vegetarians followed for one or two years did not show any decline of haemoglobin or R.B.C. level in spite of the serum vitamin B₁₂ level continuing to be low. These findings are not in complete agreement with those of Wokes *et al.* (1955), who found evidence of anaemia and macrocytosis in a significant proportion of vegans investigated by them. It must, however, be remembered that, owing to the exclusion of milk and milk products from their food, the average diet of a vegan was relatively more deficient in vitamin B₁₂ than that of a normal Indian vegetarian. Maintenance of normal haemoglobin and R.B.C. values with a serum vitamin B₁₂ value as low as 40 $\mu\text{g.}/\text{ml.}$ poses certain questions of fundamental importance. Mollin and Ross (1952) observed that a serum level of 100 $\mu\text{g.}/\text{ml.}$ was the critical concentration below which signs of vitamin B₁₂ deficiency could be recognized. They confirmed their observation later by stating that early megaloblastic change was noticed in the bone-marrow of pernicious anaemia (P.A.) patients a few days after the serum vitamin B₁₂ level had fallen below 100 $\mu\text{g.}/\text{ml.}$; at levels greater than this, the marrow remained normoblastic (Mollin and Ross, 1953).

In the present series the bone-marrow was normoblastic in tropical eosinophilia and aplastic anaemia. Bone-marrow examination was not done in normal subjects. Hence it is not possible to state whether they had any sign of vitamin B₁₂ deficiency in the marrow. Repeated examination of peripheral blood did not, however, show any evidence of erythrocytic macrocytosis or granulocytic abnormality. In N.M.A. the bone-marrow was, however, found to be megaloblastic. It must be pointed out that, in N.M.A., megaloblastic bone-marrow was also seen in all the cases with high serum vitamin B₁₂ (see Chart), suggesting that in this disease there is no direct correlation between serum vitamin B₁₂ and bone-marrow changes. It is well known that N.M.A. as seen in the eastern part of India represents deficiency of folic acid and/or vitamin B₁₂ (Das Gupta *et al.*, 1953; Chatterjea, 1958). The lack of correlation between serum vitamin B₁₂ and bone-marrow changes in N.M.A. may therefore be explained on the basis of associated folic-acid deficiency, the degree of which cannot be ascertained from the serum vitamin B₁₂ level.

In none of the cases with low serum vitamin B₁₂, including those of N.M.A., was there any neurological complications. This observation also was not in complete agreement with that of Wokes *et al.* (1955), who recorded neurological symptoms and signs in some of their subjects. In a few Indians with classical Addisonian pernicious anaemia subacute combined degeneration has, however, been found (Das Gupta and Chatterjea, 1951). In N.M.A. and in malabsorption syndrome neurological complications are conspicuous by their absence, though the serum vitamin B₁₂ level in many cases may be as low as in pernicious anaemia. The main difference between P.A. and N.M.A. (or malabsorption syndrome) is with reference to intrinsic factor and neurological complications. In the present series all the vegetarians with low serum vitamin B₁₂ had presumably no deficiency of intrinsic factor. Thus the circumstantial evidence becomes very strong in favour of intrinsic factor having some role, direct or indirect, in protecting the nervous system. If vitamin B₁₂ *per se* was responsible for the protection of the nervous system, absence or rarity of neurological signs in N.M.A., in malabsorption syndrome, and in the present series of vegetarians cannot be satisfactorily explained.

It is possible that vitamin B₁₂ absorbed from a system with an adequate amount of intrinsic factor (as in vegetarians, and in patients with N.M.A. and malabsorption syndrome) may be qualitatively different from vitamin B₁₂ absorbed from a system grossly deficient in intrinsic factor (as in P.A.). Concentration for concentration, the former vitamin B₁₂ may be more protective to the nervous system than the latter. The efficacy of parenteral vitamin B₁₂ therapy in correcting the neurological complication of P.A. may be explained by assuming that the vitamin B₁₂ as used in therapeutics is similar to the former but perhaps different from the latter. An alternative explanation would be the suggestion that therapeutically administered vitamin B₁₂ is "activated" by extragastric sources of vitamin B₁₂ (Callender and Lajtha, 1951). The above facts would tend to indicate that intrinsic factor, in addition to its role as promoter of absorption, may have an activating or maturing influence on vitamin B₁₂. Adequacy of intrinsic factor would ensure not only absorption but also activation or maturation of whatever small amount of vitamin B₁₂ may be available to the system due to dietary inadequacy and/or intestinal malabsorption (Chatterjea, 1960).

An elevated level of serum vitamin B₁₂ in chronic myeloid leukaemia, first reported by Beard *et al.* (1954), has been conclusively confirmed by a number of investigators. The exact cause of this elevated level is not clear. Lack of any differential pattern in the two groups—vegetarian or non-vegetarian—as revealed in the present study, indicates that in this disease the factor responsible for hypervitaminosis far outweighs the hypovitaminic effect of a vegetarian diet.

Summary

The serum vitamin B₁₂ level was studied in 232 subjects (196 non-vegetarians and 46 vegetarians). The subjects belonged to the following five categories: normal, tropical eosinophilia, nutritional macrocytic anaemia, aplastic anaemia, and chronic myeloid leukaemia.

Serum vitamin B₁₂ in the vegetarian group was, in general, lower than that in the non-vegetarian group. Chronic myeloid leukaemia was an exception, all the patients in this category showing a raised level as previously reported.

Low serum vitamin B₁₂ was not, however, associated with any anaemia or neurological complications. The implications of these findings are discussed.

We are indebted to Dr. R. N. Chaudhuri, Director, School of Tropical Medicine, Calcutta, for providing necessary facilities for the study. We are grateful to Mr. S. K. Ghosh and Dr. Sandhya Ghose, of the department of haematology, for their help.

REFERENCES

- Baker, S. J. (1958). *Indian J. Path. Bact.*, **1**, 11.
 Banerjee, D. K., Hosain, F., and Chatterjea, J. B. (1959). *Bull. Calcutta Sch. trop. Med.*, **7**, 147.
 Beard, M. F., Pitney, W. R., and Sanneman, E. H. (1954). *Blood*, **9**, 789.
 Callender, S. T., and Lajtha, L. G. (1951). *Ibid.*, **6**, 1234.
 Chatterjea, J. B. (1958). *J. Indian med. Ass.*, **31**, 4.
 — (1960). *Bull. Calcutta Sch. trop. Med.*, **8**, 9.
 — Banerjee, D. K., Ghose, Sandhya, and Chaudhuri, R. N. (1959). *Ibid.*, **7**, 92.
 Das Gupta, C. R. (1949). *Rep. Scient. Advisory Board for 1948, Indian Res. Fund Ass.*, p. 133. New Delhi.
 — and Chatterjea, J. B. (1951). *Blood*, **6**, 631.
 — and Basu, P. (1953). *Brit. med. J.*, **2**, 645.
 — Ghosh, S. K., and Banerjee, D. K. (1955). *Bull. Calcutta Sch. trop. Med.*, **3**, 101.
 Dhopeswarkar, G. A., Trivedi, J. C., Kulkarni, B. S., Satoskar, R. S., and Lewis, R. A. (1956). *Brit. J. Nutr.*, **10**, 105.
 Lewis, U. J., Register, U. D., Thompson, H. T., and Elvehjem, C. A. (1949). *Proc. Soc. exp. Biol. (N.Y.)*, **72**, 479.
 Mollin, D. L., and Ross, G. I. M. (1952). *J. clin. Path.*, **5**, 129.
 — (1953). *Brit. med. J.*, **2**, 640.
 Peeler, H. T., Yacowitz, H., Carlson, C. W., Miller, R. F., Norris, L. C., and Heuser, G. F. (1951). *J. Nutr.*, **43**, 49.
 Ross, G. I. M. (1952). *J. clin. Path.*, **5**, 250.
 Sokhey, S. S., Gokhale, S. K., Malandkar, M. A., and Billimoria, H. S. (1937). *Indian J. med. Res.*, **25**, 505.
 — (1938). *Ibid.*, **25**, 723.
 Wokes, F., Badenoch, J., and Sinclair, H. M. (1955). *Amer. J. clin. Nutr.*, **3**, 375.

The Italian Olympic Medical Committee is conducting a statistical study of competitors who took part in the Olympic Games in order to discover the correlations between medical characteristics and performance. To obtain this comprehensive medical data—in all, 78 medical characteristics for each athlete—for at least 2,000 of the 7,000 competitors in this year's games, the medical sections of various national Olympic Committees (including the German, Italian, and Russian) have agreed to supply the results of medical examinations of their athletes carried out in their own countries. Once this information has been collected it will be recorded on punched-cards which will then be processed by the IBM 305 RAMAC computer.

TREATMENT OF RESPIRATORY INFECTION IN SCHOOLBOYS WITH PHENETHICILLIN

BY

R. L. VOLLUM, M.A., D.Phil.

Department of Bacteriology, Radcliffe Infirmary, Oxford

AND

B. E. JUEL-JENSEN, M.A., B.M., B.Ch., Cand.Med.

Medical Officer, Summer Fields School, Oxford

The administration of penicillin to children has hitherto presented several practical problems. Crystalline penicillin by injection gives satisfactory blood levels, but is painful and very unpopular with young patients. Oral penicillins G and V are far more acceptable, but the blood levels obtained are much lower than those achieved by parenteral administration, and are indeed often insufficient for satisfactory therapeutic effect. When phenethicillin (6-(α -phenoxypropionamido)penicillanic acid) was marketed as "broxil," it was claimed that this preparation would give blood levels at least equal to those after intramuscular injection of corresponding doses of penicillin G. Knudsen and Rolinson (1959) investigated the absorption and excretion of phenethicillin in adults, and their results confirmed this claim. Garrod (1960) compared the antibacterial activity of phenethicillin, penicillin G, and penicillin V. He found that, weight for weight, phenethicillin was more active than penicillin V or G against resistant staphylococci, and very slightly less active against streptococci and pneumococci.

In February, 1960, an outbreak of severe pharyngitis, with some complications, occurred among schoolboys aged 7–13 at Summer Fields Preparatory School, Oxford. The school is residential, and it was essential to deal promptly with the infection, which spread in a few days to 31 boys. Table I shows the incidence of

TABLE I.—Clinical Diagnosis and Treatment

Condition	No.	Dose of Phenethicillin	Duration of Fever (Hours)	Failure
Pharyngitis due to β -haemolytic streptococci	13	{ 12–250 mg. q.d.s. 1–500 " "	{ 10–24 2–48	—
As above + otitis media ..	3	250 mg. q.d.s.	24	—
Pharyngitis due to other organisms	2	250 " "	{ 1–24 1–48	—
As above + otitis media ..	2	250 " "	48	—
Pharyngitis + bronchitis (2 β -haemolytic streptococci)	5	{ 2–500 mg. q.d.s. 3–250 " "	{ 4–24 1–48	—
Pharyngitis + sinusitis ..	4	{ 3–500 " " 1–250 " "	3–48	2, resistant staphs
Lobar pneumonia ..	1	250 mg. q.d.s.	73, with initial dramatic fall	—
Cellulitis of finger ..	1	250 " "	24	—

the various types of infection. Most of the boys had severe pharyngitis and cervical lymphadenopathy. The illness ran much the same course both in those infected with β -haemolytic streptococci group A and in those from whom other organisms (*Str. pneumoniae*, *Str. viridans*, neisseria) were isolated. Fourteen boys had complications. One boy had uncomplicated lobar pneumonia and one had cellulitis of a finger.

Methods

The boys were all treated with phenethicillin. The standard dose was 250 mg. (one tablet) four-hourly