

Mrs Patricia Wallace for clerical assistance, and Mrs Cathy Harwood for typing the manuscript.

- 1 AIDS Group of the United Kingdom Haemophilia Centres. Seropositivity for HIV in UK haemophiliacs. *Proc R Soc Lond [Biol]* (in press).
- 2 Peto R. Experimental survival curves for interval-censored data. *Applied Statistics* 1973;22:86-91.
- 3 Turnbull BW. The empirical distribution function with arbitrarily grouped, censored and truncated data. *Journal of the Royal Statistical Society* 1976;38:B290-5.
- 4 Brookmeyer R, Goedert JJ. Censoring in an epidemic with an application to haemophilia-associated AIDS. *Biometrics* (in press).
- 5 Cox DR, Oakes D. *Analysis of survival data*. London: Chapman and Hall, 1984.
- 6 SAS Institute. *SAS users guide: statistics*. Cary, North Carolina: SAS Institute, 1985.
- 7 Rizza CR, Spooner RJD. Treatment of haemophilia and related disorders in Britain and Northern Ireland during 1976-80: report on behalf of the directors of haemophilia centres in the United Kingdom. *Br Med J* 1983;286:929-33.
- 8 Brookmeyer R, Gail MH, Polk BF. The prevalent cohort study and the acquired immunodeficiency syndrome. *Am J Epidemiol* 1987;126:14-24.
- 9 Medley GF, Anderson RM, Cox DR, Billard L. Incubation period of AIDS in patients infected via blood transfusion. *Nature* 1987;328:719-21.
- 10 Giesecke J, Scalia-Tomba G, Berglund O, Berntorp E, Schulman S, Stigendal L. Incidence of symptoms and AIDS in 146 Swedish haemophiliacs and

blood transfusion recipients infected with human immunodeficiency virus. *Br Med J* 1988;297:99-102.

- 11 Moss AR, Bachetti P, Osmond D, et al. Seropositivity for HIV and the development of AIDS or AIDS related condition: three year follow up of the San Francisco General Hospital cohort. *Br Med J* 1988;296:745-50.
- 12 Eyster ME, Gail MH, Ballard JO, Al-Mondhry H, Goedert JJ. Natural history of human immunodeficiency virus infections in hemophiliacs: effects of T-cell subsets, platelet counts, and age. *Ann Intern Med* 1987;107:1-6.
- 13 Johnson RE, Lawrence DN, Evatt BL, et al. Acquired immunodeficiency syndrome among patients attending hemophilia treatment centres and mortality experience of hemophiliacs in the United States. *Am J Epidemiol* 1985;121:797-810.
- 14 Lui KJ, Peterman TA, Lawrence DN, Allen JR. A model-based approach to characterize the incubation period of paediatric transfusion-associated acquired immunodeficiency syndrome. *Stat Med* 1988;7:395-401.
- 15 Lui KJ, Lawrence DN, Morgan WM, Peterman TA, Haverkos HW, Bregman DJ. A model-based approach for estimating the mean incubation period of transfusion-associated acquired immunodeficiency syndrome. *Proc Nat Acad Sci USA* 1986;83:3051-5.
- 16 McCormick A. Trends in mortality statistics in England and Wales with particular reference to AIDS from 1984 to 1987. *Br Med J* 1988;296:1289-92.
- 17 Ragui MV, Kingsley LA, Kiss JE, Spero JA, Lewis JH. HIV-related deaths in HIV antibody-positive haemophilic patients. *Lancet* 1987;ii:100.

(Accepted 14 February 1989)

New regression equations for predicting peak expiratory flow in adults

A J Nunn, I Gregg

Abstract

An earlier study of peak expiratory flow (PEF) in normal adults contained too few men aged over 55 and women aged over 65 for the regression equations to be used for prediction in older people. A subsequent study was therefore carried out on an additional 23 men and 29 women aged 55 or over who were lifelong non-smokers and satisfied the same strict criteria of normality that had been used in the original study. The data from both studies were combined and a new model used to calculate equations for the regression of PEF on age and height in the two sexes. With this model predicted values could be derived for men and women aged between 15 and 85. These new equations gave predicted values in men and women aged less than 55 and 65, respectively, which were almost identical with those reported previously.

The new regression equations for PEF enable values to be predicted for people aged 15-85 and so enhance the accuracy of testing in the elderly.

Introduction

Measurement of peak expiratory flow (PEF) by the Wright meter¹ is invaluable for identifying and assessing airflow obstruction in clinical practice.² The introduction of the mini Wright meter³ has resulted in the test being used increasingly by general practitioners.

An observed PEF must be assessed by comparing it with the subject's predicted PEF, which is taken as the mean PEF attainable by "normal" people of the same ethnic origin, sex, age, and body build. The distributors of the Wright and mini Wright meters provide a nomogram for predicting PEF based on a study (reported by us in 1973)⁴ of 202 men and 199 women who were lifelong non-smokers and who fulfilled other stringent criteria of "normality." That study, however, contained few men aged over 55 and women aged over 65, so that the equations were not valid for predicting PEF in older people.

Knowledge of normal ventilatory function is particularly important in older people because of their high prevalence of chronic bronchitis and emphysema. We therefore studied additional men aged 55 or over

and women aged 65 or over who fulfilled the same selection criteria as in our original study.

In this paper we report new equations for the regression of PEF in men and women which were calculated from the combined data of the original and present studies. We compare the new regressions with those which we reported previously⁴ and also with those of other workers who did not restrict their series to non-smokers. In a subsidiary study (reported in our accompanying paper⁵) we used the new equations to investigate the effects of smoking on ventilatory function in elderly men and women who, apart from being smokers or ex-smokers, fulfilled all our other criteria of normality.

Subjects and methods

Between 1975 and 1987 one of us (IG) conducted a search in the course of day to day consultations in two general practices to identify men aged 55 or over, and women aged 65 or over who had never smoked and who denied expectoration, respiratory infection, wheeze, or shortness of breath during childhood or adulthood. The practices, in Kingston upon Thames and Southampton, were similar in their age and social class distribution, and levels of atmospheric pollution were generally low in both areas.

Measurements of PEF were made with three Wright peak flow meters, whose calibration was checked at regular intervals. All the tests were supervised by a single observer (IG), who explained the purpose of the study and then demonstrated the correct manner of performing the test. Subjects were observed while they made several trial attempts in order to detect faults in technique. Once they were able to perform the test correctly they were exhorted to make a maximum effort, and the highest value achieved in three tests in the standing position was recorded to the nearest 5 l/min. Standing height without shoes was measured to the nearest centimetre.

The data from 23 men aged 55 or over and 29 women aged 65 or over who satisfied our criteria of normality were combined with those from the 202 men and 199 women in the original series. In calculating the equations for the regression of PEF on age and height

Medical Research Council
Cardiothoracic
Epidemiology Group,
Brompton Hospital,
London SW3 6HP
A J Nunn, MSc, statistician

Faculty of Medicine,
University of Southampton
I Gregg, FRCP, senior research
fellow

Correspondence and
requests for reprints to:
Mr Nunn.

Br Med J 1989;298:1068-70

in each sex we made the following assumptions: (a) PEF increases to a maximum during early adulthood and thereafter declines with age; (b) at any given age PEF increases with height. These assumptions are satisfied by the model, $\log_e \text{PEF} = a \log_e \text{age} + b \log_e \text{age} + c/\text{height} + d$. The method of least squares analysis was used to determine the equations which gave the best fit with the data.

Results

The table gives the age and sex distribution of the 225 men and 228 women studied. Only seven men were aged 65 or over, whereas the women's ages were more evenly distributed in the older age groups. The equations which gave the best fit with the data were, for men: $\log_e \text{PEF (l/min)} = 0.544 \log_e \text{age} - 0.0151 \text{age} - 74.7/\text{height (cm)} + 5.48$; and for women: $\log_e \text{PEF (l/min)} = 0.376 \log_e \text{age} - 0.0120 \text{age} - 58.8/\text{height (cm)} + 5.63$.

Figure 1 shows the observed values of PEF in the men and women plotted against age. The figure also shows the regressions of PEF on age (obtained from the

Age distribution of 225 men and 228 women who were lifelong non-smokers and who satisfied strict criteria of normality

Age (years)	No (%) of men	No (%) of women
15-24	72 (32)	71 (31)
25-34	51 (23)	33 (14)
35-44	44 (20)	48 (21)
45-54	29 (13)	26 (11)
55-64	22 (10)	22 (10)
65-74	7 (3)	14 (6)
≥75		14 (6)
Total	225 (100)	228 (100)




FIG 1—Observed values of peak expiratory flow (PEF) in 225 men and 228 women who had never smoked and who fulfilled other strict criteria of normality (see text). Regression curves and 90% confidence intervals drawn for mean heights of men and women (177 cm and 162 cm, respectively)

new equations) for the mean heights of the men (177 cm) and women (162 cm) together with their 90% confidence intervals. Maximum values of PEF in both sexes occurred at around 30-35 years of age. After about age 50 PEF fell by about 4 l/min a year in men and 2.5 l/min a year in women. In both sexes the lower 90% confidence limits (below which only 5% of values from normal subjects would be expected to fall) were roughly 70-80 l/min below predicted and did not differ greatly with age. Of the total variation in PEF, only a small part (30% in men, 28% in women) was accounted for by differences in age and height.

Figure 2 shows the new regressions for men with a height of 175 cm and women with a height of 160 cm compared with those which we reported previously.⁴




FIG 2—Mean peak expiratory flow (PEF) in men with height of 175 cm and women with height of 160 cm derived from new regression equations compared with those described by us in 1973⁴ and by other workers. Men: 1=Present series, 2=Gregg and Nunn (1973),³ 3=Leiner et al.,⁶ 4=European Community for Coal and Steel,⁸ Women: 1=Present series, 2=Gregg and Nunn (1973),⁴ 4=European Community for Coal and Steel,⁸ 5=Pelzer and Thomson⁷

In men the largest difference was about 15 l/min at 45-55 years of age, whereas in women it was about 10 l/min during adolescence and between the ages of 25 and 35. Figure 2 also shows the regressions published by some other workers⁶⁻⁸ for men and women of the same heights.

Discussion

The problems of defining "normality" for the purpose of establishing reference values for pulmonary function have been discussed by Miller⁹ and Lebowitz et al.¹⁰ Lifelong non-smoking was the principal criterion that we used in selecting subjects for both our original⁴ and present studies.

It has been contended that predicted values obtained in non-smokers in the general population "reduce their usefulness as indicators of disease."¹¹ Nevertheless, though data from non-smoking communities of Mormons or Seventh Day Adventists¹² may be unrepresentative because of factors associated with their genealogy or lifestyle,⁹ it has been shown that smokers who deny cough or other symptoms have significantly lower levels of PEF than non-smokers,¹³⁻¹⁶ even among adolescents.¹⁷

The effects of smoking on ventilatory function are likely to be particularly great in the elderly by reason of their generally longer exposure. Among older people, especially men, only a small proportion have never smoked.¹⁸ Their inadequate representation in our original series precluded use of the regression equations for predicting PEF in men aged over 55 and women aged over 65. We suggested that approxima-

tions could be made by extending the regression curves linearly beyond these ages,⁴ an expedient that was adopted in the nomogram issued by the distributors of the Wright and mini Wright meters.

The model which we used to calculate the new regression equations was similar to one that Berglund *et al* used in a study of forced expiratory volume in one second and vital capacity.¹⁹ In being curvilinear our regressions differ from those of most other workers,^{6-8 20 21} two of which (Leiner *et al*⁶ in men, Pelzer and Thomson⁷ in women) have been recommended as reference values for predicting PEF.²² As pointed out by Brooks and Waller, linear regressions from 20 years of age give falsely high predicted values in adolescents and young adults (fig 2).²³

Reference values of PEF have been published by a working party of the European Community for Coal and Steel, convened for the purpose of standardising lung function tests.⁸ These were derived by combining data from several studies, all of which had included smokers and ex-smokers unless they admitted to expectoration or gave a history of chest disease. The regressions of the working party and those recommended by Cotes,²² which also included smokers, are substantially lower than ours at all ages in both sexes, except in adolescents (fig 2).

In a person with normal bronchi and alveoli the magnitude of PEF is determined by the pressure exerted in a forced expiration and hence the power of the expiratory muscles; it is also influenced by body build and, in particular, thoracic volume. The only conveniently measured index of body build is standing height, but this correlates poorly with thoracic volume. This explains much of the wide variation in PEF which we found among subjects (fig 1), of which our new equations accounted for only slightly more of the variation in men than our original regressions (30% *v* 23%), whereas in women the amount explained was the same (28%).

A curvilinear regression of PEF, rising to a maximum by about 30-35 years, is consistent with increasing body size and muscular power. The generally greater thoracic volumes and muscular power of men accounted for much of the difference in PEF between men and women of the same age and height. Differences of body build are also the probable explanation for the finding that, independently of smoking habit, PEF in Africans,²⁴⁻²⁷ Indians,^{28 29} and Chinese³⁰ is lower than in European subjects of corresponding age and height. That these differences are of racial rather than environmental origin is suggested by studies of immigrants to Britain. Asians from the Indian subcontinent and east Africa have been found to have lower values of PEF, forced expiratory volume in one second, and forced vital capacity than those of native Britons or immigrants from the West Indies.^{31 32}

Reduced air density at high altitude has little effect on PEF and may be ignored for clinical purposes; values at 3000 m were only 3.5% less than at sea level.³³ On the other hand, voluntary effort has a considerable influence on PEF, which is greater than in other tests of ventilatory function, and it is essential to ensure that subjects make a maximum effort. Close attention must also be paid to the correct technique of performing the test,^{1 10 34} otherwise falsely low values will be obtained.

The measurements of PEF in this study were made with the standard Wright peak flow meter,³ whereas most general practitioners use a mini Wright meter.² Though the values given by the two instruments are

generally in close agreement,³⁵ the mini Wright meter is less robust and apt to lose calibration.³⁶

It is a pleasure to thank Mrs Marion Rickman and Miss Deborah Johnson for their help in analysing the data, and the trustees of the Rudolf Friedlaender Memorial Trust for a grant to Mrs Rickman. During this study IG was supported by grants from the Department of Health and Social Security; the Chest, Heart, and Stroke Association; and Clement Clarke International Ltd.

- 1 Gregg I. The measurement of peak expiratory flow and its applications in general practice. *J R Coll Gen Pract* 1964;7:199-215.
- 2 Wright BM. A miniature Wright peak flow meter. *Br Med J* 1978;ii:1627-8.
- 3 Wright BM, McKerrrow CB. Maximal forced expiratory flow rate as a measure of ventilatory capacity with a description of a new portable instrument for measuring it. *Br Med J* 1959;ii:1041-7.
- 4 Gregg I, Nunn AJ. Peak expiratory flow in normal subjects. *Br Med J* 1973;iii:282-4.
- 5 Gregg I, Nunn AJ. Peak expiratory flow in symptomless elderly smokers and ex-smokers. *Br Med J* 1989;298:1071-2.
- 6 Leiner GC, Abramowitz S, Small MJ, Stenby VB, Lewis WA. Expiratory peak flow rate. Standard values for normal subjects. Use as a clinical test of ventilatory function. *Am Rev Respir Dis* 1963;88:644-51.
- 7 Pelzer AM, Thomson ML. Expiratory peak flow. *Br Med J* 1964;ii:123.
- 8 Quanjer PH, ed. Standardized lung function testing. *Bull Eur Physiopathol Respir* 1983;19suppl 5:1-95.
- 9 Miller A. The use of pulmonary function tests in the evaluation of disease: separating normal from abnormal. *Immunology and Allergy Practice* 1983;5:305-15.
- 10 Lebowitz MD, Quackenboss J, Camilli AE, Bronnimann D, Holberg CJ, Boyer B. The epidemiological importance of intraindividual changes in objective pulmonary responses. *European Journal of Epidemiology* 1987;3:390-8.
- 11 Morris JF, Koski A, Johnson IC. Spirometric standards for healthy non-smoking adults. *Am Rev Respir Dis* 1971;103:57-67.
- 12 Laszlo G. Standardised lung function testing. *Thorax* 1984;39:881-6.
- 13 Read J, Selby T. Tobacco smoking and ventilatory function of the lungs. *Br Med J* 1961;ii:1104-8.
- 14 Zamel N, Youssef HH, Prime FJ. Airway resistance and peak expiratory flow rate in smokers and non-smokers. *Lancet* 1963;ii:1237-8.
- 15 Huhti E. Ventilatory function in healthy non-smokers and smokers. *Scandinavian Journal of Respiratory Diseases* 1967;48:149-55.
- 16 Sluis-Cremer GK, Sichel HS. Ventilatory function in males in a Witwatersrand town. Comparison between smokers and nonsmokers. *Am Rev Respir Dis* 1968;98:229-39.
- 17 Backhouse CI. Peak expiratory flow in youths with varying smoking habits. *Br Med J* 1975;ii:360-2.
- 18 Milne JS, Williamson J. Respiratory function tests in older people. *Clin Sci* 1972;42:371-81.
- 19 Berglund E, Birath G, Bjore J, *et al*. Spirometric studies in normal subjects. Forced expirograms in subjects between 7 and 70 years of age. *Acta Med Scand* 1963;173:185-92.
- 20 Ferris BG, Anderson DO, Zickmantel R. Prediction values for screening tests of pulmonary function. *Am Rev Respir Dis* 1965;91:252-61.
- 21 Leonards AK. Der maximale Expirationsstrom. Untersuchung der Lungenfunktion bei Gesunden. *Acta Allergologica* 1966;21:99-138.
- 22 Cotes JE. *Lung function: assessment and application in medicine*. 4th ed. Oxford: Blackwell Scientific, 1979.
- 23 Brooks AGF, Waller RE. Peak flow measurements among visitors to a public health exhibition. *Thorax* 1972;27:557-62.
- 24 Elebute EA, Femi-Pearse D. Peak flow rate in Nigeria: anthropometric determinants and usefulness in assessment of ventilatory function. *Thorax* 1971;26:597-601.
- 25 Johannsen ZM, Erasmus LD. Clinical spirometry in normal Bantu. *Am Rev Respir Dis* 1968;97:585-97.
- 26 Teklu B, Seboxa T, Mills RJ. Peak expiratory flow in normal Ethiopian children and adults in Addis Ababa. *Br J Dis Chest* 1987;81:176-81.
- 27 Cookson JB, Blake GTW, Faranisi C. Normal values for ventilatory function in Rhodesian Africans. *Br J Dis Chest* 1976;70:107-11.
- 28 Malik SK, Banga N. Peak expiratory flow rates in non-smoking rural males. *Indian J Chest Dis Allied Sci* 1978;24:183-6.
- 29 Kamat SR, Tyagi NK, Rashid SSA. Lung function in adult subjects. *Lung India* 1982;1:1-11.
- 30 Lam KK, Pang SC, Allen WGL, Hill LE, Snell NJC, Nunn AJ. A survey of ventilatory capacity in Chinese subjects in Hong Kong. *Ann Hum Biol* 1982;9:459-72.
- 31 Malik MA, Moss E, Lee WR. Prediction values for the ventilatory capacity in male West Pakistani workers in the United Kingdom. *Thorax* 1972;27:611-9.
- 32 Jackson SHD, Cruickshank JK, Beevers DG, Bannan LT. Ethnic differences in peak expiratory flow rate in Birmingham factory workers. *Postgrad Med J* 1983;59:671-3.
- 33 Forster P, Parker RW. Peak expiratory flow rate at high altitude. *Lancet* 1983;ii:100.
- 34 Connolly CK. Falsely high peak expiratory flow readings due to acceleration in the mouth. *Br Med J* 1987;294:285.
- 35 Morrill CG, Dickey DW, Weiser PC, Kinsman RA, Chai H, Spector SL. Calibration and stability of standard and mini-Wright peak flow meters. *Ann Allergy* 1981;46:70-3.
- 36 Oldham HG, Bevan MM, McDermott M. Comparison of the new miniature Wright peak flow meter with the standard Wright peak flow meter. *Thorax* 1979;34:807-9.

(Accepted 27 February 1989)