

What is already known on this topic

Hysterectomy is a common operation

Little is known about the long term effects of hysterectomy

What this study adds

Hysterectomy did not significantly increase a woman's risk of mortality from all causes, cardiovascular disease, and cancer

menopausal soon after hysterectomy. We were therefore unable to carry out separate analyses using menopausal status.

Previous studies have looked at risk of specific cancers after hysterectomy, rather than all cancer mortality. The reduced risk of ovarian cancer after hysterectomy found in one study¹⁹ may have been due to a screening effect, as surgery provides an opportunity to detect abnormal ovaries. Such effects would persist for as long as it takes visible premalignant abnormalities to produce symptoms of cancer.¹⁹ This bias could have occurred in our study, although it is not clear how long such a protective effect might have influenced our risk estimates of all cancer mortality.

Most women in our study had a hysterectomy for non-malignant reasons. They would no longer be at risk of endometrial, cervical, or ovarian cancer if they also had bilateral oophorectomy. Cancers comprise more than a third of deaths in middle aged women, with many at gynaecological sites. The observed lower risk of death (although not statistically significant) from all causes and from cancer among young women who had a hysterectomy was therefore unsurprising. Our results ignore any non-fatal, physical, psychological, and social costs to the individual after hysterectomy. The results should therefore not be used to argue that hysterectomy be used as a public health measure to reduce women's risk of death later in life. Instead, patients should be reassured that hysterectomy will not put their lives at risk later on.

We thank Val Angus for extracting the data, the doctors who have contributed to the oral contraception study, and the Chief Scientist Office, Scottish Executive, who funded LI as part of a research training fellowship to complete the masters degree of which this study was a component.

Contributors: See bmj.com

Funding: The oral contraception study has received support from the British Heart Foundation, Cancer Research UK, Medical Research Council, Royal College of General Practitioners, Organon Laboratories, Ortho Pharmaceutical, Schering AG, Schering Health Care, Syntex Pharmaceuticals, GD Searle, Syntex Pharmaceuticals, John Wyeth, and Brother.

Competing interests: None declared.

Ethical approval: The study was part of a masters degree submission and received approval from the ethics committee of the London School of Hygiene and Tropical Medicine.

- 1 Vessey MP, Villard-Mackintosh L, McPherson K, Coulter A, Yeates D. The epidemiology of hysterectomy: findings in a large cohort study. *Br J Obstet Gynaecol* 1992;99:402-7.
- 2 Bush TL, Cowan LD, Barrett-Connor E, Criqui MH, Karon JM, Wallace RB, et al. Estrogen use and all-cause mortality. Preliminary results from the lipid research clinics program follow-up study. *JAMA* 1983;249:903-6.
- 3 Luoto R, Auvinen A, Pukkala E, Hakama M. Hysterectomy and subsequent risk of cancer. *Int J Epidemiol* 1997;26:476-83.
- 4 Green A, Purdie D, Bain C, Siskind V, Russell P, Quinn M, et al. Tubal sterilisation, hysterectomy and decreased risk of ovarian cancer. *Int J Cancer* 1997;79:948-51.

- 5 Riman T, Persson I, Nilsson S. Hormonal aspects of epithelial ovarian cancer: review of epidemiological evidence. *Clin Endocrinol* 1998;49:695-707.
- 6 Kreiger N, Sloan M, Cotterchio M, Kirsh V. The risk of breast cancer following reproductive surgery. *Eur J Cancer* 1999;35:97-101.
- 7 Parazzini F, Braga C, La Vecchia C, Negri E, Acerboni S, Franceschi S. Hysterectomy, oophorectomy in premenopause and risk of breast cancer. *Obstet Gynecol* 1997;90:453-6.
- 8 Gago-Dominguez M, Castela JE, Yuan J-M, Ross RK, Yu MC. Increased risk of renal cell carcinoma subsequent to hysterectomy. *Cancer Epidemiol Biomark Prevent* 1999;8:999-1003.
- 9 Royal College of General Practitioners. *Oral contraceptives and health*. Tunbridge Wells: Pitman Medical, 1974.
- 10 General Registrar Office. *Classification of occupations*. London: HMSO, 1966.
- 11 World Health Organisation. *International classification of diseases, injuries and causes of death, 8th revision*. Geneva: WHO, 1967.
- 12 Beral V, Hermon C, Kay C, Hannaford PC, Darby S, Reeves G. Mortality in relation to method of follow-up in the Royal College of General Practitioners' oral contraception study. In: Hannaford PC, Webb AMC, eds. *Evidence-guided prescribing of the pill*. Lancashire: Parthenon; 1996:327-39.
- 13 Beral V, Banks E, Reeves G. Evidence from randomised trials on the long-term effects of hormone replacement therapy. *Lancet* 2002;360:942-4.
- 14 Moorhead T, Hannaford P, Warskyj M. Prevalence and characteristics associated with use of hormone replacement therapy in Britain. *Br J Obstet Gynaecol* 1997;104:290-7.
- 15 Office for National Statistics. Smoking. In: *Living in Britain. The 2002 general household survey*. www.statistics.gov.uk/lib2002/downloads/smoking.pdf (accessed 25 Aug 2004).
- 16 Owen-Smith V, Hannaford PC, Warskyj M, Ferry S, Kay CR. Effects of changes in smoking status on risk estimates for myocardial infarction among women recruited for the Royal College of General Practitioners' oral contraception study in the UK. *J Epidemiol Community Health* 1998;52:420-4.
- 17 Rosenberg L, Hennekens CH, Rosner B, Belanger C, Rothman KJ, Speizer FE. Early menopause and the risk of myocardial infarction. *Am J Obstet Gynecol* 1981;139:47-51.
- 18 Falkeborn M, Schairer C, Naessén T, Persson I. Risk of myocardial infarction after oophorectomy and hysterectomy. *J Clin Epidemiol* 2000;53:832-7.
- 19 Irwin KL, Weiss NS, Lee NC, Peterson HB. Tubal sterilisation, hysterectomy, and the subsequent occurrence of epithelial ovarian cancer. *Am J Epidemiol* 1991;134:362-9.

(Accepted 9 May 2005)

doi 10.1136/bmj.38483.669178.8F

Corrections and clarifications

Recent developments in inhaled therapy in stable chronic obstructive pulmonary disease

In the second paragraph of the section "Long acting inhaled bronchodilators" in this Clinical Review by C B Cooper and D P Tashkin (*BMJ* 2005;330:640, 19 Mar), the final sentence should have said that tiotropium increases (not reduces) the time to first exacerbation compared with placebo. (Figure 2 in the article confirms this statement.)

Management of pregnancies with RhD alloimmunisation

We mixed up images and captions in this Clinical Review by Sailesh Kumar and Fiona Regan (*BMJ* 2005;330:1255-8, 28 May). The caption published with figure 1 should have appeared with figure 2, and the caption for figure 1 should have read: "Ultrasound image showing features of hydrops (skin oedema, hepatomegaly, and ascites)." In the text, these two figures should have been referenced in the Pathophysiology section (the third sentence from the end (fig 1)) and in the seventh sentence of the second paragraph of the Monitoring section (fig 2).

Randomised controlled trial to compare surgical stabilisation of the lumbar spine with an intensive rehabilitation programme for patients with chronic low back pain: the MRC spine stabilisation trial

An oversight in the editorial process of this paper by J Fairbank and colleagues led to the omission of the international trial number (*BMJ* 2005;330:1233-9, 28 May). The number is ISRCTN88854663.