potentially useful for patients with poor blood pressure control who are taking three or more drugs. Using the guidelines outlined in this article and restricting the use of spironolactone to 25 mg, we have found the incidence of serious hyperkalaemia and substantial renal dysfunction to be low. By careful monitoring, however, we have also identified occasional serious cases of hyperkalaemia and stopped treatment before adverse events occurred.

We thank T M Reynolds and W S A Smellie, who wrote the original guidance, and the people who reviewed the original guidance and made valuable comments in addition to those of the steering group-I S Young (Association of Clinical Biochemists), R Gama (Association of Clinical Pathologists), N Campbell (Royal College of General Practitioners), D Housley (Department of Biochemistry, Luton and Dunstable NHS Trust), and J Main (Department of Nephrology, South Tees NHS Trust).

Competing interests: None declared.

- 1 Iglesias JI, Hamburger RJ, Feldman L, Kaufman JS. The natural history of incidental renal artery stenosis in patients with aortoiliac vascular disease. Am I Med 2000:109:642-7.
- Smellie WSA. Best practice in primary care pathology: review 6. J Clin Pathol 2006; doi: 10.1136/jcp.2006.040014
- 3 Palmer BF. Renal dysfunction complicating the treatment of hypertenranner of the second completening are deduced on type test sion. N Engl J Med 2002;347:1256-61. Bakris GL, Weir MR. Angiotensin-converting enzyme inhibitor-
- 4 associated elevations in serum creatinine: is this a cause for concern? Arch Intern Med 2000;160:685-93.
- 5 Main J. When should atheromatous renal artery stenosis be considered? A guide for the general physician. Clin Med 2003;3:520-5
- Safian RD, Textor SC. Renal-artery stenosis. N Engl J Med 2001;344:431-6 42. 7
- Pitt B, Zannad F, Remme WJ, Cody R, Castaigne A, Perez A, et al. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. *N Engl J Med* 1999;341:709-17. McMurray J, Cohen-Solal A, Dietz R, Eichhorn E, Erhardt L, Hobbs R, et
- al. Practical recommendations for the use of ACE inhibitors, beta-blockers and spironolactone in heart failure: putting guidelines into practice. Eur J Heart Fail 2001;3:495-502.
- Williams B, Poulter NR, Brown MJ, Davis M, McInnes GT, Potter JF, et al. Guidelines for management of hypertension: Report of the fourth work-ing party of the British Hypertension Society, 2004–BHSIV. J Hum Hypertens 2004;18:139-85

bmjlearning.com

- 10 Palmer BF. Managing hyperkalemia caused by inhibitors of the renin-angiotensin-aldosterone system. N Engl J Med 2004;351:585-92.
- McMurray JJV, O'Meara E. Treatment of heart failure with
- Institution (J. 1997).
 spironolactone—trial and tribulations. *N Engl J Med* 2004;351:526-8.
 Juurlink DN, Mamdani MM, Lee DS, Kopp A, Austin PC, Laupacis A, et al. Rates of hyperkalemia after publication of the randomized aldactone evaluation study. N Engl J Med 2004;351:543-51.
- 13 National Institute for Health and Clinical Excellence. Hypertension: management of hypertension in adults in primary care. Partial update of NICE clinical guideline 18. London: NICE, 2006.

(Accepted 18 September 2006)

doi 10.1136/bmj.38987.502512.DE

Corrections and clarifications

Short cuts: What's new in the other general journals In the "Redistributing wealth is good for health" item by Alison Tonks (BMJ 2006;333:643-4, 23 Sep), we omitted a vital dot in the "doi" (that's direct object identifier for those who're interested). This prevented people from linking to the original Lancet article or searching successfully for it. The correct doi is 10.1016/S0140-6736(06)69341-0.

Totally transforming Auchendreich

A currency conversion went awry in this Soundings piece by Colin Douglas (BMJ 2006;333:609, 16 Sep). In the third paragraph, the consultancy cost of £30m should have been converted to €44.7m and \$56.7m.

Sudanese health minister's advocacy of condoms sparks brotest

In this article by Peter Moszynski (BMJ 2006;332:1233), we wrongly said that Sudan's health minister, Dr Tabitha Sokaya (who was appointed to the position in September 2005), was "the first woman to hold the post and the first Christian." In fact, we should have said "the first Christian woman to hold the post."

BMJ Learning Foundation Programme

Starting life as a new doctor is tough. There is lots of work to do and a seemingly infinite amount to learn in a short space of time. Pressures come from seniors and interdisciplinary staff and also from patients. If you feel overwhelmed at times then you are not alone: all doctors have gone through this experience, and thousands more are going through it this month.

So much for tea and sympathy, what can we do to help? Our latest online learning resource is the BMJ Learning Foundation Programme. If you are starting out on foundation year one or are progressing to year two then this programme is designed to help. It tells you what's in the curriculum and syllabus and what exactly you'll have to learn and achieve over the following year. It also explains what forms you need to fill in and outlines how to do your assessments. But its most important offering is its learning resources. There are a variety of interactive video modules on the site to help you learn about different aspects of the curriculum.

You can learn about the care of acutely ill patients in our diagnosis module, which was filmed at St Mary's

Hospital, Paddington. In this module you'll meet a variety of real patients with various illnesses and will be able to pit your wits against the medical staff as they establish what is wrong with the patients and start treatment. In the simulation module you'll find out what goes on at the simulation centre at Chelsea and Westminster Hospital. You may get a chance to go to such a centre during your foundation years, but, as there are relatively few simulation centres, any time that you spend there is likely to be short. Following the lead from the aviation industry, our module enables you to test and train your emergency skills online, and you can do this at any time and in any place. To use the site you'll need a computer and broadband access to the web.

You can get all of this at http://bmjlearning.com/ planrecord/foundation/index.htm. Let us know what you think.

Kieran Walsh clinical editor, BMJ Learning (bmjlearning@bmjgroup.com)