

BMJ 2018;360:k568 doi: 10.1136/bmj.k568 (Published 22 February 2018)

MINERVA



Blisters and a fever

Chris Lawrence infectious diseases registrar, Immanuel Rhema foundation doctor, Gaia Nebbia, consultant virologist

Guy's and St Thomas' Hospital, London, UK

A 23 year old woman presented with a week of coryzal illness, fever, and sore throat. She had painful vesicular lesions on the soft palate and tonsils (). Polymerase chain reaction testing of a lesion swab on day 4 of the illness detected enterovirus RNA, confirming a clinical diagnosis of herpangina.



Several enteroviruses can cause herpangina, including coxsackievirus A and B and enterovirus A71. Outbreaks occur in children, and occasionally neonates and young adults. Differential diagnoses for vesicular oral lesions include herpes simplex virus and hand, foot, and mouth disease. However, in hand, foot, and mouth, the oral lesions are usually painless, and accompanied by vesicular lesions on the hands and feet.

Herpangina is typically a self limiting illness; management is supportive and contact precautions are recommended until lesions have resolved.

Patient consent obtained.

Published by the BMJ Publishing Group Limited. For permission to use (where not already granted under a licence) please go to http://group.bmj.com/group/rights-licensing/ permissions