- ¹⁰ Rösen P, Schrör K. Increased prostacyclin release from perfused hearts of acutely diabetic rats. *Diabetologia* 1980;**18**:391-4.
- ¹¹ Davis TME, Mitchell MD, Turner RC. Prostacyclin and thromboxane metabolites in diabetes. *Lancet* 1979;iii:789-90.
- ¹² Davis TME, Mitchell MD, Dornan TL, *et al*. Plasma 6-keto-PGF_{1α} concentrations and diabetic retinopathy. *Lancet* 1980;i:373.
- ¹³ Dollery CT, Friedman LA, Hensby CN, *et al*. Circulating prostacyclin may be reduced in diabetes. *Lancet* 1979;ii:1365.
- ¹⁴ Mitchell MD. A sensitive radioimmunoassay for 6-keto-PGF_{1α}: preliminary observations on circulating concentrations. *Prostaglandins Med* 1978;**1**:13-21.
- ¹⁵ Kynoch PAM, Lehman H. Rapid estimation (2½h) of glycosylated haemoglobins for routine purposes. *Lancet* 1977;ii:16.
- ¹⁶ De Bono D. En Face organ culture of vascular endothelium. *Br J Exp Pathol* 1975;**56**:8-13.
- ¹⁷ Reinsch CH. Smoothing by spline functions. *Numerische Mathematik* 1967;**10**:177-83.
- ¹⁸ Siegel S. Non-parametric statistics. New York: McGraw Hill, 1956.
- ¹⁹ Moncada S, Higgs EA, Vane JR. Human arterial and venous tissues generate prostacyclin (prostaglandin X), a potent inhibitor of platelet aggregation. *Lancet* 1977;i:18-20.
- ²⁰ Jorgensen KA, Stoffersen E, Dyerberg J. Stability of prostacyclin in plasma. *Lancet* 1979;i:1352.
- ²¹ Pace-Asciak CR, Carrara MC. Age-dependent increase in the formation of prostaglandin I₂ by intact and homogenised aortae from the developing spontaneously hypertensive rat. *Biochim Biophys Acta* 1979;**574**:177-81.
- ²² Steer ML, MacIntyre DE, Levine L, *et al*. Is prostacyclin a physiologically important circulating anti-platelet agent? *Nature* 1980;**283**:194-5.

(Accepted 3 February 1981)

Impaired monocyte function in liver cirrhosis

AVI HASSNER, YEHUDIT KLETTER, DORIT SHLAG, MEDINA YEDVAB, MOSHE ARONSON, SHLOMO SHIBOLET

Abstract

Monocyte function in patients with cirrhosis of the liver was measured by phagocytosis and killing of *Candida pseudotropicalis* and *C. albicans*. Both variables were significantly decreased in the patients compared with controls. Control monocytes exposed for two hours to patients' serum showed a significant decrease in intracellular killing compared with control monocytes incubated in autologous serum. This suggests the presence of an inhibiting factor in the patients' serum. This inhibitory factor passed through a dialysis membrane that permitted the passage of molecules of less than 12 000 daltons. Treating monocytes from patients with trypsin significantly increased phagocytosis, indicating that the possible inhibitory factor was attached to the monocyte surface. Metabolism of monocytes during phagocytosis as determined by chemoluminescence was significantly lower in monocytes from patients compared with controls, indicating metabolic impairment.

Monocytes are a component of the monocyte-macrophage system, which includes Kupffer's cells. Impairment of the function of these cells, which have a pivotal role in clearing portal blood, might well be extremely important in the development of chronic liver disease.

Introduction

Various defects have been described in the immune defence system of patients with liver cirrhosis that might explain the increased susceptibility of these patients to infection. Impaired delayed hypersensitivity reactions,¹ impaired humoral function,² and impaired production of complement by the diseased liver³ have been described. In a review on this subject, however, Sherlock⁴ stated that there were no data showing decreased function of Kupffer's cells in liver cirrhosis but as a result of collateral circulation these cells were bypassed. While Kupffer's cells and monocytes are considered to be components of the monocyte-macrophage system,⁴ evaluation of monocyte function in patients with cirrhosis of the liver might contribute to our understanding of the factors leading to the increased tendency for infection. We therefore investigated phagocytosis and intracellular killing by blood monocytes from patients with liver cirrhosis.

Subjects and methods

We studied 40 patients (17 women, 23 men) who had liver cirrhosis that had been proved on biopsy. The mean age was 41 (range 25-67) years. Twenty-eight patients had alcoholic cirrhosis, eight post-necrotic cirrhosis, and the others cryptogenic or an undefined type of liver cirrhosis. None had signs of infection during the study, and none received antibiotics, steroids, or other immunosuppressive drugs. We also studied 32 subjects (12 women, 20 men) who were healthy or had angina and served as control group. None of them had diabetes mellitus or received drugs during the study.

Mononuclear cells from heparinised venous blood were isolated by buoyant centrifugation in Ficoll-Paque (Pharmacia).⁵ *Candida pseudotropicalis* and *C. albicans* were grown and phagocytosis and intracellular killing measured according to the method of Territo and Cline.⁶ Serum from patients was dialysed for 24 hours at 4°C against physiological saline through dialysis tubing with a 12 000 dalton cut-off (A H Thomas Co, Philadelphia, USA). Monocytes from healthy donors were incubated for 120 minutes at 4°C in autologous serum, untreated patients' serum, and patients' serum after dialysis. After incubation the cells were washed twice in Hanks's Balanced Salt Solution and heparin, and phagocytosis and intracellular killing in autologous serum were measured. Monocytes from patients were incubated for 30 minutes at 37°C in 0.25% trypsin, washed twice, and their function subsequently evaluated.

Chemoluminescence was measured in monocytes from the controls and patients 30 minutes after incubation with *C. albicans* in autologous

Department of Medicine D, Ichilov Municipal Hospital of Tel-Aviv, and Department of Histology and Cell Biology, Sackler School of Medicine, Tel-Aviv University, Tel-Aviv, Israel

AVI HASSNER, MD, house physician and instructor in internal medicine
YEHUDIT KLETTER, PHD, research fellow, department of histology and cell biology

DORIT SHLAG, MD, house physician

MEDINA YEDVAB, MD, deputy head of department of internal medicine and lecturer

MOSHE ARONSON, PHD, associate professor, department of histology and cell biology

SHLOMO SHIBOLET, MD, head of department of internal medicine and associate professor

serum. Measurements were performed in a scintillation counter (Packard Tri-Carb 3255) for 30 seconds.⁷

Statistical significance was evaluated using Student's *t* test.

Results

The tables summarise phagocytosis and intracellular killing of candida by the monocytes. Table I shows that there was a distinct decrease in phagocytosis and intracellular killing of both *Candida* species by monocytes from patients as compared with monocytes from controls.

TABLE I—Phagocytosis (mean $\pm$ SD number of yeast cells/monocyte) and intracellular killing (mean $\pm$ SD percentage of killed yeast cells) by monocytes from control subjects and patients with liver cirrhosis

	Patients (group 1)	Controls (group 2)	Patients (monocytes exposed to trypsin) (group 3)
Phagocytosis			
<i>C pseudotropicalis</i>	1.19 $\pm$ 0.07	1.27 $\pm$ 0.08	1.45 $\pm$ 0.15
Groups 1 v 2 and 1 v 3		<i>p</i> < 0.0025	<i>p</i> < 0.0005
<i>C albicans</i>	1.23 $\pm$ 0.07	1.33 $\pm$ 0.11	1.54 $\pm$ 0.11
Groups 1 v 2 and 1 v 3		<i>p</i> < 0.0025	<i>p</i> < 0.0005
Intracellular killing			
<i>C pseudotropicalis</i>	25.07 $\pm$ 6.03	39.95 $\pm$ 8.32	21.2 $\pm$ 3.75
Groups 1 v 2 and 1 v 3		<i>p</i> < 0.0005	<i>p</i> < 0.01
<i>C albicans</i>	27.81 $\pm$ 9.25	42.98 $\pm$ 7.07	24.76 $\pm$ 4.9
Groups 1 v 2 and 1 v 3		<i>p</i> < 0.0005	<i>p</i> < 0.05

Table II shows phagocytosis and intracellular killing by normal monocytes after incubation in patients' serum, dialysed patients' serum, and normal serum. Intracellular killing but not phagocytosis was reduced when the normal monocytes were incubated in patients' serum. When patients' serum was dialysed through a membrane permitting passage of molecules of less than 12 000 daltons it lost its inhibiting capacity and no difference in intracellular killing was found. After exposure of monocytes to trypsin phagocytosis was increased, possibly indicating that an inhibitory factor has been removed from the surface of these monocytes (table I). Chemoluminescence (expressed as mean $\pm$ SD cpm $\times 10^5/10^5$ monocytes in 30 seconds) was 2.64 ± 1.62 in the patients' monocytes and 7.62 ± 6.16 in the control monocytes. This difference is significant (*p* < 0.025). These data indicate an impairment of metabolism in the patients' monocytes, which might also be linked to the inhibitory factor in their serum.

TABLE II—Effect of serum on phagocytosis (mean $\pm$ SD number of yeast cells/monocyte) and intracellular killing (mean $\pm$ SD percentage of killed yeast cells) by normal monocytes

	Control monocytes exposed for two hours to:		
	Patients' serum (group 1)	Autologous serum (group 2)	Patients' serum after dialysis (group 3)
Phagocytosis			
<i>C pseudotropicalis</i>	1.32 $\pm$ 0.04	1.32 $\pm$ 0.04	1.35 $\pm$ 0.05
Groups 1 v 2 and 1 v 3		NS	NS
<i>C albicans</i>	1.39 $\pm$ 0.04	1.39 $\pm$ 0.07	1.45 $\pm$ 0.07
Groups 1 v 2 and 1 v 3		NS	<i>p</i> < 0.025
Intracellular killing			
<i>C pseudotropicalis</i>	30.78 $\pm$ 2.1	42.37 $\pm$ 4.16	40.57 $\pm$ 6.36
Groups 1 v 2 and 1 v 3		<i>p</i> < 0.0005	<i>p</i> < 0.0005
<i>C albicans</i>	34.22 $\pm$ 4.13	44.25 $\pm$ 3.6	43.79 $\pm$ 6.66
Groups 1 v 2 and 1 v 3		<i>p</i> < 0.0005	<i>p</i> < 0.0025

Discussion

There is ample evidence that in patients with cirrhosis of the liver several components of cell-mediated immunity and humoral immunity are impaired.^{1,2} Studies of monocyte function in these patients have been scarce, however, possibly because no routine methods are available for its assessment. Holdstock *et al* showed in a preliminary publication that monocytes from patients with cirrhosis of the liver have a

defect in chemotaxis.⁸ Ganguly *et al* found increased activity of acid hydrolases in monocytes from patients with cirrhosis of the liver.⁹

Our data show that in patients with cirrhosis of the liver monocyte function as determined by phagocytosis and killing of *C albicans* and *C pseudotropicalis* is significantly decreased and that metabolic activity of these cells as measured by chemoluminescence is low. In addition, a dialysable factor with a molecular weight of less than 12 000 daltons seems to exist in the patients' serum, which lowers intracellular killing of both *Candida* species by normal monocytes. We also found that proteolytic treatment of patients' monocytes with trypsin improved their phagocytosis but not their killing ability. The observed differences of action of the serum factor impairing intracellular killing and the fact that trypsinisation of monocytes improves phagocytosis would suggest that more than one factor or mechanism of action plays a part. Various serum proteins impair cellular immunity—for example, ferritin,¹⁰ α -fetoprotein,¹¹ and endotoxin.¹² Of these, only endotoxin has a molecular weight of less than 12 000 daltons.

The patients that we studied did not represent a homogeneous group but had liver disease of diverse aetiology. Monocytes have lately been considered to be a component of the monocyte-macrophage system, which includes among others Kupffer's cells. There is as yet no evidence that impaired monocyte function in a patient would also point to altered function of Kupffer's cells. While liver macrophages function poorly in liver cirrhosis as indicated by poor clearance of endotoxins from the gastrointestinal tract,¹³ Sherlock³ stated that these observations could well be explained by the altered portal circulation causing the liver and its cells to be bypassed. In view of the pivotal role of Kupffer's cells in clearing portal blood a possible decrease in their function like the one we describe in the monocytes could well be extremely important in the development of chronic liver disease. We therefore suggest that future data on the function of Kupffer's cells in patients with cirrhosis of the liver will lead to a better understanding of the pathophysiology of chronic liver disease.

This study was supported in part by grant No 598 from the Chief Scientist's Office of the Israel Ministry of Health.

References

- Berenyi MR, Straus B, Crus D. In vitro and in vivo studies of cellular immunity in alcoholic cirrhosis. *American Journal of Digestive Diseases* 1974;**19**:199-205.
- Thomas HC. The immune response in hepatic cirrhosis: animal and human studies. *Proc R Soc Med* 1977;**70**:521-5.
- Sherlock S. Immunological changes in liver disease. *Proc R Soc Med* 1977;**70**:851-5.
- Van-Furth R, Diesselhoff-Den Dulk MCC. The kinetics of promonocytes and monocytes in the bone-marrow. *J Exp Med* 1970;**132**:813-28.
- Boyum A. Isolation of mononuclear cells and granulocytes from human blood. *Scand J Clin Lab Invest* 1968;**21**, suppl 97:77-89.
- Territo MC, Cline MJ. Monocyte function in man. *J Immunol* 1977;**118**:187-92.
- Allen RC, Yevich SJ, Orth RW, Steele RH. The superoxide anion and singlet molecular oxygen: their role in the microbicidal activity of the polymorphonuclear leukocytes. *Biochem Biophys Res Commun* 1974;**60**:909-17.
- Holdstock GE, Hill S, Wright R. Peripheral monocytes in chronic liver disease. *Gastroenterology* 1979;**77**:A17.
- Ganguly NK, Lloyd B, Price CP, *et al*. Acid hydrolases in monocytes from patients with inflammatory bowel disease, chronic liver disease, and rheumatoid arthritis. *Lancet* 1978;**ii**:1073-5.
- Giller S, Moroz H. Ferritin—its structure, chemical and immunological properties and importance in malignant diseases. *Harefuah* 1978;**94**:236-8.
- Murgita RA, Andersson LC, Sherman MS, Bennich H, Wigzel H. Effects of human alpha-fetoprotein on human B and T lymphocyte proliferation in-vitro. *Clin Exp Immunol* 1978;**33**:347-56.
- Golde LM, Scher I, Osborne B, Rosenstreich L. Cellular mechanism of endotoxin unresponsiveness in C₃H/HeJ mice. *J Immunol* 1976;**116**:454-61.
- Triger DR, Boyer TD, Levin J. Portal and systemic bacteraemia and endotoxaemia in liver disease. *Gut* 1978;**19**:935-9.

(Accepted 20 February 1981)