Current Practice

TODAY'S DRUGS

With the help of expert contributors we publish below notes on a selection of drugs in current use.

Radioiodine Therapy of Thyrotoxicosis

Isotopes of iodine are used principally in the investigation of thyroid function, and the chief therapeutic application of radioiodine (^{131}I) is in the treatment of thyrotoxicosis.

Radiobiological Effects

The body does not distinguish the radioisotopes of iodine from stable iodine (^{127}I), and consequently radioiodine enters the circulating iodide pool. This pool normally consists of stable iodide derived from the diet and from the breakdown of the thyroid hormones. Radioiodine is taken up from the iodide pool by the thyroid and is incorporated into the structure of the thyroid hormones. These are released into the circulation and are transported to the tissues. After the breakdown of the hormones the radioiodine, like the stable iodine, may be taken up by the thyroid again and recycled.

The radiation dose from an isotope depends partly on the physical half-life; for ^{131}I this is eight days. Within the body, however, biological factors are also important. The proportion of the dose taken up by the thyroid, the rate of elimination of the radioiodine from the body, and the mass of the thyroid all influence the dose of radiation absorbed from a given quantity of radioiodine. These variables can be estimated and together they determine the effective biological half-life of the radioiodine.

The effects of radioiodine are immediate death of a proportion of the thyroid cells and impairment of secretion and, subsequently, failure of the surviving cells to divide. Furthermore, the ability of the cell to divide is related to the dose of radioiodine. These effects have been shown in laboratory animals^{1,2} and can be deduced from observations in man.^{3,4}

Dosage

Radioiodine is usually dispensed in a carrier solution of sodium iodide and is tasteless. The dose is taken by mouth.

There is no uniform method for calculating the ^{131}I dose. Most centres aim to give a fixed radiation dose to the thyroid and this is estimated from the 24- or 48-hour uptake of a tracer dose of ^{131}I and the mass of the thyroid. The latter is usually assessed by palpation but is often inaccurate. The mean estimate of a group of observers will be more accurate than a single observation, and experience in estimating pre-operative thyroid weights will help to improve an individual doctor's estimates.⁵ Alternatively it is possible to calculate the mass of the thyroid from a thyroid scintiscan.⁶ Some therapists still prescribe arbitrary doses based on their clinical assessment of the patient without reference to the thyroid size.

Clinical Effect

The response to radioiodine is not immediate. At least six weeks elapse before any definite improvement is observed, and clinical progress thereafter is variable. The experience of most centres is that about 20 to 30% of patients are still thyrotoxic at the end of a year.⁷

If the patient becomes euthyroid there is the problem of post- ^{131}I hypothyroidism occurring later. At first it was thought that hypothyroidism would appear in the first 12 months and that the chance of hypothyroidism developing after then was remote. Long-term follow-up studies of patients treated with radioiodine, however, have revealed an unexpected picture. The incidence of hypothyroidism rises steadily and in 10 years reaches 40 to 50% of all patients treated.^{4,8-11} Furthermore, even after this interval there is no evidence of a plateau in the incidence of hypothyroidism.

The presentation of hypothyroidism is frequently insidious and patients who fail to attend a clinic for follow-up are more likely to be hypothyroid; in fact, they probably default owing to the inertia so characteristic of their defective thyroid function. Thus, an incomplete follow-up of patients may grossly underestimate the true incidence of hypothyroidism.

The delay in the control of thyrotoxicosis and the rising incidence of hypothyroidism after radioiodine have led to a reappraisal of the methods of treatment. Many suggestions for improvements in current techniques have been made. Some have little factual evidence to support them. It has been stated that an uneven distribution of ^{131}I in the thyroid is partly responsible for the subsequent hypothyroidism,¹² and prior antithyroid medication has been claimed to produce a more uniform distribution. In a few centres fractionation of the dose is favoured; repeated small doses are given over a few weeks. More recently transient suppression of function in one lobe by external irradiation before radioiodine therapy has been proposed.¹³ It is argued that the irradiated lobe will thus survive the ^{131}I -irradiation because uptake of the isotope will be temporarily suppressed. This technique poses additional problems of dosage, and the same criticism can be made of more involved methods of calculating the ^{131}I dose.¹⁴

Many authorities advocate repeated doses of radioiodine with rapid control of the thyrotoxicosis and acceptance of a high incidence of hypothyroidism. Where regular follow-up is not practicable large doses of radioiodine have been proposed followed by the immediate institution of full replacement doses of thyroxine, thus anticipating the almost inevitable hypothyroidism. With this approach one loses all opportunity of studying the problem further, but in some areas social and economic factors prevail and make this the best compromise. If the patients are not to suffer steps must be taken to ensure that the thyroxine will be taken regularly. There is no published evidence of the reliability of patients in this clinical situation, but experience in other circumstances is not reassuring.

More recently clinical evidence has been published showing that lower ^{131}I doses produce less hypothyroidism.⁵ Experimental evidence already existed showing that the radiation dose to the thyroid was the major factor in determining the number of thyroid cells which survived.^{1,3} A clinical trial of different doses of radioiodine with over 500 patients has shown a markedly lower incidence of hypothyroidism in a five-year period when smaller doses of ^{131}I were used.⁵ Three years after treatment the proportion of euthyroid patients was the same as in the group treated with the higher dose. This study employed a single dose of radioiodine, and persisting thyrotoxic symptoms were promptly controlled with antithyroid drugs. This medication was eventually discontinued when the patient had been maintained on a low dose for some time and further treatment appeared unnecessary. The use of antithyroid drugs

after lower doses of radioiodine offers a satisfactory programme of treatment for thyrotoxicosis. The delay in the control of symptoms is overcome and the use of lower doses of radioiodine greatly reduces the incidence of hypothyroidism.

Selection of Patients

Radioiodine therapy has usually been restricted to patients over 40 years of age. This decision arose from uncertainty about the side-effects of ^{131}I therapy, and particularly the effects upon the gonads. Apart from this restriction ^{131}I therapy is suitable for all patients. The advantages of radioiodine therapy are the ease of administration, which can be arranged on an outpatient basis, and the rarity of recurrence of thyrotoxicosis. Compared with thyroid surgery there is no risk to the recurrent laryngeal nerves, tetany never supervenes, and the general hazards of major surgery are avoided. This aspect is especially important in older patients. Careful follow-up of patients after thyroid surgery reveals an incidence of hypothyroidism varying from 5% to 25%, and there is a definite risk of recurrence. Antithyroid medication alone will control the patient's symptoms, but the relapse after a prolonged course of tablets is 60–80% in patients over 40 years of age.

Side-effects

Theoretically radiation from ^{131}I might induce leukaemia or thyroid cancer, but the available evidence does not support either possibility in adults. The incidence of leukaemia is the same in those patients who have had radioiodine therapy as in a population of the same age and sex distribution.^{15,16} Likewise, neoplasms do not appear to be induced in adults, although in children ^{131}I therapy has resulted in thyroid cancers.¹⁷ When antithyroid drugs are used after radioiodine therapy the dosage should be reduced to control the symptoms without producing hypothyroidism. There is experimental evidence in rats that high doses of antithyroid drugs after ^{131}I therapy can produce neoplastic changes in the thyroid.¹⁸ Presumably the excessive stimulation by thyroid-stimulating hormone, which occurs in hypothyroidism, is partially responsible for these changes in the irradiated thyroid cells. Thyrotoxic crises after radioiodine have been described but are rare. Early transient hypothyroidism is an uncommon feature—in contrast to the late-onset hypothyroidism which is permanent. It has been suggested that an autoimmune reaction in the thyroid might be initiated by the irradiation. Although there is a slight rise in the antibody titres it is not sustained and there is no evidence that this mechanism is related to the post- ^{131}I

hypothyroidism.¹⁹ Effects on other iodide-concentrating tissues have not been described, but the relatively low uptakes are unlikely to result in any significant radiation.

Conclusion

Radioiodine therapy offers a convenient and effective form of treatment for thyrotoxicosis. Experience has shown the two main disadvantages to be delay in the control of thyrotoxicosis and a rising incidence of hypothyroidism. Other side-effects are very rare. Experimental studies have led to a better understanding of the effects of radioiodine on the thyroid cell. These observations, coupled with long-term reviews of patients treated with radioiodine, have offered both an explanation and a possible solution to the problem of hypothyroidism after ^{131}I therapy. Lower doses of radioiodine will reduce the incidence of hypothyroidism. At the same time the proportion of euthyroid patients after a few years is not diminished and recurrence of thyrotoxicosis has not proved to be a problem. Lower doses of radioiodine delay control of hyperthyroidism, but the use of antithyroid drugs to suppress symptoms until the radioiodine has been effective is a rational approach and one which avoids repeated doses of radioiodine. Further clinical trials will reveal the optimum dose of radioiodine, which may well be lower than those already studied.

REFERENCES

- Al-Hindawi, A. Y., and Wilson, G. M., *Clin. Sci.*, 1965, 28, 555.
- Philp, J. R., *Postgrad. med. J.*, 1966, 42, 437.
- Greig, W. R., *Scot. med. J.*, 1966, 11, 307.
- Green, M., and Wilson, G. M., *Brit. med. J.*, 1964, 1, 1005.
- Smith, R. N., and Wilson, G. M., *ibid.*, 1967, 1, 129.
- Myhill, J., Reeve, T. S., and Figgis, P. M., *Amer. J. Roentgenol.*, 1965, 94, 828.
- Crooks, J., *Current Topics in Thyroid Research*, ed. C. Cassano and M. Andreoli, 1966, p. 1208. London.
- Beling, U., and Einhorn, J., *Acta radiol. (Stockh.)*, 1961, 56, 275.
- Dunn, J. T., and Chapman, E. M., *New Engl. J. Med.*, 1964, 271, 1037.
- McGirr, E. M., Thomson, J. A., and Murray, I. P. C., *Scot. med. J.*, 1964, 9, 505.
- Nofal, M. N., Beierwaltes, W. H., and Patno, M. E., *J. Amer. med. Ass.*, 1966, 197, 605.
- Fraser, R., Abbott, J. D., and Stewart, F. S., *Brit. J. Radiol.*, 1954, 27, 23.
- Trotter, W. R., *Current Topics in Thyroid Research*, ed. C. Cassano and M. Andreoli, 1966, p. 1159. London.
- Myhill, J. A., Hales, I. B., and Reeve, T. S., *ibid.*, 1966, p. 1143.
- Pochin, E. E., *Brit. med. J.*, 1960, 2, 1545.
- *Proc. roy. Soc. Med.*, 1964, 57, 564.
- Kogut, M. D., et al., *New Engl. J. Med.*, 1965, 272, 217.
- Doniach, I., *Brit. med. Bull.*, 1958, 14, 181.
- Einhorn, J., Fagraeus, A., and Jonsson, J., in *Current Topics in Thyroid Research*, ed. C. Cassano, and M. Andreoli, 1966, p. 1137. London.

ANY QUESTIONS?

We publish below a selection of questions and answers of general interest.

Familial Tendency in Cancer

Q.—What is the evidence, if any, that a tendency to cancer is familial?

A.—Cancer is not a single disease but a group of many different diseases with some features in common.

There is a strong familial tendency in the susceptibility to a few, mainly rare, types of cancer—e.g., cancer of the colon and rectum associated with familial polyposis, cancer of the central and peripheral nervous system associated with von Recklinghausen's neurofibromatosis multiplex, and a number of other rare hereditary syndromes, and cancer

of the skin associated with xeroderma pigmentosum. In the case of most of the commoner types of human cancer, however, a familial tendency, if it exists at all, is very weak. In this connexion particular attention has been directed towards cancer of the breast.¹

It is perhaps interesting to consider the question also from a theoretical point of view. Studies on laboratory animals strongly suggest that genetic factors affect both the spontaneous incidence of cancers of particular types and the ease with which they can be deliberately induced by exposure to carcinogenic agents. Multiple genes tend to be

involved in the case of each type of cancer. Such differences in susceptibility may be greatly magnified by intensive inbreeding (e.g., brother-sister mating for several generations).

Different races of man differ considerably in mortality from specific types of cancer, but many of these differences appear to be determined by environmental rather than genetic factors. Between individuals of any one race, provided that there is no inbreeding, genetically determined differences are unlikely to be obvious because of the multiplicity of genes involved and the kaleidoscopic-like heterogeneity of individuals.

Studies on experimental animals also tell us that it is possible to confuse the effects of vertically transmitted cancer viruses (i.e., cancer viruses passed from parent to offspring) with those genes. At present there