Correspondence

Letters to the Editor should not exceed 500 words.

Brucellosis Still Spreading R. W. D. Turner, F.R.C.P.; I. A. B. Cathie, F.R.C.P.	685
"Five-day Courses" and Respiratory Infections A. F. Foster-Carter, D.M.	685
Deaths from Appendicitis J. A. Shepherd, F.R.C.S.	686
Spinal Injuries J. J. Walsh, F.R.C.S., and others	686
Pregnancy and Chlormadinone Acetate Christine M. Butler, M.R.C.O.G.	686
Intraocular Lenses D. P. Choyce, F.R.C.S.	686
Unconscious Patient's Airway R. K. Gilbert, L.R.C.P., F.D.S. R.C.S.	687
Control of Rabies K. Kooros, M.D.	687
Diagnosing Parotid Calculi D. H. Patey, M.S.	687

Hiatus Hernia and Reflux Oesophagitis G. J. Cole, M.CHIR.	687
Treatment of Adder Bite S. V. Humphries, F.I.C.S.	688
Spironolactone in Hyperaldosteronism J. J. Brown, M.R.C.P., and others	688
Automation in the Laboratory F. W. C. Watts, F.I.M.L.T.	688
Fibrinolysis and Toxaemia of Pregnancy A. T. Coopland, F.R.C.S.(C.)	688
Unnecessary X-rays? R. M. Scott, M.B.	689
Vaccination against Whooping-cough A. H. Griffith, M.D.	689
Anaesthesia in the Dark J. D. Whitby, F.F.A. R.C.S.	689
Voluntary Sterilization P. H. Addison, M.R.C.S.	689
Care of Myocardial Infarction J. Fry, F.R.C.S.	690

Mutants of Influenza Virus J. Mills, M.D., and others	690
Treatment of Exacerbations of Chronic Bronchitis I. W. B. Grant, F.R.C.P.ED.	690
Fascioliasis Yet Again E. W. Hardman, M.B., and others	690
"Proplis" J. M. Crawford, M.D.	690
Plight of Commonwealth Graduates in U.K. C. H. Hodge, F.R.C.S.ED.	690
Superannuation in the N.H.S. M. R. McNulty, F.F.R. R.C.S.I., and others	691
G.M.C.'s Annual Retention Fee I. M. Librach, M.B.; J. H. Reading, M.B.	691
The Consultant's Job J. N. Rimmer, F.R.C.S.	691
Salaries of Physiotherapists P. H. Corkery, F.R.C.S.ED.	691
Hunting the Caucas M. O'Donnell, M.B.	691

Brucellosis Still Spreading

SIR,—Dr. R. J. Henderson's article on brucellosis (29 November, p. 550) and your leading article (p. 512) prompt me to report a death from brucellosis which recently occurred in Edinburgh and was due to the drinking of infected milk in a town in Fife. The patient, a man aged 42, had been treated by mitral valvotomy seven years previously, and remained well until contracting this lethal infection. A neighbour who drank milk from the same dairy suffered a similar infection but recovered. It is not known how many others were infected.

We were astonished to learn that there is no law against selling infected milk, nor can anything be done to prevent the farmer distributing milk known to be infected. All that can be done in the words of the medical officer of health of the district concerned is to "twist his arm." Brucellosis is certainly one of the most serious forms of endocarditis. Though death may be rare, it is a likely complication in anyone with heart disease, and, as you point out, 1,000 preventable cases occur in this country each year, and in this respect we are indeed a backward land.—I am, etc.,

RICHARD TURNER.

Department of Medicine,
University of Edinburgh.

SIR,—May I thank Dr. R. J. Henderson (29 November, p. 550) and your leader writer (p. 512) for drawing attention to the dangers and problems of brucellosis? The frustrations and expense to the farmer in its eradication are perhaps underlined by my own experience.

Every animal on my farm, Guernseys and Herefords, has been home-bred with the exception of the bulls, every female has been vaccinated with strain 19 against brucellosis, and there has not been a case of clinical contagious abortion for 25 years; so I

thought that entering the Ministry of Agriculture's eradication scheme would be easy. Three milk ring tests on the Guernseys were negative, but the last of the 100-odd Herefords to be bled for evidence of infection was reported as a "roaring positive." This was a 16-year-old animal born and bred on the farm, duly vaccinated, which had had 12 healthy calves and was now penned with another old cow being fattened for slaughter. They were both sent forthwith.

After 60 days the Herefords were bled again, but whereas the first time the bleeding was free, this time I had to pay my private vet; and this time a heifer which had just had a second healthy calf and had been previously negative, which had had no contact with the first reactor for a year, was positive. She was immediately slaughtered. After the prescribed period I paid for a third bleeding which showed the herd to be clear, as were the Guernsey bulls. The whole procedure has taken about two years and many man hours.

Apart from making me wonder about positive serology in healthy animals, this has been a very unprofitable venture for the farm. If one is going to make a genuine contribution to the public health from one's own pocket the Ministry might be a little more co-operative with the services of its vets, while the question of compensation for the voluntary slaughter of positively reacting but healthy animals should be seriously considered.

We managed virtually to eradicate tuberculous from the national milk herd with the help of a bonus or incentive scheme and compensation, and only an approach more positive than that at present in operation can be expected to diminish the problems of brucellosis which Dr. Henderson so rightly highlights.—I am, etc.,

Moreton-in-Marsh,
Glos.

I. A. B. CATHIE.

"Five-day Courses" and Respiratory Infections

SIR,—“A cause of common Errors is the Credulity of men, that is, an easie assent to what is obtruded, or a believing at first ear, what is delivered by others.” Thus wrote Sir Thomas Browne in 1672 (*Pseudodoxia Epidemica*, Chapter 5). Yet such errors continue to appear even in our scientific age, and one of the latest and most mischievous is the belief that a “five-day course” of an antibiotic is a panacea for respiratory infections.

I drew attention to this last year in your columns (7 September, 1968, p. 614) and make no apology for doing so again at the onset of another winter. Before long we will be admitting bronchitic patients who have had several “courses” of various antibiotics over a period of weeks, each followed by a relapse; before finally, in desperation, the puzzled doctor sends the patient into hospital.

As with most common errors, the origin of this one is obscure, but it is constantly being reinforced by careless and uncritical repetition, often by people who should know better, and therefore it is unfair to blame the average doctor for accepting its validity. Already it has crept into the *British National Formulary*¹ and a particularly striking example occurs in a recent paper by Dr. R. N. Johnson and others on the chemoprophylaxis of chronic bronchitis (1 November, p. 265). These authors mention that “all groups received a five-day course of tetracycline for any acute exacerbation” (my italics). This statement appears in the Summary which heads the paper and which (let us face it) is the only part the majority of readers are likely to study. The busy doctor reading this and then glancing at the rest of the article will put it down with the comfortable feeling that these clever, scientific chaps are approving his habit of giving five days' supply of tetracycline to patients who have an exacerbation

of bronchitis. But in fact the authors produce no justification for this practice.

If we consider objectively what we are trying to do, it becomes obvious that to treat acute bronchitis, or an exacerbation of chronic bronchitis, by giving a set "course" of antibiotic, whether of five or any other number of days, is completely illogical. Our object is to eliminate the infection; therefore the antibiotic must not be stopped until there is good evidence that this has been achieved, and it is impossible to predict how long this will take. The chosen drug should be prescribed in adequate dosage for a reasonable time, say one week; but the doctor must arrange to see the patient again at the end of this period. He can then decide on clinical grounds (control of symptoms, fever, and character of sputum) whether to maintain, reduce or discontinue the drug. Concerning dosage; recent work suggests that the customary 250 mg. q.d.s. of tetracycline or ampicillin is often insufficient in the early stages of an infection, and that double this dose should be given.

If we could forget the "five-day course" and get back to these basic principles (adequate dosage of antibiotic and observation until the desired effect has been obtained), there is no doubt that many patients would be spared a great deal of prolonged illness and hospital treatment.—I am, etc.,

A. F. FOSTER-CARTER.

Brompton Hospital,
Frimley Branch,
Surrey.

REFERENCE

1. *British National Formulary*, p. 93. London, B.M.A. and the Pharmaceutical Society of Great Britain.

Deaths from Appendicitis

SIR,—In congratulating Dr. H. G. Pledger and Mr. R. Buchan on their informative, convincing, and timely paper on deaths in children with acute appendicitis (22 November, p. 466) I would like to make three comments.

Firstly, I believe that the figures they derive from their analysis of the Registrar General's statistics are probably an underestimate. The accuracy of death certification leaves much to be desired, whether in domiciliary or hospital practice. I feel that if search was made under some additional code numbers—for example, 576 (general peritonitis), the total annual mortality rate from acute appendicitis in children would prove greater than they report.

Secondly, although delay in diagnosis may explain only a small proportion of deaths in the series investigated, this factor remains significant. I have stressed the point elsewhere¹ that in most series of children under the age of 5 with acute appendicitis nearly 50% are found to have already perforated by the time operation is done. In the paper under discussion the very high incidence of "appendicitis with peritonitis" emphasizes the problem. Unfortunately it appears that in small children appendicitis progresses very rapidly and that prior to perforation the signs and symptoms may be extremely difficult to assess. Whether this has relation to the high proportion of lymphoid tissue in the child's appendix or to some other factor, perhaps viral infection, remains unknown. Without new developments in technique in diagnosis "good training and a constant suspicion of acute appendicitis" cannot produce more than marginal improvement.

Finally, although it is hinted that there is a considerable difference in the mortality rates according to the type of hospital, it is perhaps a pity that more definite figures could not be given. If, as seems all too likely, the mortality rates are high in relatively small hospitals where children with acute abdominal conditions are managed only sporadically, and if, in contrast, results are better where it is possible to have the benefits of skilled paediatric care, medical and nursing, then this supports the implications of the recent Bonham-Carter Report.² Emergency surgery on small children is best carried out either in large special hospitals for children or in district hospitals where there is a viable paediatric unit, which can offer all the assistance necessary in these difficult cases.—I am, etc.,

Liverpool 1.

JOHN SHEPHERD.

REFERENCES

1. Shepherd, J. A., *Surgery of the Acute Abdomen*, 1968, 2nd ed., p. 245. Edinburgh, Livingstone.
2. Central Health Services Council, *The Functions of the District Hospital*, 1969. London, H.M.S.O.

Spinal Injuries

SIR,—In his article on spinal injuries and fractures (15 November, p. 414) Mr. J. Ashworth rightly draws attention to the importance of careful first aid treatment. However, he makes certain statements which we feel should not pass without comment.

He states that "Injuries to the spine may be classified as stable or unstable." We feel that this classification is too simplified and that varied degrees of stability may be encountered. Immediately afterwards he comments that "it is invariably in the unstable injuries that damage to the cord occurs." This is not confirmed by our experience because, not infrequently, permanent paralysis is caused by very simple crush fractures, and indeed in some cases of traumatic paraplegia, particularly in children, there is no x-ray evidence of bony injury or dislocation.

However, the statement to which the greatest exception must be taken is that under the heading of "Treatment," where Mr. Ashworth claims that "With the exception of injuries in the cervical region, any neurological damage usually occurs at the time of impact. Hence there is nothing that can be done at the site of the accident that will have any effect on the ultimate prognosis." This is incorrect and misleading, and could well result in serious mismanagement of patients by ambulance personnel and others who may read the article. In our own experience it is not unusual to obtain a clear history of the onset of the neurological damage during transport, or indeed sometimes in hospital, and this may occur with dorsal and lumbar injuries as well as with cervical injuries. A recent example was a woman with a mid-thoracic fracture without neurological damage, who was treated by being sat upright in bed and within some minutes developed a paraplegia at T.5 (albeit sensory incomplete), which failed to recover.—We are, etc.,

J. J. WALSH.

H. L. FRANKEL.

G. M. HYSLOP.

D. O. HANCOCK.

National Spinal Injuries Centre,
Stoke Mandeville Hospital,
Aylesbury, Bucks.

Pregnancy and Chlormadinone Acetate

SIR,—I should like to comment on the advertisement for Verton (chlormadinone acetate 0.5 mg.) (22 November, pp. xix and xx). I am a little concerned that the risk of pregnancy should be more clearly stated. Verton is said to be "more reliable than mechanical methods of contraception," and doctors are requested to report any pregnancies occurring to the Dunlop Committee on Safety of Drugs. It is also suggested that Verton "is useful for all those who are unreliable at taking their tablets because of the simplicity of everyday medication."

Perhaps potential prescribers would be interested to know of the latest results in the trial of chlormadinone acetate 0.5 mg. organized by the Council for the Investigation of Fertility Control at various centres throughout this country. Two hundred and twenty-three women have been enrolled and 2,518 cycles have been completed. Twenty pregnancies have occurred, and of these only five women admitted omitting tablets. One of these women omitted the 12th, 14th, and 15th tablets in the cycle and conception followed.

When pregnancy is definitely contraindicated for medical or social reasons I would suggest that this type of contraception is not the method of choice.—I am, etc.,

CHRISTINE BUTLER,

London W.1.

Medical Assistant,
Council for the Investigation of
Fertility Control.

Intraocular Lenses

SIR,—In his typically erudite and witty account of "Spectacles in Evolution" (*The Times*, 4 November, p. 16), Mr. P. D. Trevor-Roper said: "The correcting lens can even be slipped into the eyeball itself, although the operation has generally been abandoned because the eye is often damaged irreparably in the process."

My distinguished colleague is entitled to his views, but they are not necessarily those of the rest of the ophthalmic profession. Intraocular lenses came into being about 20 years ago, with the posterior chamber lens used by Harold Ridley for the correction of aphakia. Then followed a period of frenzied activity, in which many different types of intraocular lens implants were used and many complications arose. Understandably, public opinion went against these devices and Mr. Trevor-Roper's remark, if made 10 years ago, would probably have been justified. But since then there have been various developments which suggest that such a view is now out of date. For some years now there has existed the "Intra-Ocular Implant Club," an association of like-minded eye surgeons in many different countries who regularly use such lens implants in the best interests of their patients. There are at least two British surgeons each with a personal series approaching the 1,000 mark. One of the principal implant manufacturers is British; they regularly dispatch substantial numbers to various parts of the world; they report a steady increase in demand for them.

As recently as June last the section of ophthalmology of the Royal Society of Medicine devoted an entire session to the consideration of the late results of intraocular implants. Patients (some with