

## Analgesic effect of breast feeding in term neonates: randomised controlled trial

Ricardo Carbajal, Soocramanien Veerapen, Sophie Couderc, Myriam Jugie, Yves Ville

### Abstract

**Objectives** To investigate whether breast feeding is effective for pain relief during venepuncture in term neonates and compare any effect with that of oral glucose combined with a pacifier.

**Design** Randomised controlled trial.

**Participants** 180 term newborn infants undergoing venepuncture; 45 in each group.

**Interventions** During venepuncture infants were either breast fed (group 1), held in their mother's arms without breast feeding (group 2), given 1 ml of sterile water as placebo (group 3), or given 1 ml of 30% glucose followed by pacifier (group 4). Video recordings of the procedure were assessed by two observers blinded to the purpose of the study.

**Main outcome measures** Pain related behaviours evaluated with two acute pain rating scales: the Douleur Aiguë Nouveau-né scale (range 0 to 10) and the premature infant pain profile scale (range 0 to 18).

**Results** Median pain scores (interquartile range) for breast feeding, held in mother's arms, placebo, and 30% glucose plus pacifier groups were 1 (0-3), 10 (8.5-10), 10 (7.5-10), and 3 (0-5) with the Douleur Aiguë Nouveau-né scale and 4.5 (2.25-8), 13 (10.5-15), 12 (9-13), and 4 (1-6) with the premature infant pain profile scale. Analysis of variance showed significantly different median pain scores ( $P < 0.0001$ ) among the groups. There were significant reductions in both scores for the breast feeding and glucose plus pacifier groups compared with the other two groups ( $P < 0.0001$ , two tailed Mann-Whitney U tests between groups). The difference in Douleur Aiguë Nouveau-né scores between breast feeding and glucose plus pacifier groups was not significant ( $P = 0.168$ ).

**Conclusions** Breast feeding effectively reduces response to pain during minor invasive procedure in term neonates.

### Introduction

In healthy infants, the most common painful procedures are heel lance and venepuncture. Pharmacological treatments are rarely used during these procedures because of concerns about their effectiveness and potential adverse effects. Therefore, non-pharmacological interventions are valuable alternatives.

Recent studies have reported that pain can be reduced with simple interventions such as sweet oral

solutions and non-nutritive sucking<sup>1-4</sup> or multisensory stimulation.<sup>5</sup> Skin to skin contact between mothers and infants can also be effective.<sup>6</sup> Environmental and behavioural strategies have been considered essential to the prevention and management of neonatal pain.<sup>7</sup> As breast feeding probably constitutes the most potent pleasant stimulation a newborn infant can experience we hypothesised that breast feeding could have analgesic properties in neonates.

We investigated the efficacy of breast feeding for pain relief during venepuncture in term neonates and compared any effect with that of oral glucose combined with a pacifier.

### Methods

#### Protocol

We included infants who were born at  $\geq 37$  weeks' gestation; had APGAR scores  $\geq 7$  at 5 minutes; were aged  $\geq 24$  hours; were undergoing venepuncture as part of routine medical care; were breast fed; and had not been fed for the previous 30 minutes. We excluded infants with medical instability, those who had received naloxone during the previous 24 hours, and those who had received a sedative or a major analgesic during the previous 48 hours.

#### Procedures and masking

Participating infants and their mothers were taken to a quiet nursery room for venepuncture. SV opened a consecutively numbered envelope, which contained the treatment assigned to each infant. Infants were allocated to one of four groups: in group 1 they were breast fed, starting two minutes before the procedure and continuing throughout; in group 2 they were held in their mother's arms without breast feeding, starting two minutes before the procedure; in group 3 two minutes before the procedure infants were laid on a table and given 1 ml of placebo (sterile water) without a pacifier; and in group 4 two minutes before the procedure infants were laid on a table and given 1 ml of 30% glucose followed by sucking a pacifier. The infant's legs and feet were uncovered to allow observation of movements. Infants in groups 3 and 4 lay supine on an examination table during procedures.

The water or 30% glucose was administered for about 15 seconds by a sterile syringe into the infant's mouth. In group 4 the pacifier (standard nipple stuffed with a gauze square for resistance) was held gently in



This is an abridged version; the full version is on [bmj.com](http://bmj.com)

Department of Paediatrics, Poissy-Saint Germain Hospital, 78300 Poissy, France

Ricardo Carbajal  
*paediatrician*  
Myriam Jugie  
*paediatrician*

Department of Gynaecology and Obstetrics, Poissy-Saint Germain Hospital Soocramanien Veerapen  
*general practitioner*  
Yves Ville  
*professor*

Maternity Ward, Poissy-Saint Germain Hospital Sophie Couderc  
*paediatrician*

Correspondence to: R Carbajal [carbajal@club-internet.fr](mailto:carbajal@club-internet.fr)

BMJ 2003;326:13-5

**Table 1** Median pain scores (MPS) among four groups of neonates with Douleur Aiguë Nouveau-né scale\* during venepuncture. Figures are estimated median difference (95% confidence interval)

	Breast feeding (MPS=1)	Placebo (MPS=10)
Mother's arms (MPS=10)	7 (7 to 8), P<0.0001†	0 (0 to 0), P=1†
Placebo-sterile water (MPS=10)	7 (6 to 8), P<0.0001†	NA
30% glucose plus pacifier (MPS=3)	0 (0 to 2), P=0.16†	6 (5 to 7), P<0.0001†

NA=not applicable. \*0 (no pain) to 10 (maximum pain). †For two sided Mann-Whitney U test.

**Table 2** Comparisons of median pain scores (MPS) among four groups of neonates with premature infant pain profile scale\* during venepuncture. Figures are estimated median difference (95% confidence interval)

	Breast feeding (MPS=4.5)	Placebo (MPS=12)
Mother's arms (MPS=13)	8 (6 to 9), P<0.0001†	1 (0 to 3), P=0.38†
Placebo-sterile water (MPS=12)	7 (5 to 8), P<0.0001†	NA
30% glucose plus pacifier (MPS=4)	1 (-1 to 2), P=0.28†	8 (6 to 9), P<0.0001†

NA=not applicable. \*0 (no pain) to 18 (maximum pain). †For two sided Mann-Whitney U test.

the infant's mouth by an assistant throughout the procedure. Infants' heart rate and oxygen saturation were monitored with a Nellcor monitor (model N-395). The infants and the monitor screen were video recorded during the procedure. Venepuncture was performed on the dorsal aspect of the infant's hand by one of three experienced nurses.

Two specially trained observers independently assessed the recordings using the Douleur Aiguë Nouveau-né (DAN) scale (primary outcome measure) and the premature infant pain profile (PIPP) scale (secondary outcome measure). They assessed arousal state using Precht's observational rating system.<sup>8</sup> Assessment of pain started when the needle was inserted and ended when it was removed. Observers were blinded to the purpose and hypothesis of the study. They had been told that we were assessing agreement of their scores in four different situations. For the DAN scale there was good agreement between both observers on initial evaluation. The two observers independently re-evaluated all the procedures for which scores had not been identical during their first assessment, and obtained perfect interobserver agreement.

#### Sample calculation

We calculated that we would need 40 infants in each group to detect a 2 point difference in DAN scale with 80% power and at 1% significance. We decided to include 45 neonates in each group to cover potential problems with video recordings.

#### Pain scales

The DAN scale is a behavioural scale developed to rate acute pain in term and preterm neonates.<sup>9</sup> Scores range from 0 (no pain) to 10 (maximum pain). It evaluates three items: facial expressions, limb movements, and vocal expression.

The PIPP scale is a multidimensional measure developed to assess acute pain in preterm and term infants.<sup>10</sup> It measures gestational age, behavioural state, heart rate, oxygen saturation, and three facial reactions (brow bulge, eye squeeze, nasolabial furrow). In term infants, scores range from 0 (no pain) to 18 (maximum pain).

#### Statistical analysis

We used one way analysis of variance on ranks to compare overall differences among four groups. We compared median pain scores of all groups using two

tailed Mann-Whitney U tests. Because five pairwise planned comparisons were made we considered  $P < 0.01$  as significant. We used  $\chi^2$  tests to compare categorical variables.

## Results

During the study period (February to June 2001) 351 infants met the inclusion criteria. Of these, 180 were allocated to one of four equal sized groups. The perinatal characteristics of neonates not included in the study were similar to those included. There were no substantial differences among the groups except for arousal state—the median score for state of arousal was lower in the breast fed group. The reasons for venepuncture included tests for hypothyroidism and phenylketonuria screening, bilirubin, C reactive protein, sickle cell disease screening, calcium, and blood typing.

The median pain scores (interquartile range) during venepunctures for group 1 (breast feeding), group 2 (mother's arms), group 3 (placebo), and group 4 (30% glucose plus pacifier) were 1 (0-3), 10 (8.5-10), 10 (7.5-10), and 3 (0-5) with the DAN scale and 4.5 (2.25-8), 13 (10.5-15), 12 (9-13), and 4 (1-6) with the PIPP scale. Analysis of variance showed that median pain scores were significantly different ( $P < 0.0001$ ). Tables 1 and 2 show pairwise comparisons of median pain scores.

## Discussion

We have shown that breast feeding throughout a painful procedure is analgesic in term neonates. Of 44 infants in the breastfeeding group, 16 showed no indication at all that the venepuncture and blood sampling had even occurred; 35 had a DAN pain score  $\leq 3$ , which can be considered as reflecting minimal or no pain. Our findings are clinically important as they show that natural protective mechanisms may safely and non-invasively be activated by breast feeding during medical procedures.

We detected no reduction in response to pain in infants who were simply held in their mother's arms, possibly because these infants were dressed and did not have a skin to skin contact with their mothers. Gray et al found that 10 to 15 minutes' skin to skin contact between a mother and baby reduces the infant's response to pain during heel stick.<sup>6</sup> To our knowledge, there have been only two previous reports on the analgesic effect of breast feeding. Bilgen et al compared the analgesic effects of sucrose, expressed breast milk, and breast feeding during heel pricks. Breast feeding was allowed for two minutes and stopped before a heel prick.<sup>11</sup> There was no analgesic effect of this type of intervention, possibly because breast feeding was not continued during the procedure. Gray et al reported that breast feeding before, during, and after heel prick markedly reduced crying and grimacing and prevented the increase in heart rate in term neonates compared with swaddled infants in their cots.<sup>12</sup> No other groups were included in their study design. The infants were held in full body skin to skin contact during the entire procedure.

#### Study limitations

Firstly, observers obviously recognised the four groups when they were evaluating the recordings. However,

### What is already known on this topic

Current pharmacological treatments are not appropriate for pain relief during minor procedures like venepuncture or heel prick in newborn infants

Oral sweet solutions, non-nutritive sucking, and skin to skin contact reduce procedural pain in newborn infants

### What this study adds

Breast feeding during a painful procedure effectively reduces the response to pain in newborn infants

The analgesic properties of breast feeding are at least as potent as the combination of sweet solutions and a pacifier

they did not know the purpose of the study. Moreover, high agreement among observers during initial evaluations indicates objectivity. Secondly, although the DAN scale has been shown to discriminate pain in term newborn infants, no study has yet proved that it can grade the degree of perception of pain. We assumed that the more pronounced the facial expressions, limb movements, and vocal expressions, the higher the pain in the infant. Nevertheless, the robustness of pain evaluation was supported by the fact that the simultaneous use of the PIPP scale yielded similar results. Finally, median score for state of arousal was lower in the breastfeeding groups than in the other

groups. This difference was slight and in our opinion was insufficient to explain all differences observed in pain scores among groups.

We thank the nursing staff of the maternity ward of the Poissy Hospital for their help during the study. We also thank Nicolas Simon, chief of the emergency department of the Poissy Hospital, for reading the manuscript. We are indebted to the parents for allowing their infants to participate in the study.

Contributors: See bmj.com

Funding: Fondation pour la Santé CNP, France.

Competing interests: None declared.

- 1 Stevens B, Taddio A, Ohlsson A, Einarson T. Sucrose for analgesia in newborn infants undergoing painful procedures. *Cochrane Database Syst Rev* 2001;(4):CD001069.
- 2 Skogsdal Y, Eriksson M, Schollin J. Analgesia in newborns given oral glucose. *Acta Paediatr* 1997;86:217-20.
- 3 Carbajal R, Chauvet X, Couderc S, Olivier-Martin M. Randomised trial of analgesic effects of sucrose, glucose, and pacifiers in term neonates. *BMJ* 1999;319:1393-7.
- 4 Blass EM, Watt LB. Suckling- and sucrose-induced analgesia in human newborns. *Pain* 1999;83:611-23.
- 5 Bellieni CV, Bagnoli F, Perrone S, Nenci A, Cordelli DM, Fusi M, et al. Effect of multisensory stimulation on analgesia in term neonates: a randomized controlled trial. *Pediatr Res* 2002;51:460-3.
- 6 Gray L, Watt L, Blass EM. Skin-to-skin contact is analgesic in healthy newborns. *Pediatrics* 2000;105(1):E14.
- 7 Franck L, Lawhon G. Environmental and behavioral strategies to prevent and manage neonatal pain. In: Anand KJS, Stevens BJ, McGrath PJ, eds. *Pain in neonates*. 2nd rev ed. Amsterdam: Elsevier, 2000:203-16.
- 8 Prechtl HFR. The neurological examination of the full term newborn infant. In: Prechtl HFR, ed. *Clinics in developmental medicine*. 2nd rev ed. London: Heinemann, 1977.
- 9 Carbajal R, Paupe A, Hoenn E, Lenclen R, Olivier-Martin M. Douleur Aiguë Nouveau-né: une échelle comportementale d'évaluation de la douleur aiguë du nouveau-né. [APH: evaluation behavioural scale of acute pain in newborn infants.] *Arch Pediatr* 1997;4:623-8.
- 10 Stevens B, Johnston C, Petryshen P, Taddio A. Premature infant pain profile: development and initial validation. *Clin J Pain* 1996;12:13-22.
- 11 Bilgen H, Ozek E, Cebeci D, Ors R. Comparison of sucrose, expressed breast milk, and breast-feeding on the neonatal responses to heel prick. *J Pain* 2001;2:301-5.
- 12 Gray L, Miller LW, Philipp BL, Blass EM. Breastfeeding is analgesic in healthy newborns. *Pediatrics* 2002;109:590-3. (Accepted 17 October 2002)

## Sexual behaviour of adolescents in Nigeria: cross sectional survey of secondary school students

Gail B Slap, Lucy Lot, Bin Huang, Comfort A Daniyam, Therese M Zink, Paul A Succop

### Abstract

**Objectives** To determine whether family structure (polygamous or monogamous) is associated with sexual activity among school students in Nigeria.

**Design** Cross sectional school survey with a two stage, clustered sampling design.

**Participants** 4218 students aged 12-21 years attending 39 schools in Plateau state, Nigeria. Responses from 2705 students were included in the analysis.

**Main outcome measure** Report of ever having had sexual intercourse. Variables of interest included sexual history, age, sex, religion, family polygamy, educational level of parents, having a dead parent, and sense of connectedness to parents and school.

**Results** Overall 909 students (34%) reported ever having had sexual intercourse, and 1119 (41%) reported a polygamous family structure. Sexual activity was more common among students from polygamous families (42% of students) than

monogamous families (28%) ( $\chi^2=64.23$ ;  $P<0.0001$ ). Variables independently associated with sexual activity were male sex (adjusted odds ratio 2.52 (95% confidence interval 2.05 to 3.12)), older age (1.62 (1.24 to 2.14)), lower sense of connectedness with parents (1.87 (1.48 to 2.38)), having a dead parent (1.59 (1.27 to 2.00)), family polygamy (1.58 (1.29 to 1.92)), lower sense of connectedness with school (1.25 (1.09 to 1.44)), and lower educational level of parents (1.14 (1.05 to 1.24)). Multistep logistic regression analysis showed that the effect of polygamy on sexual activity was reduced by 27% by whether students were married and 22% by a history of forced sex.

**Conclusions** Secondary school students in Nigeria from a polygamous family structure are more likely to have engaged in sexual activity than students from a monogamous family structure. This effect is partly explained by a higher likelihood of marriage during adolescence and forced sex. Students' sense of connectedness to their parents and school, regardless of family structure, decreases the likelihood of sexual

ELPS

This is an abridged version; the full version is on bmj.com

Division of Adolescent Medicine, University of Cincinnati College of Medicine, Cincinnati Children's Hospital Medical Center (ML-4000), 3333 Burnet Avenue, Cincinnati, Ohio, United States 45229-3039

Gail B Slap  
Rauh professor of paediatrics and internal medicine  
Lucy Lot  
fellow in adolescent medicine  
Bin Huang  
assistant professor of paediatrics

continued over

BMJ 2003;326:15-8