

community contraception clinics. General practitioners could also be trained.

Our study provides high quality evidence that investing in training and support for practice nurse led partner notification would be an effective use of government resources committed to improving sexual health.

We thank Philippa Matthews, general practitioner principal at Lee Bank Group Practice, Birmingham, who designed, coordinated, and delivered the training package for practice nurses; Anna Graham, who helped deliver the training; the practice nurses who participated in the chlamydia screening studies project and in this trial; and Deborah Hawking for secretarial support throughout the chlamydia screening studies project.

Contributors: See bmj.com.

Funding: The chlamydia screening studies project was funded through the NHS research and development HTA programme (project No 97/32/31). NL was supported by a NHS career scientist award until 31 March 2005. JM is supported by a NHS career scientist award.

Competing interests: None declared.

Ethical approval: South West multicentre research ethics committee (MREC/00/6/30) and all relevant local research ethics committees.

- 1 WHO/joint United Nations programme on HIV/AIDS. *Sexually transmitted diseases: policies and principles for prevention and care*. Geneva: UNAIDS, 1999. [UNAIDS best practice collection.]
- 2 Health Protection Agency. *All new episodes seen at GUM clinics: 1999-2003. Country specific tables*. London: HPA, 2004.
- 3 Adler M. Sexual health. *BMJ* 2003;327:62.

- 4 Stokes T, Schober P. A survey of contact tracing practice for sexually transmitted diseases in GUM clinics in England and Wales. *Int J STD AIDS* 1999;10:17-21.
- 5 Low N, McCarthy A, Macleod J, Salisbury C, Horner PJ, Roberts TE, et al. The chlamydia screening studies: rationale and design. *Sex Transm Infect* 2004;80:342-8.
- 6 Macleod J, Salisbury C, Low N, McCarthy A, Sterne JA, Holloway A, et al. Coverage and uptake of systematic postal screening for genital Chlamydia trachomatis and prevalence of infection in the United Kingdom general population: cross sectional study. *BMJ* 2005;330:940.
- 7 LaMontagne DS, Baster K, Emmett L. *Determinants of chlamydia re-infection: the role of partner change and treatment*. Abstract TO-405. 16th Biennial meeting of the International Society for Sexually Transmitted Diseases Research. Amsterdam: 10-13 Jul, 2005.
- 8 Nicolai LM, Ickovics JR, Zeller K, Kershaw TS, Milan S, Lewis JB, et al. Knowledge of sex partner treatment for past bacterial STI and risk of current STI. *Sex Transm Infect* 2005;81:271-5.
- 9 Curtis L, Netten A. *Units costs of health and social care*. Canterbury: University of Kent at Canterbury, Personal Social Services Research Unit, 2003.
- 10 Ross JDC, Sutherland S, Coia J. Genital Chlamydia trachomatis infections in primary care. *BMJ* 1996;313:1192-3.
- 11 Mathews C, Coetzee N, Zwarenstein M, Lombard C, Gutmacher S, Oxman A, et al. Strategies for partner notification for sexually transmitted diseases. *Cochrane Database Syst Rev* 2001;(4):CD002843.
- 12 LaMontagne DS, Fenton KA, Randall S, Anderson S, Carter P, on behalf of the National Chlamydia Screening Steering Group. Establishing the national chlamydia screening programme in England: results from the first full year of screening. *Sex Transm Infect* 2004;80:335-41.
- 13 Adams EJ, LaMontagne DS, Johnston AR, Pimenta JM, Fenton KA, Edmunds WJ. Modelling the healthcare costs of an opportunistic chlamydia screening programme. *Sex Transm Infect* 2004;80:363-70.
- 14 Golden MR, Whittington WL, Handsfield HH, Hughes JP, Stamm WE, Hogben M, et al. Effect of expedited treatment of sex partners on recurrent or persistent gonorrhoea or chlamydial infection. *N Engl J Med* 2005;352:676-85.
- 15 Department of Health. *Chlamydia screening programme roll out. Core requirements*. London: DoH, 2003.

(Accepted 1 November 2005)

doi 10.1136/bmj.38678.405370.7C

Use of a dummy (pacifier) during sleep and risk of sudden infant death syndrome (SIDS): population based case-control study

De-Kun Li, Marian Willinger, Diana B Petitti, Roxana Odouli, Liyan Liu, Howard J Hoffman

Division of Research, Kaiser Permanente Northern California, 2000 Broadway, Oakland, CA 94612, USA

De-Kun Li
senior research scientist

Liyan Liu
programmer analyst
Roxana Odouli
research associate

Research and Evaluation, Kaiser Permanente Southern California, Pasadena, CA 91188, USA

Diana B Petitti
senior scientific adviser

continued over

BMJ 2006;332:18-21

Abstract

Objectives To examine the association between use of a dummy (pacifier) during sleep and the risk of sudden infant death syndrome (SIDS) in relation to other risk factors.

Design Population based case-control study.

Setting Eleven counties in California.

Participants Mothers or carers of 185 infants whose deaths were attributed to SIDS and 312 randomly selected controls matched for race or ethnicity and age.

Main outcome measure Use of a dummy during sleep determined through interviews.

Results The adjusted odds ratio for SIDS associated with using a dummy during the last sleep was 0.08 (95% confidence interval 0.03 to 0.21). Use was associated with a reduction in risk in every category of sociodemographic characteristics and risk factors examined. The reduced risk associated with use seemed to be greater with adverse sleep conditions (such as sleeping prone or on side and sleeping with a mother who smoked), although the observed interactions were not significant. For example, infants

who did not use a dummy and slept prone or on their sides (*v* on their back) had an increased risk of SIDS (2.61, 1.56 to 4.38). In infants who used dummies, there was no increased risk associated with sleeping position (0.66, 0.12 to 3.59). While cosleeping with a mother who smoked was also associated with increased risk of SIDS among infants who did not use a dummy (4.5, 1.3 to 15.1), there was no such association among those who did (1.1, 0.1 to 13.4). **Conclusions** Use of a dummy seems to reduce the risk of SIDS and possibly reduces the influence of known risk factors in the sleep environment.

Introduction

The reduction in the incidence of sudden infant death syndrome (SIDS) after the Back to Sleep campaign in the United States and other countries showed that sleep environment strongly influences the risk of SIDS,^{1 2} although the underlying mechanisms remain



This is the abridged version of an article that was posted on bmj.com on 9 December 2005: <http://bmj.com/cgi/doi/10.1136/bmj.38671.640475.55>

poorly understood. Despite the considerable reduction in the incidence of SIDS after this public health intervention, it remains one of the leading causes of infant mortality. Some studies have reported that use of a dummy (pacifier) is associated with a reduced risk of SIDS, though few have examined the association in detail and in the context of other risk factors.³⁻⁷

Dummies usually have a bulky external handle, which could alter the infant's sleep environment by changing the configuration of the airway passage surrounding the nose and mouth. Sucking on a dummy may enhance the development of neural pathways that control the potency of the upper airway.^{8,9}

We examined data collected in a population based case-control study of risk factors for SIDS in California to determine whether use of a dummy during sleep is associated with a reduced risk of SIDS, what factors may modify the association between use and SIDS, and whether use influences other risk factors related to sleep environment.

Methods

We conducted a population based case-control study in 10 counties of northern California and Los Angeles County in southern California in 1997-2000. We identified cases of SIDS that were reported to the California Department of Health Services and to the coroner's office in Los Angeles County during the study period. Eligible participants were the mothers of any infants with a confirmed diagnosis of SIDS who lived in one of the participating counties and spoke either English or Spanish. A total of 185 mothers (50% of those eligible) participated in the study (see bmj.com).

Controls

We used data from birth certificates to identify infants randomly selected from all eligible births in the same county in which the dead infant had lived, matched for maternal race/ethnicity. Age at interview for controls was matched to age (plus or minus two weeks) at death for cases; and all mothers were asked about the index sleep, which for mothers of dead infants was defined as the last sleep and for mothers of control infants as the sleep during the night before the interview. The mean age of the infants at death was 98 days and the mean age at interview for the controls was 104 days. A total of 312 mothers of control infants (41% of those eligible) participated in the study.

Interviewers who were trained in grief counselling specifically related to SIDS obtained information on dummy use during the index sleep and on other environmental factors related to sleep, including sleep position (see bmj.com for details).

Analysis

We used odds ratios (95% confidence intervals) to estimate the relative risk of SIDS associated with use of a dummy and unconditional logistic regression analysis to examine the relation between use during sleep and the risk after adjustment for potential confounders.

Results

Table 1 shows the characteristics of cases and controls of our study population and the association between use of a dummy during the last sleep and the risk of

SIDS (table 1). After adjustment for known confounders, the odds ratio for SIDS in infants who used a dummy during the last sleep was 0.08 (0.03 to 0.21), which translates to a more than 90% reduction of risk in this study compared with infants who did not use a pacifier during the sleep.

Use of a dummy during sleep was associated with a reduced risk of SIDS in every category of those maternal and infant factors we examined. There was a consistent trend towards use of a dummy during sleep being associated with a greater reduction of risk when an infant was in an adverse sleep environment, such as sleeping prone or on the side, sleeping with a mother who was a current smoker, or sleeping on soft bedding (table 1).

Infant thumb sucking itself was associated with a reduced risk of SIDS (adjusted odds ratio 0.43, 0.25 to 0.77). However, use of a dummy was associated with a reduced risk of SIDS regardless of whether the baby sucked its thumb (0.07, 0.01 to 0.64, with thumb sucking and 0.08, 0.03 to 0.23, without (table 1). In addition, use was associated with a reduced risk in infants who ever breastfed and in infants currently breastfeeding (0.09, 0.03 to 0.25) (table 1).

In infants who did not use a dummy during the last sleep, a prone or side sleeping position, sleeping with a mother who smoked, and sleeping on soft bedding during the last sleep were associated with increased risk. In infants who did use a dummy during the last sleep, however, these known risk factors no longer had an effect (table 2). None the less, the difference in odds ratios between infants who did or did not use a dummy was not significant (table 2). (see bmj.com for details)

Discussion

In this population based case-control study, use of a dummy was associated with a substantial reduction in the risk of SIDS. This reduced risk was independent of other known risk factors. Our observation confirms several previous reports.³⁻⁷ Use of a dummy was also consistently associated with lower risk irrespective of the presence or absence of other risk factors, including those related to sleep environment. The association of use seemed stronger when an infant was already in an adverse sleep environment, such as in a prone or side sleep position, sleeping with a mother who smoked, or sleeping on soft bedding; and use of a pacifier seemed to reduce the impact of these risk factors, especially those related to adverse sleep conditions, although these differences did not reach significance. It is important that these findings be confirmed as they may provide new insight into the underlying mechanisms of the protective effect of dummies.

The use of dummies may be an effective strategy for public health intervention. While the rate of SIDS has declined in all populations in the United States, large racial disparities remain. In our study population, the incidence of use of dummies among populations at high risk of SIDS (African-Americans, young mothers, women with less than a college education, or low income) was comparable with that among those at lower risk of SIDS. The reduced risk did not seem to vary by social and demographic characteristics so advocating use of dummies for infants in high risk populations may have the potential to reduce the incidence of SIDS further.

Pregnancy and Perinatology Branch, Centre for Research on Mothers and Children, National Institute of Child Health and Human Development, NIH, Bethesda, MD 20892, USA
Marian Willinger
health scientist administrator

Epidemiology and Biostatistics Program, Division of Scientific Programs, National Institute on Deafness and Other Communication Disorders, NIH, Bethesda, MD 20892, USA
Howard J Hoffman
director

Correspondence to: Dr Li
dkl@dor.kaiser.org

Table 1 Relation of dummy use with risk of SIDS by maternal and infant characteristics and sleep environment

	SIDS cases		Controls		Crude odds ratio (95% CI)	Adjusted odds ratio* (95% CI)
	No of participants	Used a dummy (%)	No of participants	Used a dummy (%)		
All subjects	169	7 (4.1)	309	73 (23.6)	0.14 (0.06 to 0.31)	0.08 (0.03-0.21)
Maternal characteristics						
Race/ethnicity:						
White	51	2 (3.9)	118	38 (32.2)	0.09 (0.02 to 0.37)	0.04 (0.01-0.22)
African-American	32	1 (3.1)	59	16 (27.1)	0.09 (0.01 to 0.69)	0.06 (0.01-0.51)
Hispanic	59	2 (3.4)	97	14 (14.4)	0.21 (0.05 to 0.95)	0.10 (0.02-0.50)
Asian/other	27	2 (7.4)	35	5 (14.3)	0.48 (0.09 to 2.69)	0.81 (0.08-8.06)
Maternal age (years):						
<20	23	1 (4.3)	27	6 (22.2)	0.16 (0.02 to 1.44)	0.02 (0.002-0.35)
20-24	52	1 (1.9)	54	18 (33.3)	0.04 (0.005 to 0.31)	0.03 (0.004-0.26)
≥25	94	5 (5.3)	228	49 (21.5)	0.21 (0.08 to 0.53)	0.16 (0.05-0.47)
Maternal education:						
High school or less	112	6 (5.4)	155	37 (23.9)	0.18 (0.07 to 0.45)	0.08 (0.03-0.23)
At least some college	57	1 (1.8)	153	35 (22.9)	0.06 (0.01 to 0.45)	0.08 (0.01-0.62)
Marital status:						
Married	82	1 (1.2)	203	48 (23.6)	0.04 (0.005 to 0.29)	0.03 (0.004-0.24)
Cohabiting	58	4 (6.9)	72	18 (25.0)	0.22 (0.07 to 0.70)	0.08 (0.02-0.33)
Unmarried†	29	2 (6.9)	34	7 (20.6)	0.29 (0.05 to 1.50)	0.33 (0.05-2.04)
Annual income:						
<\$20 000	69	3 (4.3)	85	18 (21.2)	0.17 (0.05 to 0.60)	0.08 (0.01-0.66)
\$20 000-49 999	54	2 (3.7)	83	22 (26.5)	0.11 (0.02 to 0.48)	0.07 (0.01-0.34)
≥\$50 000	39	1 (2.6)	123	30 (24.4)	0.08 (0.01 to 0.62)	0.05 (0.01-0.24)
Parity:						
1	52	1 (1.9)	132	28 (21.2)	0.07 (0.01 to 0.55)	0.04 (0.004-0.30)
2	62	1 (1.6)	95	28 (29.5)	0.04 (0.005 to 0.30)	0.03 (0.004-0.26)
3	29	3 (10.3)	50	10 (20.0)	0.46 (0.12 to 1.84)	0.37 (0.06-2.35)
≥4	26	2 (7.7)	32	7 (21.9)	0.30 (0.06 to 1.58)	0.16 (0.02-1.13)
Gestational age (weeks) at initial prenatal care visit:						
<12	109	5 (4.6)	238	54 (22.7)	0.16 (0.06 to 0.42)	0.09 (0.03-0.27)
≥12	58	2 (3.4)	67	19 (28.4)	0.09 (0.02 to 0.41)	0.07 (0.01-0.38)
Smoking status during pregnancy:						
No (non-smoker)	128	4 (3.1)	262	54 (20.6)	0.12 (0.04 to 0.35)	0.10 (0.03 to 0.30)
Yes (smoker)	41	3 (7.3)	46	19 (41.3)	0.11 (0.03 to 0.42)	0.06 (0.01 to 0.28)
Infant characteristics						
Infant age (months):						
≤1	44	2 (4.6)	60	11 (18.3)	0.21 (0.04 to 1.01)	0.08 (0.01-0.60)
2-3	82	4 (4.9)	154	34 (22.1)	0.18 (0.06 to 0.53)	0.11 (0.03-0.38)
≥4	43	1 (2.3)	95	28 (29.5)	0.06 (0.01 to 0.43)	0.04 (0.005-0.33)
Birth weight (g):						
≥2500	135	6 (4.4)	294	71 (24.2)	0.15 (0.06 to 0.35)	0.08 (0.03-0.21)
<2500	31	1 (3.2)	14	2 (14.3)	0.20 (0.02 to 2.42)	0.39 (0.03-5.97)
Infant sickness during previous 48 hours:						
None	95	4 (4.2)	194	45 (23.2)	0.15 (0.05 to 0.42)	0.07 (0.02-0.25)
Yes	73	3 (4.1)	115	28 (24.4)	0.13 (0.04 to 0.46)	0.09 (0.02-0.38)
Sucked thumb during last week:						
No	139	6 (4.3)	210	54 (25.7)	0.13 (0.05 to 0.31)	0.08 (0.03-0.23)
Yes	30	1 (3.3)	99	19 (19.2)	0.15 (0.02 to 1.13)	0.07 (0.01-0.64)
Breastfeeding:						
No	36	1 (2.8)	43	10 (23.3)	0.09 (0.01 to 0.78)	0.05 (0.01-0.51)
Yes	133	6 (4.5)	266	63 (23.7)	0.15 (0.06 to 0.36)	0.09 (0.03-0.25)
Sleep environment						
Last sleep position:						
On back	51	3 (5.9)	174	39 (22.4)	0.22 (0.06 to 0.73)	0.20 (0.05-0.72)
Prone or on side	111	4 (3.6)	135	34 (25.2)	0.11 (0.04 to 0.32)	0.05 (0.01-0.17)
Cosleeping:						
None	98	3 (3.1)	193	46 (23.8)	0.10 (0.03 to 0.33)	0.07 (0.01-0.28)
With non-smoker	41	3 (7.3)	102	21 (20.6)	0.30 (0.09 to 1.08)	0.17 (0.04-0.69)
With mother who smoked	27	1 (3.7)	12	6 (50.0)	0.04 (0.004 to 0.38)	0.02 (0.002-0.26)
Soft bedding during last sleep:						
No	121	6 (5.0)	243	57 (23.5)	0.17 (0.07 to 0.41)	0.12 (0.04-0.31)
Yes	40	1 (2.5)	66	16 (24.2)	0.08 (0.01 to 0.63)	0.02 (0.001-0.25)

*Adjusted for maternal age, race, education, smoking during pregnancy, infant age, last sleep position, birth date, and region (study centre) unless variables themselves were examined as modifier.

†Includes unmarried/separated/divorced/widowed.

Table 2 Dummy use in relation to selected risk factors for SIDS in sleep environment. Figures are numbers* (percentages) of infants with odds ratios

	Did not use dummy			Used a dummy			P value for difference in OR
	Case (n=169)	Control (n=309)	OR† (95% CI)	Case (n=169)	Control (n=309)	OR† (95% CI)	
Last sleep position:							
On back‡	48 (31.0)	135 (57.2)	2.61	3 (42.9)	39 (53.4)	0.66	0.38
Prone/on side	107 (69.0)	101 (42.8)	(1.56 to 4.38)	4 (57.1)	34 (46.6)	(0.12 to 3.59)	
Cosleeping:							
None/with non-smoker‡	133 (83.6)	228 (97.4)	4.50	6 (85.7)	67 (91.8)	1.10	0.26
With mother who smoked	26 (16.4)	6 (2.6)	(1.32 to 15.10)	1 (14.3)	6 (8.2)	(0.10 to 13.40)	
Soft bedding:							
No‡	115 (74.7)	186 (78.8)	1.42	6 (85.7)	57 (78.1)	0.26	.15
Yes	39 (25.3)	50 (21.2)	(0.79 to 2.57)	1 (14.3)	16 (21.9)	(0.02 to 3.27)	

*Numbers in the table may not match the total numbers because of missing data.

†Adjusted for maternal age, race, education, and smoking during pregnancy, infant age and birth date, and region (study centre).

‡Reference group.

Limitations

Case-control studies are generally more susceptible to potential biases including selection and reporting biases, and results therefore require greater scrutiny and caution is needed in establishing a causal relation. Our sample sizes in some of the stratified analyses were small, the precision of the estimates being reflected in the width of corresponding 95% confidence intervals. Our discussion of potential underlying mechanisms of our observed association should not be confused with the “certainty” of a causal effect. The findings from our study and others will collectively provide support for the protective effect of dummies, but they are not “proof” of a causal effect by themselves.

In addition, the participation rate was low in our study. We examined the potential selection bias due to the low participation rate in a previous report from this study.¹ We obtained birth certificates for all eligible infants, regardless of whether their mothers participated in the study, and calculated odds ratios of known risk factors for SIDS using information from the birth certificates. The odds ratios for infants under study were comparable with those obtained from all eligible infants, had they participated.¹ Furthermore, the prevalence of use of dummies in our control group was comparable with that reported in other US populations.⁶ Finally, the magnitude of the odds ratio of SIDS

associated with use of dummies was similar to that observed in other studies.^{6,7} While these observations reduce concerns of selection bias due to low participation, it remains a possibility.

Another possibility is that our results were biased by deliberate under-reporting by mothers of infants who died. However, we did not detect any deliberate under-reporting of several more well known risk factors, such as prone sleep position, maternal smoking, and cosleeping, which were more commonly reported among infants who died. It is unlikely that mothers would have chosen to deliberately under-report only use of dummies, which, at the time of the interview was not generally known to the public as being related to SIDS in any way, while not under-reporting other well known risk factors.

Contributors: See bmj.com.

Funding: National Institute of Child Health and Human Development (NICHD) and National Institute on Deafness and Other Communication Disorders (NIDCD), National Institutes of Health, contract N01-HD-5-3227.

Competing interest: None declared.

Ethical approval: Institutional Review Boards of Kaiser Permanente in Northern and Southern California and California State Committee for the Protection of Human Subjects.

What is already known on this topic

Sleep environment has consistently been reported as influencing the risk of sudden infant death syndrome (SIDS)

Use of a dummy during sleep has been reported as reducing the risk of SIDS

What this study adds

Use of a dummy during sleep was associated with a reduced risk of SIDS consistently across a wide range of socioeconomic characteristics and risk factor profiles

Use also reduced the adverse effects of a prone sleep position, sleeping with a mother who smoked, and soft bedding

- Li DK, Petitti DB, Willinger M, McMahon R, Odouli R, Vu H, et al. Infant sleeping position and the risk of sudden infant death syndrome in California, 1997-2000. *Am J Epidemiol* 2003;157:446-55.
- Willinger M, Ko CW, Hoffman HJ, Kessler RC, Corwin MJ. Factors associated with caregivers' choice of infant sleep position, 1994-1998: the national infant sleep position study. *JAMA* 2000;283:2135-42.
- Arnestad M, Andersen M, Rognum TO. Is the use of dummy or carry-cot of importance for sudden infant death? *Eur J Pediatr* 1997;156:968-70.
- Fleming PJ, Blair PS, Pollard K, Platt MW, Leach C, Smith I, et al. Pacifier use and sudden infant death syndrome: results from the CESDI/SUDI case control study. CESDI SUDI research team. *Arch Dis Child* 1999;81:112-6.
- Mitchell EA, Taylor BJ, Ford RP, Stewart AW, Becroft DM, Thompson JM, et al. Dummies and the sudden infant death syndrome. *Arch Dis Child* 1993;68:501-4.
- Hauck FR, Herman SM, Donovan M, Iyasu S, Merrick MC, Donoghue E, et al. Sleep environment and the risk of sudden infant death syndrome in an urban population: the Chicago infant mortality study. *Pediatrics* 2003;111:1207-14.
- L'Hoir MP, Engelberts AC, van Well GT, Damste PH, Idema NK, Westers P, et al. Dummy use, thumb sucking, mouth breathing and cot death. *Eur J Pediatr* 1999;158:896-901.
- Franco P, Chabanski S, Scaillet S, Groswasser J, Kahn A. Pacifier use modifies infant's cardiac autonomic controls during sleep. *Early Hum Dev* 2004;77:99-108.
- Franco P, Scaillet S, Wermenbol V, Valente F, Groswasser J, Kahn A. The influence of a pacifier on infants' arousals from sleep. *J Pediatr* 2000;136:775-9.

(Accepted 8 November 2005)

doi 10.1136/bmj.38671.640475.55