

Effect of recolonisation with “interfering” α streptococci on recurrences of acute and secretory otitis media in children: randomised placebo controlled trial

Kristian Roos, Eva Grahn Håkansson, Stig Holm

Ear, Nose, and Throat Department, Lundby Hospital, 41717 Gothenburg, Sweden

Kristian Roos
associate professor

Department of Clinical Bacteriology, University of Umeå, 90187 Umeå, Sweden

Eva Grahn Håkansson
research associate

Department of Clinical Microbiology, University of Umeå, Stig Holm professor

Correspondence to: K Roos
kristian.roos@lundbysjukhus.se

BMJ 2001;322:210–2

Abstract

Objective To study the effect of recolonisation with α streptococci with the ability to inhibit the growth of otopathogens (“interfering” activity) on the recurrence of acute otitis media in susceptible children and the effect on the frequency of secretory otitis media.

Design Double blind, randomised, placebo controlled study.

Setting Ear, nose, and throat clinic with three doctors. **Participants** 130 children prone to otitis media aged between 6 months and 6 years, 108 of whom were eligible and followed for 3 months.

Main outcome measures Recurrence of otitis media during follow up and a normal tympanic membrane at the last valid visit.

Interventions Children with no recurrences during the last month received phenoxymethylpenicillin (n = 22), and those with a recurrence within 1 month received amoxicillin clavulanic acid (n = 86), both twice daily for 10 days. These were followed by a streptococcal or placebo solution sprayed into the nose for a further 10 days. At day 60 the same spray was started for another 10 days.

Results At 3 months 22 children (42%) given the streptococcal spray were healthy and had a normal tympanic membrane compared with 12 (22%) of those given placebo. This difference was shown separately for recurrences of both acute otitis media and secretory otitis media.

Conclusions Selected bacteria with the ability to inhibit the growth of common otopathogens can be used to protect against recurrent acute otitis media and secretory otitis media in children.

Introduction

Acute otitis media is the most common bacterial infection in young children, and large amounts of antibiotics are prescribed, especially for those with recurrent episodes. The peak incidence of acute otitis media is at 1–2 years of age. The risk of developing another episode within one month after the onset of the primary infection is estimated at 35%.¹ About 5% of children are prone to otitis media, defined as six or more episodes, or recently as three or more episodes,

during one year.^{2,3} The recent definition results in a greater number of children who are considered prone to otitis media.

The most common bacteria associated with acute otitis media are *Streptococcus pneumoniae*, *Haemophilus influenzae* and, less often, *Moraxella catarrhalis* and group A β streptococci. These bacteria originate and spread from the nasopharynx to the middle ear cavity by way of the eustachian tube. Carlin et al showed that 75% of the bacteria associated with recurrent otitis media represented new bacterial strains.¹ The remaining 25% were from either reinfection with the same bacterial strain or treatment failures.

Secretory otitis media is the most common sequela of acute otitis media. One or more of *S pneumoniae*, *H influenzae*, or *M catarrhalis* are found in about 30% of patients with secretory otitis media.³

The importance of normal flora for protecting against infection in an anatomical site has recently been shown in the upper respiratory tract, and lack of bacteria with interfering activity (the ability to inhibit the growth of the common otopathogens), especially the α streptococci, has been associated with a higher incidence of reinfections in patients with streptococcal pharyngotonsillitis.^{4,5} Lower numbers of α streptococci have been found in the nasopharynx of children who are prone to otitis media compared with those who are not prone and in those with secretory otitis media compared with healthy children.^{6–8}

We aimed to study the effect of recolonisation with α streptococci with interfering activity against the common otopathogens on the recurrence rate of acute otitis media. We also aimed to determine whether the frequency of secretory otitis media was affected by this treatment.

Participants and methods

Study design—From 1996 to 1999 we performed a double blind, randomised study with two arms on 130 children aged between 6 months and 6 years.

Patients—Children eligible for inclusion in our study were those who had had recurrent otitis media and who had been either referred by their general practitioner or a paediatrician to the open care unit of the ear, nose, and throat department at Lundby Hospital or were directly seeking medical advice for ear pain.



The full version of this paper appears on the BMJ's website

The children had had at least two episodes of acute otitis media during the past six months or five episodes during the past year. At the next occurrence of ear pain the children were examined, and those with a red or pale, bulging, thickened tympanic membrane were included in the study. We excluded those with penicillin allergy, serious underlying disease, immunological deficiency, a valvular heart defect, major lesions in the mouth or nose, a grommet in the ear, or chronic otitis media.

Informed written consent was obtained from one of the parents of each eligible child. The study protocol was approved by the Medical Products Agency and the local ethics committee in Gothenburg.

Antibiotic treatment—Those children with no recurrences during the past month but who had acute otitis media were given phenoxymethylpenicillin (Kåvepenin, AstraZeneca, Sweden) 25 mg/kg bodyweight. Those children with a recurrence within the past month were given amoxicillin clavulanic acid (Spektramox, AstraZeneca) 20 mg/kg bodyweight. Both antibiotics were given twice daily for 10 days.

Spray treatment—The streptococcal spray was made up by isolating a haemolytic streptococci from the opening of the eustachian tube of the healthy children and selecting five strains (of about 800 tested) for their superior ability to inhibit the growth of *S pneumoniae*, *H influenzae*, *M catarrhalis*, and *S pyogenes* (group A streptococci).⁹ Placebo spray comprised skimmed milk powder, with the same texture and colour as the streptococcal spray.

Participant flow and follow up analysis

At the first visit (day 1) the child's medical history and background data were recorded and a clinical examination was performed, including otomicroscopy (this is superior to otoscopy).

At the second visit (days 8-10) the children were examined by otomicroscopy and excluded from the study if signs or symptoms of infection were still present. The parent was given a fresh bottle of spray with instructions to give three puffs into each nostril twice a day for 10 days. At the end of spray treatment the children returned for a third visit (days 25-30), and the children or parents, or both, were interviewed about compliance. At the fourth visit (days 55-60) a further 10 day course of spray treatment was started. The last valid visit was between days 88 and 92.

A clinical examination was performed at all visits, the status of the tympanic membranes checked by otomicroscopy, and the clinical response classified as cured, improved, secretory otitis media, or recurrence. At the third and fifth visits the amount of unused spray in the bottle was recorded. Adverse events were recorded at follow up visits. We used Fisher's exact test and logistic regression.

Results

Of 132 children included, 108 (82%) were eligible for analysis of efficacy (53 in the α spray group and 55 in the placebo group) and 126 (95%) for analysis of adverse events.

The mean age of the children was 23 months. We found no differences between the two treatment

Cure rate in children prone to otitis either recolonised with α streptococci or given placebo. Values are numbers (percentages) unless stated otherwise

	Streptococci (n=53)*	Placebo (n=55)†	95% CI‡	P value
Cured	22 (42)	12 (22)	0.028 to 0.372	0.02
Recurrence of otitis media during follow up (92 days)	21 (40)	28 (51)	0.065 to 0.285	0.04
Secretory otitis media at last valid visit	10	15	0.074 to 0.286	0.05

*12 treated with penicillin and 41 with amoxicillin clavulanic acid.

†10 treated with penicillin and 45 with amoxicillin clavulanic acid.

‡Difference between proportion cured with streptococcal spray and proportion cured with placebo.

groups for age, number of siblings, parental proneness to otitis media, allergy, duration of breast feeding, day care, or parental smoking. Children who had had six or more episodes of acute otitis media during the past year, or two or more during the past six months, were equally distributed between the two treatment groups, and there were no differences in efficacy outcome between these two groups.

Of the children receiving the spray, 27 (22 of whom could be evaluated) were given penicillin and 103 (86 of whom could be evaluated) amoxicillin clavulanic acid. These children were equally distributed between the spray and placebo groups.

In children given the spray the rate of recurrence during the three months of follow up was significantly reduced compared with those given placebo. Overall, 22 (42%) of the children given spray experienced no acute otitis media during the study and had a normal tympanic membrane at the last valid visit compared with 12 (22%) of the children given placebo (table). Furthermore, 10 (31%) of the 32 children without recurrence who were given the spray had secretory otitis media at the last valid visit compared with 15 (56%) of the 27 children in the placebo group.

Of the 130 children included, 22 in the spray group had adverse events compared with 25 in the placebo group. One child in the placebo group got pneumonia and spray treatment was stopped, and another child in the same group stopped treatment owing to an adverse event.

Discussion

We found a 50% rate of recurrence of acute otitis media within three months of an episode in children who are prone to otitis despite adequate antibiotic treatment. We also found a high frequency of secretory otitis media. Up to 78% of the children treated with antibiotics and placebo either had a recurrence or still had secretory otitis media after three months.

Antibiotic prophylaxis has been questioned because of the increasing antibiotic resistance of respiratory tract pathogens.^{10 11} The placement of a tympanostomy tube seems to effectively prevent recurrent otitis media in these children.¹¹ The procedure, however, carries a risk, is costly, needs to be performed under general anaesthesia, and has complications and sequelae related to the tympanic membrane.¹²

People who lack interfering α streptococci seem to have more streptococcal throat infections than those with interfering α streptococci on their tonsils.^{4 5} Furthermore, patients with recurrent streptococcal pharyngotonsillitis have fewer recurrences after recolonisation with a mixture of four α streptococcal

What is already known on this topic

Lack of interfering bacteria, especially the α streptococci, has been associated with a high incidence of reinfection in streptococcal pharyngotonsillitis

Lower numbers of α streptococci have been found in the nasopharynx of children prone to otitis compared with those who are not and of children with secretory otitis media compared with healthy children

What this study adds

Recolonisation with α streptococci with inhibitory activity against common pathogens in the upper respiratory tract diminishes the recurrence rate of acute otitis media in susceptible children

This is also true for secretory otitis media, often seen as a complication of acute otitis media

Treatment with α streptococci could help reduce antibiotic consumption

strains with good growth inhibiting activity of group A streptococci.^{13 14}

Nasopharyngeal cultures from children who are prone to otitis or secretory otitis media show low numbers of α streptococci with interfering activity against common pathogens of acute otitis media. We therefore tried to recolonise children prone to recurrent acute otitis media with a mixture of five strains of interfering α streptococci. This resulted in a significantly decreased number of recurrences of acute otitis media in the treated children compared with those given placebo. This difference was also seen in the children with secretory otitis media.

Most antibiotics used to treat infections in the upper respiratory tract have an impact on the normal bacterial flora, including the dominating α streptococci. As these bacteria are part of the body's natural defence, treatment with antibiotics abates this part of the defence system and thus facilitates colonisation with pathogenic bacteria. Paradoxically, repeated courses of antibiotics might contribute to recurrent infections in children who are prone to otitis. Restoration of the normal flora would therefore be the logical way to inhibit further recurrences. It has recently been shown that other bacteria such as *Prevotella* and *Peptostreptococcus* species may also be candidates for ecological recolonisation.⁷

In conclusion, the reduction in number of recurrences after treatment with α streptococci may seem low but as otitis prone children are heavy consumers of antibiotics the impact on the development of antibiotic resistance among pathogens causing otitis media in this group of children may be considerable.

Lundby Hospital supported the clinical part of the study by giving access to necessary staff.

Contributors: KR, EGH, and SH formulated the study hypothesis, discussed core ideas, designed the protocol, and evaluated the data. KR coordinated the study and included most

of his patients in the study. SH supervised the bacteriological part of the study. EGH was responsible for the selection of interfering α streptococci and prepared the streptococcal suspension together with Marie Eklund. Hanna Eklöf monitored the study. Eva Lydén and Carl von Sydow included patients from Lundby Hospital. Hans Stenlund, University of Umeå, was responsible for the statistical analysis. The Medical Products Agency in Uppsala approved the design and suggested minor changes.

Funding: The study was supported by the Swedish National Board for Industrial and Technical Development and the Teknikbro Foundation and grants from Samariten Foundation, Stockholm.

Competing interests: We have been involved in the interference between potentially pathogenic bacteria and apathogens in the upper respiratory tract for many years. This has resulted in several theses at the University of Umeå and Gothenburg. This study is a continuation of ongoing scientific studies covering the upper respiratory tract. We believe that bacterial interference is of importance for the normal defence system and has a clinical impact. We hope that it might be routinely applied as an alternative, or supplement, to antibiotic treatment in the future. We have therefore applied for a patent in some countries for the bacterial strains used in the study.

- 1 Carlin S, Marchant C, Shurin P, Johnson C, Muddell-Panek D, Barenkamp S. Early recurrence of otitis media: reinfection or relapse. *J Pediatr* 1987;110:20-5.
- 2 Ingvarsson L, Lundgren K, Stenström C. Occurrence of acute otitis media in children: cohort studies in an urban population. *Ann Otol Rhinol Laryngol* 1990;suppl 149:17-8.
- 3 Stenfors L-E, Räisänen S. Occurrence of *Streptococcus pneumoniae* and *Haemophilus influenzae* in otitis media with effusion. *Clin Otolaryngol* 1992;17:195-9.
- 4 Roos K, Grahn E, Holm SE. Evaluation of beta-lactamase activity and microbial interference in treatment failures of acute streptococcal tonsillitis. *Scand J Infect Dis* 1986;18:313-9.
- 5 Grahn E, Holm SE. Bacterial interference in the throat during a streptococcal tonsillitis outbreak in an apartment house area. *Zbl Bact Hyg A* 1983;256:72-9.
- 6 Bernstein J, Faden H, Dryja D, Wactawski-Wende J. Micro-ecology of the nasopharyngeal bacterial flora in otitis-prone and non-otitis-prone children. *Acta Otolaryngol (Stockh)* 1993;113:88-92.
- 7 Brook I, Yocum P. Bacterial interference in the adenoids of otitis media-prone children. *Pediatr Infect Dis J* 1999;18:835-7.
- 8 Fujimori I, Hisamatsu K, Kikushima K, Goto R, Murakami Y, Yamada T. The nasopharyngeal flora in children with otitis media with effusion. *Eur Arch Otorhinolaryngol* 1996;253:260-3.
- 9 Grahn E, Holm SE, Roos K, Ekedahl C. Interference of alpha-hemolytic streptococci isolated from tonsillar surface, on hemolytic streptococci, *Streptococcus pyogenes*, a methodological study. *Zbl Bact Hyg A* 1983;254:459-68.
- 10 Casselbrant ML, Kaleida P, Rockette H, Paradise J, Bluestone C, Kurs-Lasky M, et al. Efficacy of antimicrobial prophylaxis and tympanostomy tube insertion for prevention of recurrent acute otitis media: results of a randomized clinical trial. *Pediatr Infect Dis J* 1992;11:278-86.
- 11 Prellner K, Foglé-Hansson M, Jørgensen F, Kalm O, Kamme C. Prevention of recurrent acute otitis media in otitis-prone children by intermittent prophylaxis with penicillin. *Acta Otolaryngol (Stockh)* 1994;114:182-7.
- 12 Tos M, Stangerup SE. Hearing loss in tympanosclerosis caused by grommets. *Arch Otolaryngol Head Neck Surg* 1989;115:931-5.
- 13 Roos K, Holm SE, Grahn E, Lind L. Alpha-streptococci as supplementary treatment of recurrent streptococcal tonsillitis: a randomized placebo-controlled study. *Scand J Infect Dis* 1993;25:31-5.
- 14 Roos K, Holm SE, Grahn E, Lagergren L. Recolonization with four selected alpha-streptococcal strains in the treatment of recurrent streptococcal tonsillitis—a placebo-controlled randomized multicentre study. *Scand J Infect Dis* 1996;28:459-62.

(Accepted 29 September 2000)

Endpiece

The Mystical Marriage

He was groping for the Mystical Marriage, the unity of the masculine and the feminine in himself, without which he would have been useless in his future life as an artist and as a man who understood art. Useless as any sort of man—rich man, poor man, beggar man, thief; not to speak of tinker, tailor, soldier, sailor—who is destined to see more than a few inches beyond the end of his nose.

Robertson Davies (1913-95), *What's bred in the bone*, London: Penguin, 1986