

## Comparison of cardiovascular risk between patients with type 2 diabetes and those who had had a myocardial infarction: cross sectional and cohort studies

Josie M M Evans, Jixian Wang, Andrew D Morris



The full version of this article appears on [bmj.com](http://bmj.com)

### Abstract

**Objective** To compare risks of cardiovascular outcomes between patients with type 2 diabetes and patients with established coronary heart disease.

**Design** Cross sectional study and cohort study using routinely collected datasets.

**Setting** Tayside, Scotland (population 400 000) during 1988-95.

**Subjects** In the cross sectional study, among patients aged 45-64, 1155 with type 2 diabetes were compared with 1347 who had had a myocardial infarction in the preceding 8 years. In the cohort study 3477 patients of all ages with newly diagnosed type 2 diabetes were compared with 7414 patients who had just had a myocardial infarction.

**Main outcome measures** Risk ratios for death from all causes, cardiovascular death, and hospital admission for myocardial infarction were calculated by Cox proportional hazards analysis and adjusted for age and sex.

**Results** In the cross sectional study the adjusted risk ratio for death from all causes was 2.27 (95% confidence interval 1.82 to 2.83) for patients who had had myocardial infarction compared with those with diabetes, and the risk ratio for hospital admission for myocardial infarction was 1.33 (1.14 to 1.55). In the cohort study, patients who had just had a myocardial infarction had a higher risk of death from all causes (adjusted risk ratio 1.35 (1.25 to 1.44)), cardiovascular death (2.93 (2.54 to 3.41)), and hospital admission for myocardial infarction (3.10 (2.57 to 3.73)).

**Conclusions** Patients with type 2 diabetes were at lower risk of cardiovascular outcomes than patients with established coronary heart disease.

### Introduction

The increased risk of cardiovascular disease in type 2 diabetes is well recognised<sup>1</sup> and is associated with both diabetes specific risk factors<sup>2</sup> and increased frequency of conventional risk factors for cardiovascular disease.<sup>3</sup> A recent study by Haffner et al suggested that the risk of death from coronary heart disease in patients with type 2 diabetes may even be as high as in patients who have had a myocardial infarction.<sup>4</sup> The authors of this study proposed that patients with diabetes should be

treated as if they had existing coronary heart disease. Important weaknesses of this study, however, were the lack of power to detect differences between the two groups and that patients were self selected.

The aim of the present investigation was to compare risks of cardiovascular outcomes between patients with diabetes and patients with established coronary heart disease. We carried out two studies: in the first we used a design similar to that of Haffner et al,<sup>4</sup> and in the second we compared patients with newly diagnosed diabetes for cardiovascular risk with those who had just had a myocardial infarction.

### Subjects and methods

#### Data sources

We used routine data for residents of Tayside, Scotland, including data of all patients with type 1 and type 2 diabetes from 1988, a record of all inpatient hospital admissions in Tayside from 1980 with diagnostic codes from ICD-9 (international classification of diseases, ninth revision), and records of death certificates from the registrar general with ICD-9 coded causes of death from 1988.<sup>5,6</sup>

#### Cross sectional study

We identified a cross section of patients aged 45-64 years at the study index date (1 January 1988). We defined a diabetic group, which included all patients with type 2 diabetes diagnosed before the index date. We also defined a group of patients, aged 45-64, who had an inpatient hospital admission with myocardial infarction between January 1980 and December 1987. We excluded any patient who was in both groups. Follow up was from January 1988 to December 1995.

The outcomes were mortality (any cause), ascertained from death certification records, and hospital admission with a primary diagnosis of myocardial infarction. We used a Cox proportional hazards model to calculate risk ratios (with 95% confidence intervals), adjusted for age and sex, for the risks of the outcomes in the myocardial infarction group compared with those in the diabetic group.

#### Cohort study

We identified two cohorts of patients. The diabetic cohort included any patient in the study population

Department of Epidemiology and Public Health, Ninewells Hospital, Dundee DD1 9SY

Josie M M Evans  
*lecturer*

Department of Clinical Pharmacology, Ninewells Hospital, Dundee

Jixian Wang  
*statistician*

Department of Medicine, Ninewells Hospital, Dundee  
Andrew D Morris  
*reader in medicine*

Correspondence to:  
J M M Evans  
[j.m.m.stansfield@dundee.ac.uk](mailto:j.m.m.stansfield@dundee.ac.uk)

BMJ 2002;324:939-42

with type 2 diabetes diagnosed between January 1988 and December 1995. Their study index date was their date of diagnosis of diabetes. The myocardial infarction cohort included any patient who had a first hospital admission with a primary or secondary diagnosis of myocardial infarction between January 1988 and December 1995. Their study index date was their date of hospital admission.

The outcomes in this study during follow up to December 1995 were mortality (any cause), cardiovascular mortality, and hospital admission for myocardial infarction (primary or secondary diagnosis). Any patient who died within 28 days after their index date was excluded from further analyses. We constructed Kaplan-Meier survival curves to compare probability of death between the study cohorts. We used a Cox proportional hazards model to compare the risks, with adjustment for age and sex.


## Results

Our study population consisted of 263 175 patients resident in Tayside.

### Cross sectional study

We identified 1155 patients (42% women) with diabetes, with a mean age of 57 years and a mean duration of diabetes of 6.04 years. We identified 1347 patients (28% women) with myocardial infarction, with a mean age of 57 and a mean time since infarction of 3.5 years.

In the eight year follow up 438 (33%) of the patients in the myocardial infarction group died, and 274 (20%) were hospitalised for a further infarction. In the diabetic group 284 (25%) died, and 113 (10%) were



Kaplan-Meier survival curve showing time to death from any cause in patients with newly diagnosed type 2 diabetes and patients who had just had a myocardial infarction (MI)

hospitalised for an infarction. The risk of death was higher in the myocardial infarction group compared with the diabetic group. Hospital admission for myocardial infarction also increased more than twofold in this group (table 1). The risks of these outcomes seemed to be greater for men than women, and the risks increased with age (although age was a weaker predictor of hospital admission for myocardial infarction than of death).

### Cohort study

For the diabetic cohort, we identified 3477 patients (49% women), with a mean age of 66, of whom 3403 (98%) were included in the statistical analyses (patients who did not die within 28 days after their index date). For the myocardial infarction cohort, we identified 7414 patients (46% women), with a mean age of 71, of whom 5350 (72%) were included in the statistical analyses.

During the subsequent follow up, there were 1196 deaths in the diabetic cohort (35% of the patients in the analyses), of which 222 were cardiovascular deaths, and 142 patients were hospitalised for myocardial infarction. In the myocardial infarction cohort there were 2596 deaths (49% of patients in the analyses), of which 1077 were cardiovascular deaths, and 656 patients were hospitalised for myocardial infarction. The Kaplan-Meier survival curve shows that the diabetic cohort had lower risks for death from all causes (figure). The reduced risk seemed to be mainly in the early period of follow up, with the gradients of the curves converging after the first 1000 days or so.

Patients in the myocardial infarction cohort had an increased risk of death from any cause compared with patients in the diabetic cohort (table 2). The risk increased with age, and men were at greater risk than women. Patients in the myocardial infarction cohort had a threefold higher risk of death from a cardiovascular cause than patients in the diabetic cohort, and the risk of cardiovascular death increased with age, but there was no difference between men and women. The myocardial infarction cohort also had an increased risk of hospital admission for myocardial infarction. The risks of hospital admission were higher in the age groups 60-69 and 70-79 than in the group aged  $\geq 80$ , but there was no difference between men and women.

**Table 1** Risk of death from all causes and hospital admission for myocardial infarction in cross sectional study of patients with type 2 diabetes and those who had had a myocardial infarction

	Diabetic group		MI group		Risk ratio (95% CI)*
	No of patients	No of events	No of patients	No of events	
<b>Death from all causes</b>					
Age (years):					
45-49	102	8	106	23	0.34 (0.24 to 0.49)
50-54	260	39	237	60	0.48 (0.38 to 0.59)
55-59	299	68	366	100	0.61 (0.51 to 0.73)
60-64	494	169	638	255	1.00
Sex:					
Men	668	182	974	334	1.35 (1.15 to 1.61)
Women	487	102	373	104	1.00
Diagnosis:					
MI	—	—	1347	438	1.33 (1.14 to 1.55)
Diabetes	1155	284	—	—	1.00
<b>Hospital admission for myocardial infarction</b>					
Age (years):					
45-49	102	5	106	22	0.71 (0.47 to 1.06)
50-54	260	19	237	50	0.81 (0.61 to 1.06)
55-59	299	27	366	75	0.85 (0.67 to 1.08)
60-64	497	62	638	127	1.00
Sex:					
Men	668	71	974	208	1.28 (1.03 to 1.61)
Women	487	42	373	66	1.00
Diagnosis:					
MI	—	—	1347	274	2.27 (1.82 to 2.83)
Diabetes	1155	113	—	—	1.00

MI=myocardial infarction.

\*Adjusted for all covariates, derived from Cox regression analysis.

## Discussion

We found that patients with type 2 diabetes were at lower risk of death from all causes, death from cardiovascular causes, and hospital admission for myocardial infarction than patients with established coronary heart disease. Thus, our results do not support the hypothesis that patients with diabetes have as high a cardiovascular risk as patients with established coronary heart disease.

### Comparison with other studies

For our cross sectional study, we used a design similar to that used by Haffner et al in their Finnish study.<sup>4</sup> The patient groups were slightly different in that we included patients whose diabetes was treated by diet only, and patients in the myocardial infarction group were restricted to those whose myocardial infarction occurred in the preceding eight years. Patients in the myocardial infarction group were clearly at an increased risk of death and hospital admission for myocardial infarction, after adjustment for the differences in age and sex between the two groups. Although we were unable to adjust for differences in cardiovascular risk factors at baseline (smoking status, hypertension, low density and high density lipoprotein cholesterol concentrations, and total cholesterol and triglyceride concentrations), such adjustments in the Finnish study reduced the hazard ratio for death from coronary heart disease only slightly, from 1.4 to 1.2. In our study a substantial decrease would be required for the difference in risk of hospital admission for myocardial infarction to be no longer significant. An important point is the narrow width of the confidence intervals in our study.

Although we could not make a direct comparison between the results from the two studies, as we were unable to determine mortality from coronary heart disease in our patient groups, it is interesting that the result for overall mortality in our study falls within the confidence interval for mortality from coronary heart disease in the Finnish study. We believe that our study highlights the lack of power in the earlier study. It also shows that the cardiovascular risk profile is clearly different in patients with diabetes compared with those who have had a myocardial infarction.

### What is already known on this topic

A recent influential study suggested that patients with type 2 diabetes without established cardiovascular disease have as high a risk of cardiovascular events and death as non-diabetic patients who have had a myocardial infarction

Some clinicians therefore advocate aggressive treatment of cardiovascular risk factors in the presence of diabetes

### What this study adds

Patients with type 2 diabetes are at lower risk of death from all causes or cardiovascular causes and of hospital admission for myocardial infarction than patients with established coronary heart disease

**Table 2** Risk of death from all causes, death from cardiovascular causes, and hospital admission for myocardial infarction in cohort study of patients with newly diagnosed type 2 diabetes and those who had just had a myocardial infarction

	Diabetic group		MI group		Risk ratio (95% CI)*
	No of patients	No of events	No of patients	No of events	
<b>Death from all causes</b>					
Age (years):					
<50	441	30	397	41	0.05 (0.04 to 0.07)
50-59	677	92	870	191	0.12 (0.10 to 0.13)
60-69	1005	295	1457	548	0.25 (0.23 to 0.27)
70-79	840	434	1565	920	0.51 (0.47 to 0.55)
>79	440	345	1061	896	1.00
Sex:					
Men	1734	604	3062	1352	1.19 (1.12 to 1.26)
Women	1669	592	2288	1244	1.00
Diagnosis:					
MI	—	—	5350	2596	1.35 (1.25 to 1.44)
Diabetes	3403	1196	—	—	1.00
<b>Death from cardiovascular causes</b>					
Age (years):					
<50	441	6	397	17	0.10 (0.07 to 0.16)
50-59	677	12	870	86	0.22 (0.18 to 0.27)
60-69	1005	61	1457	235	0.42 (0.36 to 0.48)
70-79	840	87	1565	389	0.70 (0.61 to 0.80)
>79	440	56	1061	350	1.00
Sex:					
Men	1734	113	3062	559	1.07 (0.95 to 1.19)
Women	1669	109	2288	518	1.00
Diagnosis:					
MI	—	—	5350	1077	2.93 (2.54 to 3.41)
Diabetes	3403	222	—	—	1.00
<b>Hospital admission for myocardial infarction</b>					
Age (years):					
<50	441	11	397	48	1.18 (0.85 to 1.62)
50-59	677	20	870	106	1.25 (0.96 to 1.62)
60-69	1005	49	1457	213	1.61 (1.28 to 1.99)
70-79	840	50	1565	189	1.41 (1.13 to 1.75)
>79	440	12	1061	100	1.00
Sex:					
Men	1734	69	3062	385	1.00 (0.87 to 1.15)
Women	1669	73	2288	271	1.00
Diagnosis:					
MI	—	—	5350	656	3.10 (2.57 to 3.73)
Diabetes	3403	142	—	—	1.00

MI=myocardial infarction.

\*Adjusted for all covariates, derived from Cox regression analysis.

Another concern with the Finnish study was its cross sectional design. Patients were recruited into the study on the basis of existing diabetes (mean duration eight years) and previous myocardial infarction (mean time since infarction not given).<sup>4</sup> It is possible that the rates of cardiovascular disease in the two groups varies with time since diagnosis of diabetes and time since myocardial infarction. Our second study therefore compared rates between patients with newly diagnosed diabetes and those who had just had a myocardial infarction. This comparison may have more clinical relevance. Again, the results showed higher risks of death from all causes, cardiovascular deaths, and hospital admission for myocardial infarction in patients who had had a myocardial infarction compared with patients with diabetes.

### Implications of results

Our results have important implications for clinical practice, in that we should be cautious about basing

treatment decisions on individual risk factors for cardiovascular disease in isolation.<sup>7</sup>

We thank the members of the DARTS/MEMO Collaboration and members of the DARTS Steering Group for their support. MEMO is part of the MRC Health Services Research Collaboration.

Contributors: See [bmj.com](http://bmj.com)

Funding: JMME held a Wellcome Trust training fellowship in health services research (Ref 050212). JW is funded by an Aventis fellowship in health economics.

Competing interests: None declared

- 1 Stamler J, Vaccaro O, Neaton JD, Wentworth D. Diabetes, other risk factors, and 12-year cardiovascular mortality for men screened in the multiple risk factor intervention trial. *Diabetes Care* 1993;16:434-44.
- 2 Stratton IM, Adler AI, Neil AW, Matthews DR, Manley SE, Cull CA, et al. Association of glycaemia with macrovascular and microvascular compli-

cations of type 2 diabetes (UKPDS 35): prospective observational study. *BMJ* 2000;321:405-12.

- 3 Claudi T, Midthjell K, Holmen J, Fougner K, Kruger O, Wiseth R. Cardiovascular disease and risk factors in persons with type 2 diabetes diagnosed in a large population screening: the Nord-Trøndelag diabetes study, Norway. *J Intern Med* 2000;248:492-500.
- 4 Haffner SM, Lehto S, Ronnemaa T, Pyörälä K, Laasko M. Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. *N Engl J Med* 1998;339:229-43.
- 5 Evans JMM, McDevitt DG, MacDonald TM. The Tayside Medicines Monitoring Unit (MEMO): a record-linkage system for pharmacovigilance. *Pharm Med* 1995;9:177-84.
- 6 Morris AD, Boyle DIR, MacAlpine R, Emslie-Smith A, Jung RT, Newton RW, et al. The diabetes audit and research in Tayside, Scotland (DARTS) study: electronic record-linkage to create a diabetes register. *BMJ* 1997;315:524-8.
- 7 Byrne CD, Wild SH. Diabetes care needs evidence based interventions to reduce risk of vascular disease. *BMJ* 2000;320:1554-5.

(Accepted 13 November 2001)



The full version of this paper appears on [bmj.com](http://bmj.com)

## Consumer demand for caesarean sections in Brazil: population based birth cohort study linking ethnographic and epidemiological methods

Dominique P Béhague, Cesar G Victora, Fernando C Barros

Editorial by Langer and p 945

Department of Social Medicine, Federal University of Pelotas, CP 464-96001-970, Pelotas, Rio Grande do Sul, Brazil  
Dominique P Béhague  
*anthropologist*  
Cesar G Victora  
*professor of epidemiology*

Centro Latinoamericano de Perinatología, Montevideo, Uruguay  
Fernando C Barros  
*consultant*

Correspondence to: D P Béhague, Department of Anthropology, McGill University, Montreal, Quebec H3A 2T7, Canada  
[dbehague@aol.com](mailto:dbehague@aol.com)

*BMJ* 2002;324:942-5

### Abstract

**Objectives** To investigate why some women prefer caesarean sections and how decisions to medicalise birthing are influenced by patients, doctors, and the sociomedical environment.

**Design** Population based birth cohort study, using ethnographic and epidemiological methods.

**Setting** Epidemiological study: women living in the urban area of Pelotas, Brazil who gave birth in hospital during the study. Ethnographic study: subsample of 80 women selected at random from the birth cohort. Nineteen medical staff were interviewed.

**Participants** 5304 women who gave birth in any of the city's hospitals in 1993.

**Main outcome measures** Birth by caesarean section or vaginal delivery.

**Results** In both samples women from families with higher incomes and higher levels of education had caesarean sections more often than other women. Many lower to middle class women sought caesarean sections to avoid what they considered poor quality care and medical neglect, resulting from social prejudice. These women used medicalised prenatal and birthing health care to increase their chance of acquiring a caesarean section, particularly if they had social power in the home. Both social power and women's behaviour towards seeking medicalised health care remained significantly associated with type of birth after controlling for family income and maternal education.

**Conclusions** Fear of substandard care is behind many poor women's preferences for a caesarean section. Variables pertaining to women's role in the process of redefining and negotiating medical risks were much stronger correlates of caesarean section rates than income or education. The unequal distribution of medical technology has altered concepts of good and

normal birthing. Arguments supporting interventionist birthing for all on the basis of equal access to health care must be reviewed.

### Introduction

Brazil has a very high rate of caesarean sections<sup>1</sup>; 55% of women from families earning more than \$1000 (£700, €1120) per month had a caesarean section, above the 15% recommended by the World Health Organization.<sup>2,3</sup>

Women's requests for caesarean section may be an important determinant of birth outcome, particularly in countries with growing privatisation and options for patient choice.<sup>4-8</sup> Most research focuses on women's fears of the physiological consequences of vaginal delivery.<sup>5,9</sup> For this reason the debate has focused on providing consumers with knowledge on the risks associated with vaginal and operative deliveries so that decisions about birth may be rationally informed.<sup>10-14</sup> But how do women and obstetricians conclude that the risk of attempting a vaginal delivery is too high?<sup>13,14</sup>

We investigated why some women seek out medicalised care. In particular, we explored how medicalisation is linked to social power in a society where the everyday experience of inequalities is profound.

### Methods

We used a linked ethnographic and epidemiological approach for our study. For the ethnographic study we selected 80 mothers from a birth cohort of 5304 women who gave birth in 1993 in any of the hospitals in Pelotas, Brazil.<sup>2</sup> We interviewed women two or three times during the first five months after birth, using a semistructured guide to explore women's health seeking behaviour, interactions with medical institutions, social integration and activities, and dynamics within