



The full version of
this paper appears
on bmj.com

Reliability of N-terminal pro-brain natriuretic peptide assay in diagnosis of heart failure: cohort study in representative and high risk community populations

F D R Hobbs, R C Davis, A K Roalfe, R Hare, M K Davies, J E Kenkre

Department of
Primary Care and
General Practice,
Primary Care
Clinical Sciences
Building, University
of Birmingham,
Birmingham
B15 2TT

F D R Hobbs
head of department
R C Davis
clinical research fellow
A K Roalfe
statistician
R Hare
research associate
J E Kenkre
senior research fellow

Department of
Cardiology, Selly
Oak Hospital,
Birmingham
B29 2PE
M K Davies
*consultant
cardiologist*

Correspondence to:
F D R Hobbs
[f.d.r.hobbs@
bham.ac.uk](mailto:f.d.r.hobbs@bham.ac.uk)

BMJ 2002;324:1498–500

Abstract

Objective To investigate the performance of a novel assay for N-terminal pro-brain natriuretic peptide (NT-proBNP) in diagnosing heart failure in various randomly selected general and high risk community populations.

Design Community cohort study (substudy of the echocardiographic heart of England screening study).

Setting Four randomly selected general practices in the West Midlands of England.

Participants 591 randomly sampled patients over the age of 45, stratified for age and socioeconomic status and falling into four cohorts (general population, patients with an existing clinical label of heart failure, patients prescribed diuretics, and patients deemed at high risk of heart failure).

Main outcome measure Sensitivity, specificity, positive and negative predictive values, likelihood ratios, and area under receiver operating characteristic curve for NT-proBNP assay in the diagnosis of heart failure.

Results For NT-proBNP in the diagnosis of heart failure in the general population (population screen), a level of >36 pmol/l had a sensitivity of 100%, a specificity of 70%, a positive predictive value of 7%, a negative predictive value of 100%, and an area under the receiver operating characteristic curve of 0.92 (95% confidence interval 0.82 to 1.0). Similar negative predictive values were found for patients from the three other populations screened.

Conclusions This NT-proBNP assay seems to have value in the diagnosis of heart failure in the community. High negative predictive values indicate that the assay's chief use would be to rule out heart failure in patients with suspected heart failure with normal concentrations of NT-proBNP. Positive results may identify patients who need cardiac imaging.

Introduction

Heart failure is an increasingly common disorder,¹ with a prevalence of around 2% in developed countries.² It is characterised by poor prognosis and quality of life for patients,^{3 4} and is responsible for high healthcare costs. As outcomes in heart failure are linked to the stage of disease, early and accurate diag-

nosis is needed to guide appropriate treatment strategies. Despite the need for accurate diagnosis many doctors, especially in primary care, rely on diagnosing heart failure on clinical grounds alone.⁵

Clinical diagnosis of heart failure is not possible on the basis of European and American guidelines,^{6 7} as objective evidence of cardiac dysfunction is needed, along with the presence of appropriate symptoms, before a diagnosis is confirmed. Improving the reliability of diagnosis is essential, as determining the aetiology and stage of heart failure leads to different management choices, such as angiotensin converting enzyme inhibitors and β blockers in most patients with left ventricular systolic dysfunction,^{8 9} spironolactone in patients with severe heart failure,¹⁰ or surgery if appreciable valve disease exists. These treatments improve symptoms, quality of life, and prognosis of disease and reduce healthcare utilisation and costs.

Symptomatic heart failure has a major impact on patients and healthcare systems: its prognosis is worse than those of breast cancer or prostate cancer, and its treatment costs are second only to those for stroke, mainly owing to high admission rates.¹¹ If diagnosis is made early, when there may be no symptoms, treatment can delay or reverse progression of disease.¹² Diagnostic methods may therefore need to encompass screening strategies,^{13 14} as well as identification of patients with symptoms.

The investigation most commonly used to confirm the diagnosis of heart failure is echocardiography, which offers a detailed, semiobjective assessment of ventricular and valve function. However, access to echocardiography is limited in many healthcare systems,⁵ especially for primary care physicians, who see most potential cases. Not surprisingly, the validity of a clinical diagnosis of heart failure in primary care is poor, with high rates of misdiagnosis when patients are assessed against objective criteria, ranging from 25% to 50% accuracy.^{15 16}

Electrocardiography is another option, as a normal recording will, in most cases, exclude left ventricular dysfunction.¹⁷ However, changes on the electrocardiogram may be subtle, and primary care physicians¹⁸ (and hospital doctors¹⁹) are unreliable in assessing such changes.

One potential diagnostic aid is the assessment of patients by assay of brain natriuretic peptide. Brain natriuretic peptide is a biologically active peptide of 32

European Society of Cardiology criteria for diagnosis of heart failure

To satisfy the diagnosis of heart failure, there must be:

- Appropriate symptoms of heart failure
- Objective evidence of cardiac dysfunction on echocardiography and electrocardiography
- Appropriate response to relevant treatment, in cases of doubt

Echocardiography is the most practical tool to demonstrate cardiac dysfunction

amino acids, with vasodilator and natriuretic properties, which is cleaved from the 108 amino acid pro-brain natriuretic peptide released from the cardiac ventricles in response to stretching of the chamber. The second remnant after cleavage, N-terminal pro-brain natriuretic peptide (NT-proBNP), is a 76 amino acid peptide with no known biological function, which circulates at higher concentrations than brain natriuretic peptide and may represent cardiac status over longer periods. Studies of brain natriuretic peptide to date have been small, and data in different subgroups of patients are conflicting.^{20 21} No reliable data exist on the performance of assays for brain natriuretic peptide in the diagnosis of heart failure in the general population. NT-proBNP has been even less investigated.¹⁶ This study provides original data on the utility of a novel NT-proBNP assay in the diagnosis of heart failure.

Methods

This is a prospective substudy of the echocardiographic heart of England screening (ECHOES) study of the prevalence of left ventricular dysfunction and heart failure.² We conducted this substudy in four practices, containing 607 consecutively screened patients randomly selected from the four target groups (307 in the general population, 103 with an existing clinical diagnosis of heart failure, 87 taking diuretics, and 134 at high risk of heart failure, with some sampled from more than one cohort).

We screened patients by history, New York Heart Association functional class, clinical examination, quality of life (SF-36 health status questionnaire), spirometry, resting 12 lead electrocardiography, and echocardiography, including Doppler studies. We diagnosed heart failure on the basis of the agreed gold standard of the European and American guideline criteria (box). A research fellow took blood for the peptide assay after participants had rested supine for five minutes. Blood

was kept at room temperature for up to 24 hours before transport to a local laboratory for centrifugation and freezing of plasma to -20°C . A central independent laboratory performed the NT-proBNP assays.

Results

Five hundred and ninety one (97%) of 607 eligible patients consented to supply a blood sample. Table 1 gives the overall performance characteristics of the assay.

The assay detected definite heart failure in a total of 52 people from the four diagnostic groups—seven from the random general population, 35 from the patients with an existing diagnosis of heart failure, 14 from patients on diuretic treatment, and nine in the high risk group (13 patients were in more than one group). The most frequent causes of heart failure were left ventricular systolic dysfunction, atrial fibrillation, and significant valve disease, or combinations of the three.

In the general population sample, an NT-proBNP concentration of >36 pmol/l had a sensitivity of 100%, a specificity of 70%, a positive predictive value of 7%, and a negative predictive value of 100%. The likelihood ratio of a positive test result was 3.37, and the likelihood ratio of a negative result was 0 (table 1). Similar performance characteristics for diagnosis of heart failure were found when we used the same cut-off values in the other three screened population cohorts. Negative predictive values ranged from 97% to 100%, and likelihood ratios of a negative result ranged from 0 to 0.18 (table 1).

Discussion

Although several studies on assays for brain natriuretic peptide in select patient groups have been published, these are the first data on the performance characteristics of an assay for NT-proBNP in a large generalisable series of randomly selected adults with validated diagnoses of heart failure and with a comparator normative population randomly selected from the same populations as the cases. The assay was also used under the same conditions and constraints as occur in routine clinical practice.

We have shown NT-proBNP assays to be of value in diagnosing heart failure within a representative community based general adult population and also within at risk populations. Importantly, we have shown that the natriuretic peptide assay retains its utility despite samples being handled as they would be in the everyday practice of primary care—that is, collection by

Table 1 Area under receiver operating characteristic curve, sensitivity, specificity, predictive values, and likelihood ratios for NT-proBNP, with a cut-off value of 36 pmol/l, in the diagnosis of definite heart failure. Values in parentheses are 95% confidence intervals

Cohort sampled	Area under curve	Sensitivity (%)	Specificity (%)	Positive predictive value (%)	Negative predictive value (%)	Likelihood ratio of positive result	Likelihood ratio of negative result
General population aged over 45 (n=307)	0.92 (0.82 to 1)	100 (65 to 100)	70 (65 to 75)	7 (3 to 14)	100 (99 to 100)	3.37 (2.83 to 4.01)	0 (0 to 0.89)
Patients with existing diagnosis of heart failure (n=103)	0.8 (0.72 to 0.88)	100 (92 to 100)	18 (10 to 29)	39 (28 to 49)	100 (78 to 100)	1.21 (1.09 to 1.36)	0 (0 to 0.90)
Patients taking diuretics (n=87)	0.87 (0.76 to 0.99)	93 (66 to 100)	40 (28 to 52)	23 (13 to 36)	97 (83 to 100)	1.54 (1.22 to 1.95)	0.18 (0.03 to 1.21)
Patients at high risk for heart failure (n=133)	0.84 (0.76 to 0.93)	100 (72 to 100)	44 (35 to 54)	12 (5 to 21)	100 (96 to 100)	1.8 (1.54 to 2.1)	0 (0 to 1.16)

What is already known on this topic

Many primary care doctors rely on clinical diagnosis of heart failure, although the validity of such diagnosis is poor

Electrocardiography can be used to triage patients for echocardiography, but specialist interpretation may be needed

Assay of brain natriuretic peptide is a potential aid in the diagnosis of heart failure

What this study adds

Assay of N-terminal pro-brain natriuretic peptide (NT-proBNP) seems to be a reliable test for ruling out heart failure

Normal concentrations virtually exclude the diagnosis of heart failure, and very high levels effectively diagnose the condition; intermediate values require confirmation by echocardiography

Assay of NT-proBNP has potential as part of a diagnostic triage in patients presenting with symptoms suggestive of heart failure or in screening populations at high risk

practising nurses and storage at room temperature for some hours before transfer to laboratory settings or freezing at domestic freezer temperatures. Previous studies have mostly used specialised in-hospital handling of samples, including immediate chilled centrifugation and rapid freezing of samples to -70°C .

Another important finding is that NT-proBNP levels are elevated in all major causes of heart failure—left ventricular systolic dysfunction, atrial fibrillation, and valve disease. However, we provide no data on the utility of the assay in diagnosing diastolic heart failure.

The very high negative predictive values, and corresponding low likelihood ratios of a negative result, of the assay make heart failure very unlikely with concentrations below the cut-off value, suggesting that the most appropriate use of the assay in routine practice would be as a rule out or exclusion test. The low likelihood ratios of a negative result are comparable to those for colposcopy, which range from 0.002 to 0.38.²² Elevated concentrations of NT-proBNP should therefore trigger further cardiac investigation, including echocardiography. If left ventricular systolic dysfunction is excluded, elevated concentrations of NT-proBNP may indicate other problems, such as diastolic dysfunction,²³ left ventricular hypertrophy, unstable angina, or pulmonary hypertension.²⁴

These data suggest that, in clinical practice, the assay would have three practical uses: screening patients with existing clinical labels of heart failure (70 of the 103 patients so categorised in this study had heart failure ruled out on NT-proBNP testing); triaging patients presenting with symptoms suggestive of heart failure (shortness of breath, lethargy) for echocardiography; and screening patients at high risk of heart failure. We suspect the assay would perform well in these settings, but the first indication was not formally tested in this study, and the third indication was tested in only 134 patients. These data are sufficiently positive to justify formal trials of the assays in real life strategies for screening symptomatic patients and those at high risk.

Contributors: See bmj.com

Funding: The ECHOES study was funded by the NHS R&D cardiovascular disease and stroke programme. The costs of collecting and handling the samples were supported by an NHS Regional R&D grant, and the assays and technical support were donated by J Trawinski and J Baumann of Roche Diagnostic Solutions.

Competing interests: FDRH is a member of the European Society of Cardiology (ESC) Working Party on Heart Failure, chair of the British Primary Care Cardiovascular Society, and Treasurer of the British Society for Heart Failure. MKD is chairman of the British Society for Heart Failure. FDRH and MKD have received travel sponsorship and honorariums from several biotechnology and pharmaceutical companies with cardiovascular products for plenary talks and attendance at major cardiology scientific congresses and conferences.

- Bonneaux L, Barendregt JJ, Meeter K, Bonsel GJ, van der Maas PJ. Estimating clinical morbidity due to ischemic heart disease and congestive heart failure: the future rise of heart failure. *Am J Public Health* 1994;84:20-8.
- Davies MK, Hobbs FDR, Davis RC, Kenkre JE, Roalfe AK, Hare R, et al. Prevalence of left ventricular systolic dysfunction and heart failure in the general population: main findings from the ECHOES (echocardiographic heart of England screening) study. *Lancet* 2001;358:439-45.
- Ho KKL, Anderson KM, Kannel WB, Grossman W, Levy D. Survival after onset of congestive heart failure in Framingham heart study subjects. *Circulation* 1993;88:107-15.
- Davis R, Hobbs FDR, Kenkre JE, Roalfe AK, McLeod S, Hare R, et al. Quality of life in heart failure, as measured by SF-36 health status questionnaire. *Eur Heart J* 1998;19:S639.
- Hobbs FDR, Jones MI, Allan TF, Wilson S, Tobias R. European survey of primary care physician perceptions on heart failure diagnosis and management (Euro-HF). *Eur Heart J* 2000;21:1877-87.
- The Task Force on Heart Failure of the European Society of Cardiology. Guidelines for the diagnosis of heart failure. *Eur Heart J* 1995;16:741-51.
- American College of Cardiology, American Heart Association. Guidelines for the evaluation and management of heart failure. Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Evaluation and Management of Heart Failure). *Circulation* 1995;92:2764-84.
- Garg R, Yusuf S for the Collaborative Group on ACE Inhibitor Trials. Overview of randomised trials of angiotensin-converting enzyme inhibitors on mortality and morbidity in patients with heart failure. *JAMA* 1995;273:1450-56.
- CIBIS-II Investigators and Committees. The cardiac insufficiency bisoprolol study II. *Lancet* 1999;353:9-13.
- Pitt B, Zannad F, Remm WJ, Cody R, Castaigne A, Perez A, et al for the Randomised Aldactone Evaluation Study Investigators. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. *N Engl J Med* 1999;341:709-17.
- Gillum RF. Heart and stroke facts. *Am Heart J* 1993;126:1042-7.
- Pfeffer MA, Braunwald E, Moy LA, Basta L, Brown EJ, Cuddy TE for the SAVE Investigators. Effect of captopril on mortality and morbidity in patients with left ventricular dysfunction after myocardial infarction: results of the survival and ventricular enlargement trial. *N Engl J Med* 1992;327:669-77.
- Hobbs FDR, Davis RC, McLeod S, Marshall T, Kenkre J, Lancashire R, et al. Prevalence of heart failure in "high risk" groups. *J Am Coll Cardiol* 1998;31(S5):85C.
- Hobbs FDR. Primary care physicians: champions of or an impediment to optimal care of the patient with heart failure? *Eur J Heart Fail* 1999;1:11-5.
- Agency for Health Care Policy and Research. *Heart failure: evaluation and care of patients with left-ventricular systolic dysfunction*. Rockville, MD: Agency for Health Care Policy and Research, 1994.
- Hunt PJ, Richards AM, Nicholls MG, Yandle TG, Doughty RN, Espiner EA. Immunoreactive amino-terminal pro brain natriuretic peptide (NT-proBNP): a new marker of cardiac impairment. *Clin Endocrinol* 1997;47:287-96.
- Davie AP, Francis CM, Love MP, Caruana L, Starkey IR, Shaw TRD, et al. Value of an electrocardiogram in identifying heart failure due to left ventricular systolic dysfunction. *BMJ* 1996;312:222.
- Sur DK, Kaye L, Mikus M, Goad J, Morena A. Accuracy of electrocardiogram reading by family practice residents. *Fam Med* 2000;32:315-9.
- Weston MJ, Bett JHN, Over R. Consensus opinion and observer accuracy in electrocardiography with reference to coronary arteriographic information. *Aust NZ J Med* 1976;6:429-32.
- McClure SJ, Caruana L, Davie AP, Goldthorp S, McMurray JJ. Cohort study of plasma natriuretic peptides for identifying left ventricular systolic dysfunction in primary care. *BMJ* 1998;317:516-9.
- Landray MJ, Lehman R, Arnold I. Measuring brain natriuretic peptide in suspected left ventricular systolic dysfunction in general practice: cross-sectional study. *BMJ* 2000;320:985-6.
- Mitchell MF, Schottenfeld D, Tortolero-Luna G, Cantor SB, Richards-Kortum R. Colposcopy for the diagnosis of squamous intraepithelial lesions: a meta-analysis. *Obstet Gynecol* 1998;91:626-31.
- Fruhwald F, Fahrleitner A, Watzinger N, Fruhwald S, Dobnig H, Schumacher M, et al. Natriuretic peptides in patients with diastolic dysfunction due to dilated cardiomyopathy. *Eur Heart J* 1999;20:1415-23.
- Nagaya N, Nishikimi T, Okano Y, Uematsu M, Satoh T, Kyotani S, et al. Plasma brain natriuretic peptide levels increase in proportion to the extent of right ventricular dysfunction in pulmonary hypertension. *J Am Coll Cardiol* 1998;31:202-8.

(Accepted 4 April 2002)