

## A strategy to reduce cardiovascular disease by more than 80%

N J Wald, M R Law

### Abstract

**Objectives** To determine the combination of drugs and vitamins, and their doses, for use in a single daily pill to achieve a large effect in preventing cardiovascular disease with minimal adverse effects. The strategy was to simultaneously reduce four cardiovascular risk factors (low density lipoprotein cholesterol, blood pressure, serum homocysteine, and platelet function) regardless of pretreatment levels.

**Design** We quantified the efficacy and adverse effects of the proposed formulation from published meta-analyses of randomised trials and cohort studies and a meta-analysis of 15 trials of low dose (50-125 mg/day) aspirin.

**Outcome measures** Proportional reduction in ischaemic heart disease (IHD) events and strokes; life years gained; and prevalence of adverse effects.

**Results** The formulation which met our objectives was: a statin (for example, atorvastatin (daily dose 10 mg) or simvastatin (40 mg)); three blood pressure lowering drugs (for example, a thiazide, a  $\beta$  blocker, and an angiotensin converting enzyme inhibitor), each at half standard dose; folic acid (0.8 mg); and aspirin (75 mg). We estimate that the combination (which we call the Polypill) reduces IHD events by 88% (95% confidence interval 84% to 91%) and stroke by 80% (71% to 87%). One third of people taking this pill from age 55 would benefit, gaining on average about 11 years of life free from an IHD event or stroke. Summing the adverse effects of the components observed in randomised trials shows that the Polypill would cause symptoms in 8-15% of people (depending on the precise formulation).

**Conclusion** The Polypill strategy could largely prevent heart attacks and stroke if taken by everyone aged 55 and older and everyone with existing cardiovascular disease. It would be acceptably safe and with widespread use would have a greater impact on the prevention of disease in the Western world than any other single intervention.

### Introduction

Heart attacks, stroke, and other preventable cardiovascular diseases kill or seriously affect half the population of Britain. Randomised trials show that drugs to lower three risk factors—low density lipoprotein (LDL) cholesterol,<sup>1</sup> blood pressure,<sup>2-6</sup> and platelet function (with

aspirin)<sup>7 8</sup>—reduce the incidence of ischaemic heart disease (IHD) events and stroke. Evidence that lowering serum homocysteine (with folic acid) reduces the risk of these diseases is largely observational but still compelling.<sup>9 10</sup>

Drug treatment to prevent IHD events and stroke has generally been limited to single risk factors, to targeting the minority of patients with values in the tail of the risk factor distribution, and to reducing the risk factors to “average” population values. This policy can achieve only modest reductions in disease.<sup>11</sup> A large preventive effect would require intervention in everyone at increased risk irrespective of the risk factor levels; intervention on several reversible causal risk factors together; and reducing these risk factors by as much as possible.<sup>11</sup>

We describe a strategy to prevent cardiovascular disease based on these three principles<sup>12</sup>: a daily treatment, the Polypill, containing six components, each lowering one of the above four risk factors. We also quantify its overall preventive effect. The pill would be suitable for people with cardiovascular disease and for everyone over a specified age (say 55).

### Methods

We identified categories of drugs or vitamins used to modify LDL cholesterol, blood pressure, homocysteine, and platelet function. For LDL cholesterol, statins are the drugs of choice.<sup>1 13 14</sup> For lowering blood pressure, we considered all five main categories of drugs: thiazides,  $\beta$  blockers, angiotensin converting enzyme (ACE) inhibitors, angiotensin II receptor antagonists, and calcium channel blockers.<sup>13</sup> Serum homocysteine is most effectively reduced by folic acid.<sup>15</sup> Aspirin is the most widely used and least expensive antiplatelet agent.

The choices of statin and of the categories and doses of blood pressure lowering drugs were determined from the meta-analysis of short term randomised trials in our companion papers.<sup>1 16</sup> The dose of folic acid was the minimum needed to ensure the maximum reduction in serum homocysteine.<sup>15 17</sup> The long term effect of a specified absolute reduction in LDL cholesterol, blood pressure, and homocysteine expressed as the proportional reduction in the incidence of IHD events and stroke was based on systematic reviews of cohort studies.<sup>1 9 14 18</sup> Both cohort studies and trials have shown that a specified reduction



This is an abridged version; the full version is on [bmj.com](http://bmj.com)

Editorial by Rodgers

Department of Environmental and Preventive Medicine, Wolfson Institute of Preventive Medicine, Barts and the London, Queen Mary's School of Medicine and Dentistry, University of London, London EC1M 6BQ

N J Wald  
professor  
M R Law  
professor

Correspondence to:  
N J Wald  
[n.j.wald@qmul.ac.uk](mailto:n.j.wald@qmul.ac.uk)

BMJ 2003;326:1419-23



Further tables appear on [bmj.com](http://bmj.com)

**Table 1** Effects of the Polypill on the risks of ischaemic heart disease (IHD) and stroke after two years of treatment at age 55-64

Risk factor	Agent	Reduction in risk factor	% reduction in risk (95% CI)*		Source of evidence
			IHD event	Stroke	
LDL cholesterol	Statin†	1.8 mmol/l (70 mg/dl) reduction in LDL cholesterol	61 (51 to 71)	17 (9 to 25)	Law et al <sup>1</sup>
Blood pressure	Three classes of drug at half standard dose	11 mm Hg diastolic	46 (39 to 53)	63 (55 to 70)	Law et al <sup>16</sup>
Serum homocysteine	Folic acid (0.8 mg/day)	3 µmol/l	16 (11 to 20)	24 (15 to 33)	Wald et al <sup>9</sup>
Platelet function	Aspirin (75 mg/day)	Not quantified	32 (23 to 40)	16 (7 to 25)	Table A on bmj.com
Combined effect	All		88 (84 to 91)	80 (71 to 87)	

LDL=low density lipoprotein.

\*95% confidence intervals include imprecision of the estimates of both the agent reducing the risk factor and the risk factor reduction decreasing risk.

†Atorvastatin 10 mg/day, or simvastatin or lovastatin 40 mg/day taken in the evening or 80 mg/day taken in the morning.

in blood pressure or serum cholesterol produces a constant proportional reduction in risk that is independent of the initial value of the risk factor.<sup>1 3 4 11 14</sup> The average age at which cardiovascular event occurred in the studies was around 60-65 years.

We used direct evidence from randomised trials of the effects of low dose aspirin on disease events. Since the necessary information on stroke and adverse effects was not available from published meta-analyses, we conducted one. We identified trials of ≥6 months' duration from previous meta-analyses,<sup>7 8</sup> from Medline, and the Cochrane Collaboration and Web of Science databases. This yielded 15 trials. We determined the average proportional reduction in IHD events and stroke, and the prevalence and incidence of adverse effects.

We calculated the combined effect of changing the four risk factors (the effect of the Polypill) by multiplying the relative risks associated with each. Using a simple Markov model, we calculated the years of life gained without a heart attack or stroke if people without a previous cardiovascular event used the Polypill from age 55.

## Results

### Efficacy

Table 1 shows the effects of the individual agents. By use of statins, LDL cholesterol concentration can be reduced by an average of 1.8 mmol/l. Atorvastatin 10 mg taken at any time of day or simvastatin (or lovastatin) 40 mg taken in the evening or 80 mg taken in the morning after about two years of treatment can reduce the incidence of IHD events at age 60 by an estimated 61%.<sup>1</sup> The overall reduction in stroke from an LDL cholesterol reduction of 1.8 mmol/l is about 17%.<sup>1</sup>


The five main categories of blood pressure lowering drugs (thiazide, β blockers, ACE inhibitors, angiotensin II receptor antagonists, and calcium channel blockers), and the individual drugs within the categories, produce similar reductions in blood pressure, given dose as a ratio of standard dose.<sup>16</sup> A combination of three drugs from different categories in low dose has greater efficacy and fewer adverse effects than using one or two drugs in standard dose.<sup>16</sup> The blood pressure reduction with three drugs in combination at half standard dose is about 11 mm Hg diastolic, reducing the incidence of IHD events by 46% and stroke by 63%.<sup>16 17</sup>

The maximum effect of folic acid, achieved at a dose of about 0.8 mg/day,<sup>15 18</sup> lowers serum homo-

cysteine by 3 µmol/l (about 25%) and reduces IHD events by about 16% and stroke by 24%.<sup>9</sup>

Figure 1 shows our meta-analysis of the 15 randomised trials of low dose aspirin (50-125 mg/day). IHD events were reduced by 32% and strokes by 16% (details in web table A).

Table 1 shows that changing all four risk factors together reduces the risk of IHD events by 88% and stroke by 80%. These results are obtained by multiplying the relative risk estimates relating to the intervention on each risk factor, which is the complement of the proportion of events prevented;



**Fig 1** Relative risks (95% confidence intervals) of ischaemic heart disease events and all strokes (fatal and non-fatal) in 15 randomised trials of low dose aspirin

thus, preventing, say, 61% is equivalent to a relative risk of 0.39. The following example illustrates the calculation. The relative risks of an IHD event for the four interventions in table 1 are 0.39, 0.54, 0.84, and 0.66, the product of which yields a combined relative risk of 0.12 or an 88% preventive effect (if 100 people who would have had IHD events without intervention were treated, statins would prevent 61 of the 100 events, leaving 39; 46% of these would be prevented with blood pressure lowering drugs, leaving 21; 16% of these would be prevented with folic acid, leaving 18; and 34% of these would be prevented with aspirin, leaving 12; 88% have thus been prevented). Reducing one risk factor has a similar proportional effect on risk irrespective of the level of other risk factors, as confirmed by cohort studies and randomised trials.<sup>19-22</sup>

For example, trials of LDL cholesterol reduction show similar proportional reductions in risk in people with high and low blood pressure and in people taking and not taking aspirin.<sup>20, 21</sup>

Other than the statin (in respect of IDH), omitting a single component has a relatively minor impact on the combined effect of the residual components, illustrating the robustness of the Polypill concept. Compared with the reductions in IHD events and stroke of 88% and 80% respectively with all six components, the reductions were 86% and 74% without folic acid, 85% and 73% without one blood pressure lowering drug (two instead of three), and 83% and 77% without aspirin. So, for example, aspirin prevents 32% of IHD events when used alone but prevents only an additional 5% of the original number of expected events when added to the other components in the combination.

Table 2 shows the expected proportion of people who would avoid an IHD event or stroke by taking the Polypill from age 55 and, in those, the average number of event-free life years gained. The estimates take account of deaths from causes other than IHD and stroke. About a third of people taking the Polypill would benefit. On average each will gain 11-12 years of life free from a heart attack or stroke. The gain in life is substantial at all ages.

#### Adverse effects

See [bmj.com](http://bmj.com) for a table of the extracranial adverse effects of low dose aspirin from our meta-analysis of 15 randomised trials. Table 3 uses these data together with those published in our companion papers<sup>1, 16</sup> to show the proportions of people reporting symptoms attributable to any of the components of the Polypill (percentage with symptoms in treated groups minus percentage in placebo groups in trials). If we included the three classes of blood pressure lowering drugs with the lowest prevalence of adverse effects (thiazide, angiotensin II receptor antagonist, and calcium channel blocker<sup>16</sup>) in a Polypill formulation, 8% would be expected to have symptoms attributable to one or more of the six components of the pill, mostly due to aspirin. If we used the three least expensive blood pressure lowering drugs (a thiazide, a  $\beta$  blocker, and an ACE inhibitor) instead, a Polypill including these would cause symptoms in about 15% of people taking the pill.

Of all the components, aspirin has the most serious adverse effects, mainly due to haemorrhage (see tables

**Table 2** Expected benefits in 100 men and 100 women without a known vascular disease who start taking the Polypill at age 55. Calculations are based on a Markov model and allow for other causes of death

Age (years)	Men		Women	
	No who benefit*	Years gained†	No who benefit*	Years gained†
Up to 65	7	21	3	24
Up to 75	18	16	11	18
Up to 85	30	13	24	14
Up to any age	36	12	35	11

\*Ischaemic heart disease event or stroke avoided or delayed.

†Average years of life gained without an IHD event or stroke in those who benefit.

**Table 3** Prevalence of participants in randomised trials reporting symptoms attributable to the Polypill components (in doses specified in table 1)

Drug or vitamin	% of participants with symptom*	
	Any symptoms	Symptoms sufficient to stop treatment in short term trials
Statin <sup>1</sup>	0.1	<0.1
Thiazide <sup>16</sup>	2.0	0.1
Angiotensin II receptor antagonist	<0.1	<0.1
Calcium channel blocker <sup>16</sup>	1.6	0.1
Folic acid	<<0.1	<<0.1
Aspirin	3.9	1.6

\*Percentage in treated group minus percentage in placebo group.

on [bmj.com](http://bmj.com)). In our meta-analysis of the trials of low dose aspirin the increase in haemorrhagic stroke (table A on [bmj.com](http://bmj.com)) was exceeded by the reduction in thrombotic strokes, producing an overall 16% reduction in stroke. There was no excess risk of fatal extracranial haemorrhage, with 13 and 15 deaths in the aspirin and placebo groups respectively in about 17 000 people in each (table B on [bmj.com](http://bmj.com)), and an excess risk of major non-fatal extracranial haemorrhage (mainly gastric) of 1.2 per 1000 person years (tables on [bmj.com](http://bmj.com)).

## Discussion

The Polypill strategy, based on a single daily pill containing six components as specified, would prevent 88% of heart attacks and 80% of strokes. About 1 in 3 people would directly benefit, each on average gaining 11-12 years of life without a heart attack or stroke (20 years in those aged 55-64).

We are confident that the estimated effect is accurate. There is substantial evidence on the individual components of the Polypill, both for risk factor reduction and disease reduction. Extensive evidence exists that reducing the four risk factors by any means lowers the risk of cardiovascular disease. The consistency between evidence from observational studies and trials is persuasive.

The percentage reduction in stroke will be greater for non-fatal than fatal events (about 82% and about 75%, respectively) because statins and aspirin have different effects on thrombotic and haemorrhagic stroke and haemorrhagic strokes are more often fatal.

#### Who should take the Polypill

In people with a previous heart attack or a stroke, without any treatment, cardiovascular disease mortality is about 5% per year for life.<sup>23</sup> About half of all cardiovascular deaths occur in individuals with a previous myocardial infarction or cerebral thrombosis. All such individuals should be offered treatment to reduce the

reversible risk factors and would benefit from taking the Polypill. Patients with angina pectoris, transient ischaemic attacks, peripheral arterial disease, and diabetes mellitus should also consider taking the Polypill.

Among people without existing disease, the most discriminatory screening factor is age. As 96% of deaths from ischaemic heart disease or stroke occur in people aged 55 and over, treating everyone in this group would prevent nearly all such deaths. Using different age cut-offs for men and women, or smokers and non-smokers, or combining several risk factor values with age and sex to produce individual estimates of overall risk would add little discrimination and would probably not justify the added complexity and cost.<sup>11 24 25</sup> These factors, though aetiologically important, are poor predictors of future cardiovascular disease events (see [bmj.com](http://bmj.com) for a figure illustrating this with serum cholesterol, blood pressure, and serum homocysteine). There is little separation between the distributions of the risk factors in people who over a specified period do or do not have a disease event. Cut-off levels that identify the 5% of the unaffected population who have the most extreme values of the risk factors identify only about 15% of the disease events (24% for blood pressure and stroke). The screening performance of cardiovascular risk factors in combination is little better.<sup>22</sup>

The best approach is therefore to treat people with known occlusive vascular disease and everyone aged 55 and over. There is no need to measure the four risk factors before starting treatment, because intervention is effective whatever the initial levels of the risk factors,<sup>11</sup> nor to monitor the effect of the treatment, because fluctuations within individuals tend to mask variations between individuals in the systematic effects of the interventions.

#### Adverse effects

The Polypill may not be suitable for some people.  $\beta$  blockers are unsuitable for people with asthma, and some people are intolerant of aspirin. Monitoring to prevent rare serious adverse effects of treatment might be considered, measuring serum creatine kinase and transaminase (for rhabdomyolysis and hepatitis caused by statins), and serum potassium and creatinine (for acute renal failure caused by ACE inhibitors and angiotensin II receptor antagonists). However, the value of such monitoring is uncertain. The complications are rare, it is not known whether monitoring will avoid them, and the tests lack specificity, so the increased risk of cardiovascular disease after stopping the drug in people positive on monitoring may outweigh any benefit.

#### Cost and acceptability

A low cost Polypill could use generic components that are not subject to patent protection (simvastatin (from mid-2003), hydrochlorothiazide, atenolol, enalapril, folic acid, and aspirin). This formulation does not have the lowest rate of adverse effects, but even if about 10% of people were intolerant of the formulation it would still have considerable public health merit. Those found to be intolerant could be prescribed alternatives to avoid the side effects. Controlled trials of different formulations of the Polypill would provide direct estimates of acceptability.

### What is already known on this topic

Four risk factors (LDL cholesterol, blood pressure, homocysteine, and platelet function) that can be reduced by drugs or vitamins account for most cardiovascular disease

Apart from aspirin, the use of such agents has focused on people with high levels of the risk factor

### What this study adds

Intervening on all four risk factors reduces heart attacks and strokes by over 80%

To achieve this large effect in a population requires a combination treatment taken by everyone above a specified age (say 55) and younger people with a clinical history of occlusive arterial disease

A combination pill containing six active components could be widely used

Each component has been used in medical practice for more than 10 years with substantial evidence on safety and efficacy

### Conclusions

The preventive strategy outlined is radical. But a formulation that prevented all cancer and was safe would undoubtedly be widely used, and one that prevented more than 80% of cardiovascular disease would be even more important, because such deaths are more common than cancer deaths. It is time to discard the view that risk factors need to be measured and treated individually if found to be "abnormal." Instead it should be recognised that in Western society the risk factors are high in us all, so everyone is at risk; that the diseases they cause are common and often fatal; and that there is much to gain and little to lose by the widespread use of these drugs.

We thank Joan Morris and Alicja Rudnicka for statistical help, and Leo Kinlen, Jeffrey Aronson, Mark Caulfield, David Collier, James Haddow, and Frank Speizer for their helpful comments on drafts of this paper. This paper is based on a lecture given by Nicholas Wald on 18 September 2000 at a meeting in Israel of the Israel National Institute for Health Services Policy and Health Services Research.

Contributors: See [bmj.com](http://bmj.com).

Funding: None.

Competing interests: The authors have filed a patent application on the formulation of the combined pill described here (application Nos GB 0100548.7 and GB 008791.6, priority date 10 April 2000) and a trademark application for the name Polypill.

- 1 Law MR, Wald NJ, Rudnicka A. Quantifying effect of statins on low density lipoprotein cholesterol, ischaemic heart disease, and stroke: systematic review and meta-analysis. *BMJ* 2003;326:1423-7.
- 2 MacMahon S, Peto R, Cutler J, Collins R, Sorlie P, Neaton J, et al. Blood pressure, stroke and coronary heart disease. Part 1: prolonged differences in blood pressure: prospective observational studies corrected for the regression dilution bias. *Lancet* 1990;335:765-74.
- 3 Collins R, Peto R, MacMahon S, Hebert P, Fiebach NH, Eberlein KA, et al. Blood pressure, stroke and coronary heart disease. Part 2: short-term reductions in blood pressure: overview of randomised drug trials in their epidemiological context. *Lancet* 1990;335:827-38.
- 4 Law MR, Wald NJ, Morris JK. Lowering blood pressure to prevent myocardial infarction and stroke: a new prevention strategy. *Health Technol Assess* (in press).

- 5 PATS Collaborating Group. Post-stroke antihypertensive treatment study. *Chin Med J* 1995;108:710-17.
- 6 PROGRESS Collaborative Group. Randomised trial of a perindopril-based blood-pressure-lowering regimen among 6105 individuals with previous stroke or transient ischaemic attack. *Lancet* 2001;358:1033-41.
- 7 Antiplatelet Trialists' Collaboration. Collaborative overview of randomised trials of antiplatelet therapy. I: Prevention of death, myocardial infarction, and stroke by prolonged antiplatelet therapy in various categories of patients. *BMJ* 1994;308:81-105.
- 8 Antithrombotic Trialists' Collaboration. Collaborative meta-analysis of randomised trials of antiplatelet therapy for prevention of death, myocardial infarction, and stroke in high risk patients. *BMJ* 2002;324:71-86.
- 9 Wald DS, Law M, Morris JK. Homocysteine and cardiovascular disease: evidence on causality from a meta-analysis. *BMJ* 2002;325:1202-6. (See full version on bmj.com).
- 10 Schnyder G, Roffi M, Pin R, Flammer Y, Lange H, Eberli FR, et al. Decreased rate of coronary restenosis after lowering of plasma homocysteine levels. *N Engl J Med* 2001;345:1593-600.
- 11 Law MR, Wald NJ. Risk factor thresholds: their existence under scrutiny. *BMJ* 2002;324:1570-6.
- 12 Wald NJ, Law MR. Formulation for the prevention of cardiovascular disease. UK patent application No 0008791.6.2000.
- 13 British Medical Association, Royal Pharmaceutical Society of Great Britain. *British national formulary*. London: BMA, RPS, 2002. (No 44.)
- 14 Law MR, Wald NJ, Thompson SG. By how much and how quickly does reduction in serum cholesterol concentration lower risk of ischaemic heart disease? *BMJ* 1994;308:367-72.
- 15 Homocysteine Lowering Trialists Collaboration. Lowering blood homocysteine with folic acid based supplements: meta-analysis of randomised trials. *BMJ* 1998;316:894-8.
- 16 Law MR, Wald NJ, Morris JK, Jordan R. Value of low dose combination treatment with blood pressure lowering drugs: analysis of 354 randomised trials. *BMJ* 2003;326:1427-31.
- 17 Wald DS, Bishop L, Wald NJ, Law M, Hennessy E, Weir D, et al. Randomized trial of folic acid supplementation and serum homocysteine levels. *Arch Intern Med* 2001;161:695-700.
- 18 Prospective Studies Collaboration. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet* 2002;360:1903-13.
- 19 ISIS-2 Collaborative Group. Randomised trial of intravenous streptokinase, oral aspirin, both or neither among 17,187 cases of suspected acute myocardial infarction: ISIS-2. *Lancet* 1988;ii:349-59.
- 20 Sacks FM, Pfeffer MA, Moye LA, Rouleau JL, Rutherford JD, Cole TG, et al. The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. *N Engl J Med* 1996;335:1001-9.
- 21 Heart Protection Study Collaborative Group. MRC/BHF heart protection study of cholesterol lowering with simvastatin in 20,536 high-risk individuals: a randomised placebo-controlled trial. *Lancet* 2002;360:7-22.
- 22 Wald NJ, Law M, Watt H, Wu T, Bailey A, Johnson M, et al. Apolipoproteins and ischaemic heart disease: implications for screening. *Lancet* 1994;343:75-9.
- 23 Law MR, Watt HC, Wald NJ. The underlying risk of death after myocardial infarction in the absence of treatment. *Arch Intern Med* 2002;162:2405-10.
- 24 Wald NJ, Hackshaw AK, Frost CD. When can a risk factor be used as a worthwhile screening test? *BMJ* 1999;319:1562-5.
- 25 Screening brief: screening for ischaemic heart disease by serum homocysteine measurement. *J Med Screen* 2001;8:220.

## Quantifying effect of statins on low density lipoprotein cholesterol, ischaemic heart disease, and stroke: systematic review and meta-analysis

M R Law, N J Wald, A R Rudnicka



This is an abridged version; the full version is on bmj.com

### Abstract

**Objectives** To determine by how much statins reduce serum concentrations of low density lipoprotein (LDL) cholesterol and incidence of ischaemic heart disease (IHD) events and stroke, according to drug, dose, and duration of treatment.

**Design** Three meta-analyses: 164 short term randomised placebo controlled trials of six statins and LDL cholesterol reduction; 58 randomised trials of cholesterol lowering by any means and IHD events; and nine cohort studies and the same 58 trials on stroke.

**Main outcome measures** Reductions in LDL cholesterol according to statin and dose; reduction in IHD events and stroke for a specified reduction in LDL cholesterol.

**Results** Reductions in LDL cholesterol (in the 164 trials) were 2.8 mmol/l (60%) with rosuvastatin 80 mg/day, 2.6 mmol/l (55%) with atorvastatin 80 mg/day, 1.8 mmol/l (40%) with atorvastatin 10 mg/day, lovastatin 40 mg/day, simvastatin 40 mg/day, or rosuvastatin 5 mg/day, all from pretreatment concentrations of 4.8 mmol/l. Pravastatin and fluvastatin achieved smaller reductions. In the 58 trials for an LDL cholesterol reduction of 1.0 mmol/l the risk of IHD events was reduced by 11% in the first year of treatment, 24% in the second year, 33% in years three to five, and by 36% thereafter ( $P < 0.001$  for trend). IHD events were reduced by 20%, 31%, and 51% in trials grouped by LDL cholesterol reduction (means 0.5 mmol/l, 1.0 mmol/l, and 1.6 mmol/l) after

results from first two years of treatment were excluded ( $P < 0.001$  for trend). After several years a reduction of 1.8 mmol/l would reduce IHD events by an estimated 61%. Results from the same 58 trials, corroborated by results from the nine cohort studies, show that lowering LDL cholesterol decreases all stroke by 10% for a 1 mmol/l reduction and 17% for a 1.8 mmol/l reduction. Estimates allow for the fact that trials tended to recruit people with vascular disease, among whom the effect of LDL cholesterol reduction on stroke is greater because of their higher risk of thromboembolic stroke (rather than haemorrhagic stroke) compared with people in the general population.

**Conclusions** Statins can lower LDL cholesterol concentration by an average of 1.8 mmol/l which reduces the risk of IHD events by about 60% and stroke by 17%.

### Introduction

Statins are highly effective in lowering serum cholesterol concentrations and preventing ischaemic heart disease (IHD).<sup>1-3</sup> Three issues remain. We do not know by how much different statins at different doses reduce low density lipoprotein (LDL) cholesterol concentrations. Secondly, the full effect of statins in preventing IHD events has been underestimated because IHD events in the first two years (before the full effect is attained<sup>4</sup>) were not censored, many trials used less effective statins, and trials were affected by non-adherence to the allocated regimen. Thirdly, there

Department of Environmental and Preventive Medicine, Wolfson Institute of Preventive Medicine, Barts and the London, Queen Mary's School of Medicine and Dentistry, University of London, London EC1M 6BQ  
M R Law  
professor  
N J Wald  
professor  
A R Rudnicka  
lecturer

Correspondence to: M R Law  
m.r.law@qmul.ac.uk

BMJ 2003;326:1423-7



A full list of references and two extra tables can be found on bmj.com