

Risk of cancer after low doses of ionising radiation—retrospective cohort study in 15 countries

E Cardis, M Vrijheid, M Blettner, E Gilbert, M Hakama, C Hill, G Howe, J Kaldor, C R Muirhead, M Schubauer-Berigan, T Yoshimura and the international study group

Abstract

Objectives To provide direct estimates of risk of cancer after protracted low doses of ionising radiation and to strengthen the scientific basis of radiation protection standards for environmental, occupational, and medical diagnostic exposures.

Design Multinational retrospective cohort study of cancer mortality.

Setting Cohorts of workers in the nuclear industry in 15 countries.

Participants 407 391 workers individually monitored for external radiation with a total follow-up of 5.2 million person years.

Main outcome measurements Estimates of excess relative risks per sievert (Sv) of radiation dose for mortality from cancers other than leukaemia and from leukaemia excluding chronic lymphocytic leukaemia, the main causes of death considered by radiation protection authorities.

Results The excess relative risk for cancers other than leukaemia was 0.97 per Sv, 95% confidence interval 0.14 to 1.97. Analyses of causes of death related or unrelated to smoking indicate that, although confounding by smoking may be present, it is unlikely to explain all of this increased risk. The excess relative risk for leukaemia excluding chronic lymphocytic leukaemia was 1.93 per Sv (<0 to 8.47). On the basis of these estimates, 1-2% of deaths from cancer among workers in this cohort may be attributable to radiation.

Conclusions These estimates, from the largest study of nuclear workers ever conducted, are higher than, but statistically compatible with, the risk estimates used for current radiation protection standards. The results suggest that there is a small excess risk of cancer, even at the low doses and dose rates typically received by nuclear workers in this study.

Introduction

The 15 country study, an international collaborative study of cancer risk among radiation workers in the nuclear industry, was carried out to directly estimate the risk of cancer after protracted low dose exposures and to strengthen the scientific basis of radiation protection.¹ We present risk estimates for mortality from all cancers, excluding leukaemia, and from leukaemia excluding chronic lymphocytic leukaemia, and compare them with estimates derived from data on survivors of the A bomb. We have used the term nuclear industry to refer to facilities engaged in production of nuclear power, manufacture of nuclear weapons, enrichment and processing of nuclear fuel, production of radioisotopes, or nuclear reactor or weapons research. Uranium mining is not included.

Methods

This multinational retrospective cohort study used a common protocol in 15 countries and collected information on nearly 600 000 workers. Study cohorts were defined from employment and dosimetric records of participating facilities or, where available, from centralised national dose registries. The a priori eligibility criteria for inclusion of cohorts² were essentially complete and non-selective follow-up for mortality; availability of individual annual recorded estimates of dose for all monitored workers; and availability of information on historical monitoring policies and practices. We included all workers who had been monitored for external photon (x and γ) radiation exposure through the use of personal dosimeters. Details of country specific methods are described elsewhere.³

Vital status and cause of death ascertainment

We established vital status through linkage with national or regional death registries or, where this was not possible, appropriate records of local authorities. Completeness of follow-up ranged from 87% to nearly 100%. Vital statistics registries provided cause of death, which was known for over 90% of workers who died.

Adequacy of dosimetric records

We reconstructed each worker's dosimetric history using recorded doses from individual facilities or national dose registries. A study evaluated the comparability of dose estimates across facilities and time³ and quantified sources of bias (see bmj.com). Doses from higher energy photons (100-3000 keV), which constituted most of the dose in most cohorts, were judged to have been measured in a comparable way over time and across facilities.³⁻⁴ We excluded workers with potential for substantial doses ($\geq 10\%$ of their whole body dose) from other radiation types (neutrons, internal exposures).

Main study population

The main study population was workers who had been employed in one or more facilities for at least one year, who had been monitored for external radiation exposure, and whose doses resulted predominantly from higher energy photon radiation (190 677 workers who did not fit these criteria were excluded).

Dosimetric errors and derivation of organ doses

Organ doses were derived by dividing recorded doses by the appropriate organ dose bias factor (see bmj.com). We used doses to the colon and active bone marrow for analyses of mortality from all cancers excluding leukaemia and from leukaemia, respectively.

International Agency for Research on Cancer, Lyon, France
E Cardis
head of radiation group
M Vrijheid
scientist

Institute of Medical Biostatistics, Epidemiology and Informatics, University of Mainz, Germany
M Blettner
professor

Radiation Epidemiology Branch, Division of Epidemiology and Genetics, National Cancer Institute, Bethesda, Maryland, USA
E Gilbert
expert

University of Tampere, Tampere, Finland
M Hakama
research professor

Institut Gustave-Roussy, Villejuif, France
C Hill
epidemiologist
Mailman School of Public Health, Columbia University, New York, USA
G Howe
professor of epidemiology

continued over

BMJ 2005;331:77-80



This is the abridged version of an article that was posted on bmj.com on 29 June 2005: <http://bmj.com/cgi/doi/10.1136/bmj.38499.599861.E0>

National Centre in HIV Epidemiology and Clinical Research, Sydney, Australia

J Kaldor
deputy director

Radiation Protection Division, Health Protection Agency, Chilton, Didcot, Oxfordshire

C R Muirhead
group leader
epidemiology

Division of Surveillance, Hazard Evaluations and Field Studies, National Institute for Occupational Safety and Health, Cincinnati, USA

M Schubauer-Berigan
senior research epidemiologist,
industrywide studies branch

Fukuoka Institute of Health and Environmental Sciences, Fukuoka, Japan

T Yoshimura
director

Details of members of the international study group can be found on bmj.com

Correspondence to: E Cardis
cardis@iarc.fr

Statistical methods

Analyses were based on a linear relative risk Poisson regression model (see bmj.com). Analyses used only underlying cause of death. To allow for a latent period between exposure and death, doses were lagged by two years for leukaemia and 10 years for other cancers. Sensitivity analyses were conducted with a range of different lags.

We focused on the main causes of death for which radiation protection committees have provided risk estimates: all cancers excluding leukaemia and leukaemia excluding chronic lymphocytic leukaemia. Chronic lymphocytic leukaemia is thought to be less inducible by ionising radiation than other leukaemias.⁵ We have also presented risk estimates for solid cancers to compare with recent data for A bomb survivors⁶ and for all cancers excluding leukaemia, lung, and pleural cancers (which have the greatest potential for confounding by smoking, internally incorporated radionuclides, and other occupational carcinogens). We investigated confounding by smoking by separately analysing solid cancers related or unrelated to smoking and two groupings of smoking related outcomes other than cancer (all non-malignant respiratory diseases and chronic obstructive bronchitis and emphysema).

Analysis of data from survivors of A bomb

We analysed mortality data from the A bomb survivors for solid cancer to 1997⁶ and leukaemia to 1990⁷ using similar methods to provide risk estimates for comparison.

Results

The main study population comprised 407 391 workers (table 1). Most workers in the study were men (90%), and men received 98% of the collective dose. The distribution of recorded doses was skewed (fig 1). The average cumulative dose was 19.4 mSv. Ninety per cent




Fig 1 Distribution of cumulative radiation doses among workers included in the analyses

of workers received cumulative doses < 50 mSv and less than 0.1% received cumulative doses > 500 mSv.

For all cancers excluding leukaemia, the excess relative risk was significantly different from zero (table 2). This estimate corresponds to a relative risk of 1.10 for a radiation dose of 100 mSv. For solid cancers, the excess relative risk was higher than but statistically compatible with the estimate for A bomb survivors. The excess relative risk for leukaemia excluding chronic lymphocytic leukaemia corresponds to a relative risk of 1.19 for a dose of 100 mSv. This estimate is between the linear and linear quadratic extrapolations from data on A bomb survivors.

Indirect analyses of the possible confounding effect of smoking yielded excess relative risks that ranged between 0.59 per Sv (-0.29 to 1.70) for all cancers excluding leukaemia and lung and pleural cancer, and 0.91 per Sv (-0.11 to 2.21) for smoking related cancers (see bmj.com).

The increased risk for smoking related cancers was mainly due to an increased risk of lung cancer (1.86 per

Table 1 Cohorts included in the 15 country study

	No of facilities	First year of operations	Follow-up period	No of workers	Person years	Deaths			Average individual cumulative dose (mSv)	
						All causes	All cancers excluding leukaemia	Leukaemia excluding CLL		Collective cumulative dose (Sv)
Australia	1	1959	1972-98	877	12 110	56	17	0	5.4	6.1
Belgium	5	1953	1969-94	5 037	77 246	322	87	3	134.2	26.6
Canada	4	1944	1956-94	38 736	473 880	1 204	400	11	754.3	19.5
Finland	3	1960	1971-97	6 782	90 517	317	33	0	53.2	7.8
France CEA-COGEMA	9	1946	1968-94	14 796	224 370	645	218	7	55.6	3.8
France EDF	22	1956	1968-94	21 510	241 391	371	113	4	340.2	15.8
Hungary	1	1982	1985-98	3 322	40 557	104	39	1	17.0	5.1
Japan	33*	1957	1986-92	83 740	385 521	1 091	413	19	1526.7	18.2
Korea (south)	4	1977	1992-97	7 892	36 227	58	21	0	122.3	15.5
Lithuania	1	1984	1984-2000	4 429	38 458	102	24	1	180.2	40.7
Slovak Republic	1	1973	1973-93	1 590	15 997	35	10	0	29.9	18.8
Spain	10	1968	1970-96	3 633	46 358	68	25	0	92.7	25.5
Sweden	6	1954	1954-96	16 347	220 501	669	190	4	291.8	17.9
Switzerland	4	1957	1969-95	1 785	22 051	66	24	0	111.2	62.3
UK	32	1946	1955-92	87 322	1 370 101	7 983	2201	54	1810.1	20.7
US Hanford	1	1944	1944-86	29 332	678 833	5 564	1279	35	695.4	23.7
US INEL	1	1949	1960-96	25 570	505 236	3 491	886	26	254.6	10.0
US NPP	15	1960	1979-97	49 346	576 682	983	314	19	1336.0	27.1
US ORNL	1	1943	1943-84	5 345	136 673	1 029	225	12	81.1	15.2
Total	154	—	—	407 391	5 192 710	24 158	6519	196	7892.0	19.4

CEA-COGEMA=Commissariat à l’Energie Atomique-Compagnie Générale des Matières Nucléaires; EDF=Electricité de France; NPP=nuclear power plants; INEL=Idaho National Engineering Laboratory; ORNL=Oak Ridge National Laboratory; CLL=chronic lymphocytic leukaemia.

*No information available to allow separation of different facilities.

Sv, 0.26 to 4.01). Other smoking related cancers showed little evidence of an increased risk (0.21 per Sv, <0 to 2.01). Risk estimates for mortality from non-malignant respiratory diseases and from chronic obstructive bronchitis and emphysema were raised but not significantly different from zero (excess relative risk per Sv 1.16, -0.53 to 3.84, and 2.12, -0.57 to 7.46, respectively).

Discussion

Results from our study suggest that in workers in the nuclear industry an excess risk of cancer exists, albeit small, even at low doses and dose rates. The 15 country study allowed the compilation of the largest body of direct evidence to date concerning the effects of low dose chronic exposure to ionising radiation.

All cancer excluding leukaemia

We found a significantly increased risk for all cancers (excluding leukaemia). The central risk estimate was higher than the linear extrapolation from the A bomb survivors. It is unlikely that this could be due to ascertainment bias, as the excess relative risk for all non-cancer mortality was weakly positive (0.20, -0.26 to 0.72).

Information was not available to adjust directly for possible confounding by variables such as smoking, diet, and occupational exposures. Some of these factors—particularly smoking and diet—are strongly related to socioeconomic status and adjustment for this will have partially controlled for their effects. Some studies have found an association between radiation dose and smoking,^{8,9} while others have not.¹⁰⁻¹²

The central risk estimate for cancers unrelated to smoking was higher than that for smoking related cancers other than lung cancer, indicating that confounding by smoking is unlikely to explain all of the relation found between all cancer risk and radiation dose. On the other hand, the non-significantly increased risks for mortality from non-malignant smoking related diseases indicate a possible effect of smoking. The risk estimates for mortality from cancers related and unrelated to smoking, however, are consistently two to three times higher than, but statistically compatible with the risk estimate for solid cancers from the A bomb analyses. Taken together, these findings indicate that a confounding effect by smoking may be partly, but not




Fig 2 Excess relative risks per Sv for all cancer excluding leukaemia in cohorts with more than 100 deaths (NPP=nuclear power plants, ORNL=Oak Ridge National Laboratory)

entirely, responsible for the estimated increased risk for mortality from all cancers other than leukaemia.

Formal tests for heterogeneity provided no evidence for differences in risk between countries, cohorts, or groups of facilities ($P > 0.20$). Figure 2 shows the excess relative risk per Sv in the larger cohorts (> 100 cancer deaths). Analyses excluding one cohort or country at a time produced excess relative risks per Sv ranging from 0.58 (excluding Canada) to 1.25 (excluding the UK), all consistently higher than but compatible with the estimate from A bomb analyses. Only when we excluded Canada was the excess relative risk no longer significantly different from zero (0.58, -0.22 to 1.55).

Sensitivity analyses of different lag periods showed that both the risk estimates and their uncertainties increased with increasing lag. The excess relative risk per Sv ranges from 0.76 (0.07 to 1.59) with a lag of five years to 1.68 (0.22 to 3.48) with a lag of 20 years. The estimates are all statistically compatible with the linear extrapolation from the A bomb survivors.

Leukaemia excluding chronic lymphocytic leukaemia

Although our estimate of risk of leukaemia is not significantly different from zero, it is similar to estimates from previous large scale studies of nuclear workers.^{13,14} Furthermore, it is intermediate between estimates obtained by fitting a linear and a linear quadratic dose-response model to data on men exposed to the A bomb at age 20-60. The excess relative risk per

Table 2 Estimates of excess relative risk per Sv (95% confidence interval) for all cancers excluding leukaemia, solid cancers, and leukaemia excluding chronic lymphocytic leukaemia, for nuclear workers and survivors of A bomb in Japan*

	15 country study		Atomic bomb survivors (men exposed at age 20-60)	
	No of cancers	Risk	No of cancers	Risk†
All cancers excluding leukaemia	5024	0.97 (0.14 to 1.97)		
Solid cancers	4770	0.87 (0.03 to 1.88)	3246	0.32‡ (0.01 to 0.50)
Leukaemia excluding CLL:				
Linear model	196	1.93 (<0§ to 8.47)	83	3.15¶ (1.58 to 5.67)
Linear quadratic model				1.54** (-1.14 to 5.33)

CLL=chronic lymphocytic leukaemia.

*Colon dose used for all cancers and solid cancer analyses, bone marrow dose for leukaemia.

†Note that because analyses were restricted to men aged 20-60 at exposure the confidence intervals are much wider than those presented by other investigators,^{13,14} based on the full cohort.

‡Analyses carried out at IARC with excess relative risk model that allows for age at exposure modification, adjusted for attained age, calendar period, and city. Estimate for men exposed at age 35.

§Estimate on boundary of parameter space.

¶Analyses carried out at IARC with constant excess relative risk model, adjusted for attained age, calendar period, and city.

**Analyses carried out at IARC—linear term of linear quadratic model—preferred model for describing leukaemia mortality in analyses of data on A bomb survivors.¹⁴

What is already known on this topic

Current radiation protection standards are based mainly on data from the survivors of the atomic bomb in Japan

The estimation of risks after low dose protracted or fractionated exposures to ionising radiation is controversial

What this study adds

A small excess risk of cancer exists, even at the low doses typically received by nuclear industry workers in this study

Sv shows only a small increase with increasing lag periods, from 1.93 (<0 to 8.47) with a lag of two years to 2.53 (<0 to 10.45) with a lag of 10 years.

Implications for radiation protection

Current recommendations are to limit occupational doses to 100 mSv over five years (not to exceed 50 mSv in any one year) and doses to the public to 1 mSv per year.¹⁵ Our estimates suggest that a cumulative exposure of 100 mSv would lead to a 9.7% (1.4 to 19.7%) increased mortality from all cancers excluding leukaemia and a 5.9% (-2.9 to 17.0%) increased mortality from all cancers excluding leukaemia, lung, and pleura compared with background rates. The corresponding figure is 19% (<0 to 84.7%) for mortality from leukaemia excluding chronic lymphocytic leukaemia. Less than 5% of workers in this study received cumulative doses of the order of 100 mSv over their entire career. We estimate that 1-2% of deaths from cancer (including leukaemia) among workers in this cohort may be attributable to radiation.

Conclusions

We have provided radiation risk estimates from the largest study of nuclear industry workers conducted so far. These estimates are higher than but statistically compatible with the current bases for radiation protection standards. The confidence intervals range from values lower than those derived by linear extrapolation from data from A bomb survivors up to values that exceed this extrapolation by a factor of six for cancers other than leukaemia and nearly three for leukaemia. These results suggest that an excess risk of cancer exists, albeit small, even at the low doses and dose rates typically received by nuclear workers in this study.

We are grateful to Richard Doll, Jacques Estève, and Bruce Armstrong who helped to start this study; to the late Len Salmon who inspired the study of errors in doses; to past members of the dosimetry and epidemiology subcommittees (William Murray, Robert Rinsky) and of the international study group (the late E Gubéran, Y Hosoda, T Iwasaki, G Kendall, M Murata, T Rytömaa); to everyone in the participating countries who worked in the collection and validation of the data used in the study; and to the representatives and staff of the nuclear facilities included in the study for their open collaboration. This report uses data obtained from the Radiation Effects Research Foundation (RERF), Hiroshima and Nagasaki, Japan. RERF is a private, non-profit foundation funded by the Japanese Ministry of Health, Labour and Welfare and the US Department of Energy through the National Academy of Sciences. The conclusions in this report are those of the authors and do not necessarily reflect the scientific judgment of RERF or its funding agencies. The other members of the international study group

were: F Bermann, G Cowper, J Fix, C Hacker, B Heinmiller, M Marshall, I Thierry-Chef, D Utterback, Y-O Ahn, E Amoros, P Ashmore, A Auvinen, J-M Bae, J Bernar Solano, A Biau, E Combalot, P Deboodt, A Diez Sacristan, M Eklof, H Engels, G Engholm, G Gulis, R Habib, K Holan, H Hyvonen, A Kerekes, J Kurtinaitis, H Malke, M Martuzzi, A Mastauskas, A Monnet, M Moser, M S Pearce, D B Richardson, F Rodriguez-Artalejo, A Rogel, H Tardy, M Telle-Lamberton, I Turai, M Usel, K Veress. Contributors: See bmj.com

Funding: European Union (contracts F13P-CT930066, F14P-CT96-0062, FIGH-CT1999-20001); US Centers for Disease Control (Co-operative agreement U50/CCU011778); Canadian Nuclear Safety Commission; Japanese Institute for Radiation Epidemiology, Australian Nuclear Science and Technology Organisation; Nuclear Research Center (SCKCEN), Belgium; Health Canada and Statistics Canada; La Ligue Nationale contre le Cancer, France; La Compagnie Générale des Matière Nucléaire, France; Electricité de France; Ministry of Education, Culture, Sports, Science and Technology in Japan; Ministry of Science and Technology (MOST) in Korea; Spanish Nuclear Safety Council; US Department of Energy. These sponsors had no role in study design, data collection, analysis, or interpretation. IFC received funding from the Association pour la Recherche contre le Cancer (ARC, France); MSP and DBR were the recipients, respectively, of an IARC postdoctoral fellowship and of an ORAU fellowship during their stay at IARC.

Competing interests: C Hacker, B Heinmiller, H Hyvonen, M Marshall, A Rogel, J Bernar Solano, M Eklof, and K Holan are (or have been in the past five years) employees of the nuclear industry or have links to the nuclear industry in their country. They were appointed to the international study group and/or the dosimetry subcommittee as experts because of their critical knowledge and experience in historical radiation protection practices and dosimetry. None of them had influence on decisions concerning analysis of the results.

Ethical approval: The study was approved by the IARC ethical review committee and by the relevant ethical committees of the participating countries.

- 1 BEIR V. Committee on the Biological Effects of Ionizing Radiation. *The effects on populations of exposure to low levels of ionizing radiation*. Washington, DC: National Academy of Sciences, 1990.
- 2 Cardis E, Esteve J. *International collaborative study of cancer risk among nuclear industry workers. I—Report of the feasibility study. II—Protocol report 92/001*. Lyon: International Agency for Research on Cancer, 1992.
- 3 Cardis E, Martuzzi M, Amoros E. *International collaborative study of cancer risk among nuclear industry workers. III—Procedures document, rev 1*. Lyon: International Agency for Research on Cancer, 1997. (97/002.)
- 4 Fix JJ, Salmon L, Cowper G, Cardis E. A retrospective evaluation of the dosimetry employed in an international combined epidemiological study. *Radiat Protect Dosimetry* 1997;74:39-53.
- 5 United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR). *Sources and effects of ionizing radiation*. New York: United Nations, 2000.
- 6 Preston DL, Shimizu Y, Pierce DA, Suyama A, Mabuchi K. Studies of mortality of atomic bomb survivors. Report 13: Solid cancer and noncancer disease mortality: 1950-1997. *Radiat Res* 2003;160:381-407.
- 7 Pierce DA, Shimizu Y, Preston DL, Vaeth M, Mabuchi K. Studies of the mortality of atomic bomb survivors. Report 12, Part I. Cancer: 1950-1990. *Radiat Res* 1996;146:1-27.
- 8 Gribbin MA, Weeks JL, Howe GR. Cancer mortality (1956-1985) amongst male employees of Atomic Energy of Canada Limited with respect to occupational exposure to external low linear energy transfer ionizing radiation. *Radiat Res* 1993;133:375-80.
- 9 Murata M, Miyake T, Inoue Y, Ohshima S, Kudo S, Yoshimura T, et al. Life-style and other characteristics of radiation workers at nuclear facilities in Japan: base-line data of a questionnaire survey. *J Epidemiol* 2002;12:310-9.
- 10 Peterson G, Gilbert ES, Buchanan JA, Stevens RG. A case-cohort study of lung cancer ionizing radiation, and tobacco smoking among males at the Hanford sites. *Health Phys* 1990;58:3-11.
- 11 Carpenter L, Fraser P, Booth M, Higgins C, Beral V. Smoking habits and radiation exposure. *J Radiol Protection* 1989;9:286-7.
- 12 Auvinen A, Pukkala E, Hyvönen H, Hakama M, Rytömaa T. Cancer incidence among Finnish nuclear reactor workers. *J Occup Environ Med* 2002;44:634-8.
- 13 IARC Study Group on Cancer Risk among Nuclear Industry Workers. Direct estimates of cancer mortality due to low doses of ionising radiation: an international study. *Lancet* 1994;344:1039-43.
- 14 Muirhead CR, Goodill AA, Haylock RGE, Vokes J, Little MP, Jackson DA, et al. Occupational radiation exposure and mortality: second analysis of the national registry for radiation workers. *J Radiol Prot* 1999;19:3-26.
- 15 International Commission on Radiological Protection ICRP. *Recommendations of the international commission on radiological protection*. Oxford: Pergamon Press, 1991. (ICRP publication 60.)

(Accepted 23 May 2005)

doi 10.1136/bmj.38499.599861.E0