

- 3 Lang S, Lawrence CJ, Orme RL. Cup feeding: an alternative method of infant feeding. *Arch Dis Child* 1994;71:365-9.
- 4 Stine MJ. Breastfeeding the premature newborn: a protocol without bottles. *J Hum Lact* 1990;6:167-70.
- 5 Kliethermes PA, Cross ML, Lanese MG, Johnson KM, Simon SD. Transitioning preterm infants with nasogastric tube supplementation: increased likelihood of breastfeeding. *J Obstet Gynecol Neonatal Nurs* 1999;28:264-73.
- 6 Rocha NM, Martinez FE, Jorge SM. Cup or bottle for preterm infants: effects on oxygen saturation, weight gain, and breastfeeding. *J Hum Lact* 2002;18:132-8.
- 7 Pinelli J, Symington A. Non-nutritive sucking for promoting physiologic stability and nutrition in preterm infants. *Cochrane Database Syst Rev* 2004;(4): CD001071.
- 8 Lang S. Cup feeding: an alternative method. *Midwives Chronicle* 1994;107:171-6.
- 9 World Health Organization. *Indicators for assessing breast-feeding practices*. Geneva: WHO, 1991.
- 10 Hill PD, Hanson KS, Mefford AL. Mothers of low birthweight infants: breastfeeding patterns and problems. *J Hum Lact* 1994;10:169-76.
- 11 Labbok M, Krasovec K. Toward consistency in breastfeeding definitions. *Stud Fam Plann* 1990;21:226-30.
- 12 Altman DG. Confidence intervals for the number needed to treat. *BMJ* 1998;317:1309-12.
- 13 Kaufman KJ, Hall LA. Influences of the social network on choice and duration of breast-feeding in mothers of preterm infants. *Res Nurs Health* 1989;12:149-59.
- 14 Yip E, Lee J, Sheehy Y. Breast-feeding in neonatal intensive care. *J Paediatr Child Health* 1996;32:296-8.
- 15 Boo NY, Goh ES. Predictors of breastfeeding in very low birthweight infants at the time of discharge from hospital. *J Trop Pediatr* 1999;45:195-201.
- 16 Nyqvist KH, Ewald U. Infant and maternal factors in the development of breastfeeding behaviour and breastfeeding outcome in preterm infants. *Acta Paediatr* 1999;88:1194-203.
- 17 Killersreiter B, Grimmer I, Buhner C, Dudenhausen JW, Obladen M. Early cessation of breast milk feeding in very low birthweight infants. *Early Hum Dev* 2001;60:193-205.
- 18 Furman L, Minich N, Hack M. Correlates of lactation in mothers of very low birth weight infants. *Pediatrics* 2002;109:e57. www.pediatrics.org/cgi/content/full/109/4/e57 (accessed Sep 2002).
- 19 Prescott R, Counsell C, Gillespie W, Grant A, Russell I, Kiauka S, et al. Factors that limit the quality, number and progress of randomised controlled trials. *Health Technol Assess* 1999;3:1-143.
- 20 Lang S. *Breastfeeding special care babies*. London: Bailliere Tindall, 1997. (Accepted 6 May 2004)

doi 10.1136/bmj.38131.675914.55

Infertility among male UK veterans of the 1990-1 Gulf war: reproductive cohort study

Noreen Maconochie, Pat Doyle, Claire Carson

Department of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, University of London, London WC1E 7HT

Noreen Maconochie senior lecturer in epidemiology and medical statistics
Pat Doyle reader in epidemiology
Claire Carson research assistant

Correspondence to: N Maconochie noreen.maconochie@lshtm.ac.uk

BMJ 2004;329:196-200

Abstract

Objectives To examine the hypothesis that, theoretically at least, exposure to toxicants of the type present in the Gulf war could affect spermatogenesis, which might be observed as increased levels of infertility.

Design Retrospective reproductive cohort analysis.

Setting Male UK Gulf war veterans and matched comparison group of non-deployed servicemen, surveyed by postal questionnaire.

Participants 42 818 completed questionnaires were returned, representing response rates of 53% for Gulf veterans and 42% for non-Gulf veterans; 10 465 Gulf veterans and 7376 non-Gulf veterans reported fathering or trying to father pregnancies after the Gulf war.

Main outcome measures Failure to achieve conceptions (type I infertility) or live births (type II infertility) after the Gulf war, having tried for at least a year and consulted a doctor; time to conception among pregnancies fathered by men not reporting fertility problems.

Results Risk of reported infertility was higher among Gulf war veterans than among non-Gulf veterans (odds ratio for type I infertility 1.41, 95% confidence interval 1.05 to 1.89; type II 1.50, 1.18 to 1.89). This small effect was constant over time since the war and was observed whether or not the men had fathered pregnancies before the war. Results were similar when analyses were restricted to clinically confirmed diagnoses. Pregnancies fathered by Gulf veterans not reporting fertility problems also took longer to conceive (odds ratio for >1 year 1.18, 1.04 to 1.34).

Conclusions We found some evidence of an association between Gulf war service and reported infertility. Pregnancies fathered by Gulf veterans with

no fertility problems also reportedly took longer to conceive.

Introduction

In late 1990 and early 1991 around 53 000 UK armed service personnel were deployed to the Gulf war. Compared with the many reports on adult health after service in the Gulf, relatively few epidemiological studies have been conducted on reproductive outcomes.¹⁻⁹ Only two small studies specifically examined infertility, with inconsistent results.⁵⁻⁸

We now report findings relating to infertility from the only epidemiological survey of reproductive outcomes among UK Gulf veterans. Analyses of fetal death and congenital malformation have been reported elsewhere.⁹

Methods

Main survey

Detailed information about the study is given elsewhere.^{9 10} This was a retrospective cohort study of the reproductive health of all UK armed forces personnel deployed to the Gulf region between August 1990 and June 1991 (51 581 men, 1230 women) and a comparison group (matched on service, sex, age, fitness to be deployed, serving status, and rank) who were in service at that time (January 1991) but were not deployed (51 688 men, 1236 women). A postal questionnaire, sent from August 1998 (with reminders until early 2001), requested detailed information on reproductive history, including questions on infertility



This is the abridged version of an article that was posted on bmj.com on 14 July 2004: <http://bmj.com/cgi/doi/10.1136/bmj.38163.620972.AE>

problems and their diagnosis and treatment. After adjustment for undelivered mail, response rates among men were 53% (24 379) for Gulf veterans and 42% (18 439) for non-Gulf veterans. An analysis of non-responders found no evidence of bias with respect to reproduction. Of these men, 18 924 (44%) had fathered or tried unsuccessfully to father children since the Gulf war, 5711 (30%) of whom also reported pregnancies before the war.¹⁰

Investigation of infertility

The figure summarises the construction of the dataset. We restricted the analyses to men who had tried to have children since the Gulf war, regardless of whether they had conceived pregnancies before that. We further restricted the analyses to men whose first post-Gulf conception or pregnancy attempt was before August 1997.

We defined participants as having fertility problems if they had tried for a pregnancy unsuccessfully for more than a year since the Gulf war and had consulted a doctor before 1 August 1997. We defined infertility as having fertility problems and either never achieving a recognised pregnancy (type I infertility) or never achieving a pregnancy ending in a live birth (type II infertility). We included participants as non-infertility cases if they had fathered at least one pregnancy after the Gulf war before the reported unsuccessful pregnancy attempt.

Coding and clinical verification


Where relevant permission was available, we attempted to verify reported fertility problems by contacting both partners' general practitioner or relevant clinician. We requested diagnostic details, with a copy of semen analysis results if available. We received clinical information for similar proportions in the two groups (245/732 (33%) Gulf veterans, 117/370 (32%) non-Gulf veterans; $P=0.53$ for difference). We coded diagnoses according to World Health Organization definitions¹¹; we based our coding on information received from the clinician where available and otherwise on the participant's description only. The three main indicators of semen quality were oligospermia (low sperm count) (at least one result of <20 million sperm/ml), asthenospermia (poor motility) (at least one result of $<50\%$ motility), and teratospermia (abnormal morphology) (at least one result of $>70\%$ abnormal forms).

Time to conception

We asked participants whether each pregnancy was planned and, if yes, how long it had taken to conceive, grouped as <3 , 3-6, 7-12, and >12 months. We included in these analyses all pregnancies fathered after the Gulf war (and before 1 August 1997) by participants not reporting fertility problems.

Statistical methods

We used Stata statistical software for all analyses. We used logistic regression to estimate the effect of Gulf war service (or certain specific agents) on risk of infertility,¹² with non-Gulf veterans as baseline. We adjusted odds ratios for age of both participant and female partner at first consultation for infertility or post-Gulf conception, year of first consultation or post-Gulf conception, having fathered pregnancies before the war, and participant's service and rank at the time of the



Creation of dataset for analysis of infertility among male Gulf veterans and non-Gulf veterans. *After Gulf war deployment (Gulf veterans) or 1 January 1991 (non-Gulf veterans) and before start of main study (1 August 1998). †At least one pregnancy fathered after Gulf war and before 1 August 1997. ‡Congenital malformation of the urogenital system, sterilisation as a result of cancer treatment or injury, and unsuccessful reversal of (elective) sterilisation do not fulfil case definition for infertility

Gulf war. We examined smoking at time of first consultation or conception and number of weekly units of alcohol consumed at survey, but as neither acted as confounders we excluded them from the final models.

For time to conception the unit of analysis was a pregnancy; we estimated the effect of Gulf war service on risk of infertility by using logistic regression, with non-Gulf veterans as baseline. We adjusted odds ratios in all analyses for pregnancy order, maternal age, year of conception, and service and rank at the time of the Gulf war. Smoking and alcohol again did not confound the effect of interest and were excluded from final models.

Results

Study population

In total, 10 465 Gulf veterans and 7376 non-Gulf veterans who had conceived or attempted to conceive a child after the Gulf war satisfied the eligibility criteria.

Infertility

Seven hundred and thirty two (7%) Gulf veterans and 370 (5%) non-Gulf veterans had consulted a doctor for fertility problems arising since the Gulf war (odds ratio 1.38, 95% confidence interval 1.20 to 1.60). More than 60% (n = 721) of these men had succeeded in fathering one or more pregnancy, and more than 50% (n = 580) had fathered one or more live birth by the time of survey (table 1).

Prevalence of type I infertility was higher in Gulf veterans (2.5% (n = 259) in Gulf veterans and 1.7% (n = 122) in non-Gulf veterans; odds ratio 1.41, 95% confidence interval 1.05 to 1.89). The effect was

Table 1 Characteristics of men included in the analysis. Values are numbers (percentages) unless stated otherwise

	Gulf veterans (n=10 465)	Non-Gulf veterans (n=7376)
Age at survey (years)		
<30	2642 (25.3)	1491 (20.2)
30-34	4403 (42.1)	3134 (42.5)
35-39	2545 (24.3)	2046 (27.7)
≥40	875 (8.4)	705 (9.6)
Mean (SD)	33.4 (4.51)	33.9 (4.53)
Year of first attempt or conception after Gulf war*		
1991†	2296 (21.9)	1679 (22.8)
1992-3	3717 (35.5)	2691 (36.5)
1994-5	2726 (26.1)	1848 (25.1)
1996-7‡	1726 (16.5)	1158 (15.7)
Age at first attempt or conception after Gulf war* (years)		
<25	2807 (26.8)	1790 (24.3)
25-29	4535 (43.3)	3188 (43.2)
30-34	2363 (22.6)	1803 (24.4)
≥35	760 (7.3)	595 (8.1)
Mean (SD)	28.0 (4.50)	28.4 (4.55)
Age of wife or female partner§ at first attempt or conception after Gulf war (years)		
<25	3279 (31.8)	2116 (29.0)
25-29	4226 (41.0)	3026 (41.5)
30-34	2221 (21.6)	1708 (23.4)
≥35	581 (5.6)	449 (6.2)
Mean (SD)	27.0 (4.54)	27.2 (4.49)
Pregnancies before Gulf war		
All	2829 (27.0)	2492 (33.8)
Fertility problems after Gulf war¶		
All:	732 (7.0)	370 (5.0)
Reported pregnancies before Gulf war	117 (16.0)	72 (19.5)
Achieved recognised pregnancies	473 (64.6)	248 (67.0)
Achieved live births	376 (51.4)	204 (55.1)
No conceptions:	259 (2.5)	122 (1.7)
With pregnancies before Gulf war	45 (17.4)	27 (22.1)
No live births:	356 (3.4)	166 (2.3)
With pregnancies before Gulf war	61 (17.1)	35 (21.1)

*First unsuccessful attempt after Gulf war for those reporting fertility problems; first post-Gulf war conception for those reporting no fertility problems. Reproductive events assigned pre-Gulf or post-Gulf war status by date of first deployment (Gulf veterans) or 1 January 1991 (non-Gulf veterans and Gulf veterans where date of first deployment not known).

†Includes 135 Gulf veterans who conceived pregnancies or first attempted unsuccessfully to have children in 1990, after deployment to the Gulf.

‡Attempts and conceptions included only up to 31 July 1997 (see methods).

§Excluding 158 Gulf veterans and 77 non-Gulf veterans with missing data.

¶Tried for a pregnancy after Gulf war without success for more than a year and consulted a doctor before 1 August 1997 about failure to conceive. With or without conceptions before Gulf war, but no conceptions after Gulf war before unsuccessful attempt. Excludes 14 men reporting fertility problems due to unsuccessful reversal of (elective) sterilisation, congenital urogenital anomaly, injuries (for example, paralysis), or cancer treatment.

stronger for type II infertility (3.4% (n = 356) and 2.3% (n = 166); odds ratio 1.50, 1.18 to 1.89) (table 2). For most men, this was primary infertility; only 45 (17%) Gulf veterans and 27 (22%) non-Gulf veterans with type I infertility had fathered one or more pregnancies before the war (table 1). These effects did not vary with time since the war (P values for interaction 1.00 for type I infertility and 0.56 for type II infertility) or with whether the infertility was primary (no previous conceptions) or secondary (P values for interaction 0.83 for type I and 0.82 for type II).

We found only weak evidence of an association between Gulf war service and a diagnosis of any male factor infertility (odds ratio for type I infertility 1.16, 0.74 to 1.82), though the effect was slightly stronger for type II infertility (odds ratio 1.45, 0.98 to 2.14) (table 2). We found evidence for a more specific effect on risk of teratospermia (odds ratios 2.02, 0.79 to 5.14 for type I; 2.55, 1.03 to 6.30 for type II), and an association was also suggested between Gulf war service and risk of oli-

goasthenoteratospermia. The numbers of these cases were, however, extremely small. Results were similar when restricted to clinically confirmed diagnoses.

We examined risk of infertility in relation to four self reported Gulf war exposures (vaccination against anthrax or plague, nerve agent pretreatment sets, depleted uranium, and pesticides). A high proportion (up to 37.3%) of men could not recall or did not know their exposure. Overall, the analyses revealed little or no evidence of an increased risk of infertility in relation to any specific exposure (data tabulated on bmj.com).

Time to pregnancy

The 9733 Gulf veterans and 7006 non-Gulf veterans who did not have fertility problems had fathered 15 593 and 11 023 pregnancies respectively since the Gulf war (table 3). Pregnancies fathered by Gulf veterans were more often reported as unplanned (adjusted odds ratio 1.12, 1.05 to 1.19). Among planned pregnancies, those fathered by Gulf veterans took longer to conceive; 9.1% (n = 845) took more than a year compared with 7.8% (528) of those fathered by non-Gulf veterans (adjusted odds ratio 1.18, 1.04 to 1.34).

Discussion

Principal findings

We found a small increased risk of infertility associated with service in the 1990-1 Gulf war, which was strengthened when we extended the definition to include men reporting fertility problems who had fathered only pregnancies ending in fetal death. We also found that pregnancies fathered by Gulf veterans who did not report fertility problems took longer to conceive, although again the effect was not large

We found a weak and non-statistically significant association between Gulf war service and a diagnosis of male factor infertility overall. We also found some suggestion that teratospermia and oligoasthenoteratospermia were increased in Gulf war veterans. These results should, however, be treated with extreme caution, given the small numbers and the fact that we were able to clinically validate only around 40% of cases of infertility.

Consistency with other studies

Our results are consistent with the findings of a study of male Australian veterans, in which Gulf veterans had a 40% increased risk of having fertility problems.⁸ However, the results conflict with a study of male Danish veterans, the only other study on this topic, which found no evidence of an effect of Gulf war service on markers of male fertility.⁵

Limitations

As discussed in previous reports,^{9 10} our study does have a fairly low response rate, raising the possibility of selective participation according to adverse reproductive outcome. However, comparison of reproductive patterns among responders to the main study with those in a study of intensively traced non-responders provided no strong evidence of selection bias related to reproduction.¹⁰ The proportions of Gulf veterans and non-Gulf veterans who had consulted a doctor for infertility were almost identical among responders and non-responders.¹⁰

What is already known on this topic

Relatively few epidemiological studies have examined reproductive outcomes in veterans of the 1990-1 Gulf war, and only two of these have specifically examined infertility

One study found no effect of Gulf war service on reproductive hormones or reported fertility; the second found that male Gulf veterans were more likely than the comparison group to report difficulties with fertility

In both of these studies expected numbers were small and power consequently low

What this study adds

Male UK veterans of the 1990-1 Gulf war had a small increased risk of reported infertility

Pregnancies fathered by veterans of the 1990-1 Gulf war who did not report fertility problems were reported to take longer to conceive

These results should be interpreted with caution, and we cannot at this stage conclude that the associations are causal

Differential recall of infertility by the Gulf veterans or the comparison group is also a possibility. It could be argued that Gulf veterans had more incentive to report infertility if they perceived that it might be related to their Gulf war service. Unfortunately, few reliable British population based estimates of infertility are available for comparison.

Interpretation

Despite our data's limitations, the reported observations seem robust. The increases in risk did not decline with time since the Gulf war, nor were they different among men who had fathered children before the Gulf war. The constant effect over time argues in favour of either paternal germ cell mutation or other damage to spermatogenic stem cells or the testicular cells necessary for supporting spermatogenesis. In previous work we found no increase in genetic syndromes and chromosomal anomalies in the offspring of Gulf veterans, but we did observe an increase in reported miscarriage.⁹ Furthermore, the suggestion in our data that the effect of Gulf war service might be strongest for teratospermia, and perhaps oligoasthenoteratospermia, is consistent with findings of apoptosis in testicular germ cells, Sertoli cells, and Leydig cells in rats subjected to combined exposure to pyridostigmine bromide, the insect repellent DEET, and the insecticide permethrin (all of which are reported to have been present during the Gulf war).¹³ Some of our analyses involved extremely small numbers, however, and no firm conclusions can be drawn from them.

Gulf war veterans are known to report more illnesses than non-veterans.¹⁴ To what extent these illnesses have a direct effect on infertility and time to conception is not known, but Gulf related illness may have played a part in our findings.

Table 2 Reported prevalence of infertility* after Gulf war. Values are numbers (percentages) unless stated otherwise

Outcome	Gulf veterans (n=10 465)	Non-Gulf veterans (n=7376)	Adjusted odds ratio† (95% CI)
Reported fertility problems after Gulf war‡:	732 (7.0)	370 (5.0)	1.38 (1.20 to 1.60)
No conceptions (type I infertility)	259 (2.5)	122 (1.7)	1.41 (1.05 to 1.89)
No live births (type II infertility)	356 (3.4)	166 (2.3)	1.50 (1.18 to 1.89)
Type I infertility—no conceptions			
Any male infertility factor§:	97 (0.9)	47 (0.6)	1.16 (0.74 to 1.82)
Azoospermia¶	10 (0.1)	5 (0.1)	0.95 (0.21 to 4.26)
Oligospermia¶	63 (0.6)	27 (0.4)	1.38 (0.75 to 2.53)
Asthenospermia¶	39 (0.4)	19 (0.3)	0.97 (0.51 to 1.85)
Teratospermia¶	21 (0.2)	6 (0.1)	2.02 (0.79 to 5.14)
Oligoasthenoteratospermia¶	8 (0.1)	2 (0.03)	2.17 (0.43 to 10.89)
Type II infertility—no live births			
Any male infertility factor§:	125 (1.2)	54 (0.7)	1.45 (0.98 to 2.14)
Azoospermia¶	10 (0.1)	5 (0.1)	0.95 (0.21 to 4.26)
Oligospermia¶	80 (0.8)	32 (0.4)	1.58 (0.94 to 2.65)
Asthenospermia¶	51 (0.5)	23 (0.3)	1.19 (0.68 to 2.08)
Teratospermia¶	26 (0.3)	6 (0.1)	2.55 (1.03 to 6.30)
Oligoasthenoteratospermia¶	9 (0.1)	2 (0.03)	2.47 (0.51 to 12.00)

*Infertility defined as trying for a pregnancy after Gulf war without success for more than a year and consulting a doctor before 1 August 1997 about failure to conceive (no conceptions after Gulf war before unsuccessful attempt), with no subsequent conceptions achieved (type I) or no live births achieved (type II). Excludes 14 men reporting fertility problems due to unsuccessful reversal of (elective) sterilisation, congenital urogenital anomaly, injuries (for example, paralysis), or cancer treatment.

†Adjusted for age of both participant and female partner at first consultation for infertility or first post-Gulf conception (not infertile), year of first consultation or post-Gulf conception, previous pregnancies (before Gulf war), and service and rank at time of Gulf war.

‡Whether or not conceptions subsequently achieved (that is, whether or not infertile, as defined in this paper).

§Excludes 136 Gulf veterans and 61 non-Gulf veterans for whom diagnosis not known.

¶Participants may appear in more than one category if diagnosed as having more than one infertility factor.

Table 3 Pregnancies fathered by Gulf war veterans (n=9733) and non-Gulf war veterans (n=7006) not reporting fertility problems. Values are numbers (percentages) unless stated otherwise

	Pregnancies fathered by		Adjusted* odds ratio (95% CI)
	Gulf veterans (n=15 593)	Non-Gulf veterans (n=11 023)	
Planned†:			
Yes	9968 (68.0)	7408 (71.5)	1.12‡ (1.05 to 1.19)
No	4700 (32.0)	2951 (28.5)	
Planned pregnancies only:			
Time to conception§:			
<3 months	5647 (61.0)	4410 (65.5)	1.18¶ (1.04 to 1.34)
3-6 months	1907 (20.6)	1224 (18.2)	
7-12 months	858 (9.3)	572 (8.5)	
>12 months	845 (9.1)	528 (7.8)	

*Adjusted for pregnancy order, maternal age, year of conception, and service and rank at time of Gulf war. Standard errors adjusted for repeated pregnancies fathered by same participant.

†Excluding 925 pregnancies reported by Gulf veterans and 664 by non-Gulf veterans with missing data.

‡Unplanned pregnancy as opposed to planned pregnancy in Gulf veterans relative to non-Gulf veterans.

§Excluding 711 planned pregnancies reported by Gulf veterans and 674 by non-Gulf veterans with missing data.

¶More than 12 months to conception as opposed to less than 12 months to conception in Gulf veterans relative to non-Gulf veterans.

Conclusion and recommendations

This study found evidence of a small increased risk of infertility among veterans of the 1990-1 Gulf war, strengthened by the finding that pregnancies fathered by men with no fertility problems reportedly took longer to conceive than did those fathered by non-Gulf veterans. Put together with the previous finding of increased risk of miscarriage among pregnancies fathered by male UK Gulf veterans,⁹ we feel that the results of this study justify further research into the reproductive health of men deployed to the Gulf region, including a prospective investigation of semen quality and fecundability among veterans of the recent conflict in Iraq.

We thank the many people who supported the conduct of this study: representatives of the armed services; the British Legion, in particular Terry English; National Gulf Veterans and Families Association; Gulf Veterans Association; and, most importantly, the study participants for the time and effort involved. We also acknowledge the skills and commitment of those who worked on the study, particularly Tommy Clarke, Graham Davies, Haydon Hughes, Juliet Jain, Sam Lewis, Margo Pelerin, Susan Prior, Darren Reed, Patrick Sampson, and Janet Sullivan. For supplying cohort data and for invaluable help with queries, we thank all members of the Gulf Veterans Illness Unit (now the Veterans Policy Unit) at the Ministry of Defence, in particular Nick Blatchley, John Graham, Philip Bolton, Linda Walpole, and Chris Baker. We also thank Steve McManus and colleagues at the British Forces Post Office for providing valuable information on serving status and addresses.

Contributors: See bmj.com

Funding: The study was funded by the UK Ministry of Defence and administered by the UK Medical Research Council.

Competing interest: None declared.

Ethical approval: West Midlands multi-centre research ethics committee; London School of Hygiene and Tropical Medicine ethics committee.

- 1 Penman AD, Currier MM, Tarver RS. No evidence of increase in birth defects and health problems among children born to Persian Gulf veterans in Mississippi. *Mil Med* 1996;161:1-6.
- 2 Cowan DN, DeFraites RF, Gray GC, Goldenbaum MB, Wishik SM. The risk of birth defects among children of Persian Gulf veterans. *N Engl J Med* 1997;336:1650-6.
- 3 Araneta MRG, Moore CA, Olney RS, Edmonds LD, Karcher JA, McDonough C, et al. Goldenhar syndrome among infants born in military hospitals to Gulf veterans. *Teratology* 1997;56:244-51.
- 4 *Health study of Canadian forces personnel involved in the 1991 conflict in the Persian Gulf*. Vol 1. Canadian Department of National Defence, 1998.

(Prepared for the Gulf War Illness Advisory Committee, Department of National Defence, by Goss Gilroy Inc. Management Consultants.) www.dnd.ca/site/reports/Health/vol1_TOC_e.htm (accessed 10 May 2004).

- 5 Ishoy T, Andersson AM, Suadican P, Guldager B, Appleyard M. Major reproductive health characteristics in male Gulf veterans. *Dan Med Bull* 2001;48:29-32.
- 6 Kang H, Magee C, Mahan C, Lee K, Murphy F, Jackson L, et al. Pregnancy outcomes among U.S. Gulf veterans: a population-based survey of 30,000 veterans. *Ann Epidemiol* 2001;11:504-11.
- 7 Araneta MR, Schlangen KM, Edmonds LD, Destiche DA, Merz RD, Hobbs CA, et al. Prevalence of birth defects among infants of Gulf veterans in Arkansas, Arizona, California, Georgia, Hawaii, and Iowa, 1989-1993. *Birth Defects Res Part A Clin Mol Teratol* 2003;67:246-60.
- 8 Sim M, Abramson M, Forbes A, Glass D, Ikin J, Ittak P, et al. *Australian Gulf veterans' health study 2003*. Vol 2. Commonwealth Department of Veterans' Affairs, 2003. www.dva.gov.au/media/publicat/2003/gulfwarhs (accessed 10 May 2004).
- 9 Doyle P, Maconochie N, Davies G, Maconochie I, Pelerin M, Prior S, et al. Miscarriage, stillbirth and congenital malformation in the offspring of UK veterans of the first Gulf war. *Int J Epidemiol* 2004;33:74-86.
- 10 Maconochie N, Doyle P, Davies G, Lewis S, Pelerin M, Prior S, et al. The study of reproductive outcome and the health of offspring of UK veterans of the Gulf war: methods and description of the study population. *BMC Public Health* 2003;3:4.
- 11 World Health Organization. *WHO laboratory manual for the examination of human semen and sperm-cervical mucus interaction*. 3rd ed. Cambridge, UK: Cambridge University Press on behalf of WHO, 1992.
- 12 Breslow NE, Day NE. *Statistical methods in cancer research. Vol II. The analysis of cohort studies*. Lyon: International Agency for Research on Cancer, 1980.
- 13 Abou-Donia MB, Suliman HB, Khan WA, Abdel-Rahman AA. Testicular germ-cell apoptosis in stressed rats following combined exposure to pyridostigmine bromide, N-diethyl m-toluamide (DEET) and permethrin. *J Toxicol Environ Health A*. 2003;66:57-73.
- 14 Military Health Research Advisory Group (MHRAG). *MRC review of research into UK Gulf veterans' illnesses 2003*. London: Medical Research Council, 2003. www.mrc.ac.uk/pdf/gulf-illness-review.pdf (10 May 2004).

(Accepted 16 June 2004)

doi 10.1136/bmj.38163.620972.AE

Passive smoking and risk of coronary heart disease and stroke: prospective study with cotinine measurement

Peter H Whincup, Julie A Gilg, Jonathan R Emberson, Martin J Jarvis, Colin Feyerabend, Andrew Bryant, Mary Walker, Derek G Cook

Abstract

Objective To examine the associations between a biomarker of overall passive exposure to tobacco smoke (serum cotinine concentration) and risk of coronary heart disease and stroke.

Design Prospective population based study in general practice (the British regional heart study).

Participants 4729 men in 18 towns who provided baseline blood samples (for cotinine assay) and a detailed smoking history in 1978-80.

Main outcome measure Major coronary heart disease and stroke events (fatal and non-fatal) during 20 years of follow up.

Results 2105 men who said they did not smoke and who had cotinine concentrations < 14.1 ng/ml were divided into four equal sized groups on the basis of cotinine concentrations. Relative hazards (95% confidence intervals) for coronary heart disease in the second (0.8-1.4 ng/ml), third (1.5-2.7 ng/ml), and fourth (2.8-14.0 ng/ml) quarters of cotinine concentration compared with the first (≤ 0.7 ng/ml) were 1.45 (1.01 to 2.08), 1.49 (1.03 to 2.14), and 1.57 (1.08 to 2.28), respectively, after adjustment for established risk factors for coronary heart disease.

Hazard ratios (for cotinine 0.8-14.0 $v \leq 0.7$ ng/ml) were particularly increased during the first (3.73, 1.32 to 10.58) and second five year follow up periods (1.95, 1.09 to 3.48) compared with later periods. There was no consistent association between cotinine concentration and risk of stroke.

Conclusion Studies based on reports of smoking in a partner alone seem to underestimate the risks of exposure to passive smoking. Further prospective studies relating biomarkers of passive smoking to risk of coronary heart disease are needed.

Introduction

Meta-analyses examining the effect of living with a cigarette smoker on risk of coronary heart disease (CHD) among non-smokers have shown an overall increase in risk of about one quarter, after adjustment for potential confounding factors.^{1,2} Passive smoking may also be related to risk of stroke.³

Department of
Community Health
Sciences, St
George's Hospital
Medical School,
London SW17 0RE

Peter H Whincup
professor of
cardiovascular
epidemiology

Julie A Gilg
research statistician

Derek G Cook
professor of
epidemiology

continued over

BMJ 2004;329:200-4



This is the abridged version of an article that was posted on bmj.com on 30 June 2004: <http://bmj.com/cgi/doi/10.1136/bmj.38146.427188.55>