

### What is already known on this topic

Prevention of obesity is an increasingly important aspect of health promotion

Few trials have investigated school based primary prevention programmes directed at obesity

### What this study adds

The programme was successful in producing school level changes to tackle risk factors for obesity

High levels of participation indicated support from schools, staff, parents, and pupils

Positive changes were seen in school meals, tuck shops, and playground activities

Our programme used a population based approach underpinned by the Health Promoting Schools philosophy.<sup>10</sup> The population approach ensured that all children including those at risk of developing obesity were included. As the prediction of adult obesity from childhood measures is poor,<sup>11</sup> a population approach should have more effect at the public health level than targeting children who are already obese. Furthermore, behavioural and environmental conditions that contribute to the development and maintenance of obesity might be modified. Given

the positive effects achieved in all 10 schools, we believe that our programme could be successful in other primary schools.

We thank the staff, pupils, and parents of Brodetsky, Cookridge, Fir Tree, Horsforth St Mary's, Moortown, Morley The Newlands, Pudsey Lowtown, Shadwell, Springbank, and Whartons Otle primary schools for their participation, enthusiasm, and support. We also thank Nazia Chaudary and Julia Bartrop for conducting the focus groups and Gail Cook for helping with the evaluation of school meals.

Funding: Northern and Yorkshire Region Research and Development Unit funded the research.

Competing interests: None declared.

- 1 World Health Organization. *Preventing and managing the global epidemic*. Geneva: WHO, 1997.
- 2 Reilly JJ, Dorosty AR, Emmett PM. Prevalence of overweight and obesity in British children: cohort study. *BMJ* 1999;319:1039.
- 3 Rudolf MCJ, Sahota P, Barth JH, Walker J. Increasing prevalence of obesity in primary school children: cohort study. *BMJ* 2001;322:1094-5.
- 4 Fruhbeck G. Childhood obesity: time for action, not complacency. *BMJ* 2000;320:328-9.
- 5 Renisow K. School-based obesity prevention. *Ann N Y Acad Sci* 1993;699:154-6.
- 6 Story M. School-based approaches for preventing and treating obesity. *Int J Obes* 1999;23(suppl 2):S43-51.
- 7 NHS Centre for Reviews and Dissemination. The prevention and treatment of obesity. *Effective Health Care Bull* 1997;3 (No 2).
- 8 Tones K. The health promoting school: some reflections on evaluation. *Health Educ Res* 1996;11(4):i-viii.
- 9 Sahota P, Rudolf MCJ, Dixey R, Hill AJ, Barth JH, Cade J. Randomised controlled trial of primary school based intervention to reduce risk factors for obesity. *BMJ* 2001;323:1029-32.
- 10 Young I, Williams T, eds. *The healthy school*. Edinburgh: Scottish Health Education Group, World Health Organization, 1989.
- 11 Power C, Lake JL, Cole TJ. Measurement and longterm risks of child and adolescent fatness. *Int J Obes Metab Dis* 1997;21:507-26.

(Accepted 12 July 2001)

## Randomised controlled trial of primary school based intervention to reduce risk factors for obesity

Pinki Sahota, Mary C J Rudolf, Rachael Dixey, Andrew J Hill, Julian H Barth, Janet Cade

### Abstract

**Objective** To assess if a school based intervention was effective in reducing risk factors for obesity.

**Design** Group randomised controlled trial.

**Setting** 10 primary schools in Leeds.

**Participants** 634 children aged 7-11 years.

**Intervention** Teacher training, modification of school meals, and the development of school action plans targeting the curriculum, physical education, tuck shops, and playground activities.

**Main outcome measures** Body mass index, diet, physical activity, and psychological state.

**Results** Vegetable consumption by 24 hour recall was higher in children in the intervention group than the control group (weighted mean difference 0.3 portions/day, 95% confidence interval 0.2 to 0.4), representing a difference equivalent to 50% of baseline consumption. Fruit consumption was lower in obese children in the intervention group (-1.0, -1.8 to -0.2) than those in the control group. The three day diary showed higher consumption of high sugar foods (0.8, 0.1 to 1.6) among overweight children in the intervention group than the control group. Sedentary behaviour was higher in overweight

children in the intervention group (0.3, 0.0 to 0.7). Global self worth was higher in obese children in the intervention group (0.3, 0.3 to 0.6). There was no difference in body mass index, other psychological measures, or dieting behaviour between the groups. Focus groups indicated higher levels of self reported behaviour change, understanding, and knowledge among children who had received the intervention. **Conclusion** Although it was successful in producing changes at school level, the programme had little effect on children's behaviour other than a modest increase in consumption of vegetables.

### Introduction

School staff have access to large numbers of children in an environment that has the potential to support healthy behaviour and is favourable for the delivery of health promotion programmes.<sup>1</sup> Primary schools are particularly suitable for such programmes as children in this age group are responsive to health messages and behavioural changes may be maintained into adolescence and adulthood.<sup>2</sup>

Recent reports have indicated that over 17% of 11 year old children are obese and 30% overweight.<sup>3</sup>



The full version of this article appears on [bmj.com](http://bmj.com)

Editorial  
by Atkinson

School of Health Sciences, Leeds Metropolitan University, Leeds LS1 3HE

Pinki Sahota  
senior lecturer in nutrition and dietetics  
Rachael Dixey  
principal lecturer in health promotion

Leeds Community and Mental Health Trust, Belmont House, Leeds LS2 9DE

Mary C J Rudolf  
consultant community paediatrician

continued over

*BMJ* 2001;323:1029-32

School of Medicine,  
Leeds University,  
Leeds LS2 9LT  
Andrew J Hill  
senior lecturer in  
behavioural sciences

Leeds General  
Infirmary, Leeds  
LS1 3EX

Julian H Barth  
consultant in chemical  
pathology and  
metabolic medicine

Nuffield Institute of  
Health, Leeds  
LS2 9PL

Janet Cade  
senior lecturer in  
nutritional  
epidemiology

Correspondence to:  
M C J Rudolf  
Mary.Rudolf@  
leedsth.nhs.uk

School based programmes might be able to reverse the increase in obesity. Programmes targeted at obese children and adolescents have reported positive results, particularly those aimed at primary school aged children<sup>1</sup> and those combining diet and exercise with parental involvement. However, most of these studies not only had inadequate methods<sup>5</sup> but also had the problem that they targeted obese children (often volunteered by parents). Such targeting may increase stigmatisation of children at school. An alternative strategy is to implement a health promotion programme aimed at all pupils or pupils at high risk of becoming overweight

Only a few primary prevention studies of school based interventions have been reported from the United States,<sup>6</sup> and none in the United Kingdom.<sup>7</sup> We designed a multicomponent health promotion programme, based on the Health Promoting Schools concept,<sup>8</sup> aimed at reducing risk factors for obesity in primary schools. It was devised as a group randomised trial so that its effect could be appropriately assessed. We targeted children aged 8-10 because they were cognitively able to complete questionnaires and because levels of obesity start to rise around this age.<sup>3</sup>

## Participants and methods

We recruited 10 state primary schools in Leeds and paired them according to size, ethnicity, and level of social disadvantage (as reflected by numbers of free school meals). We randomised them to receive the intervention or to serve as the comparison school using the toss of a coin. A power calculation indicated

that with five schools in each arm, the study would have 80% power to detect an underlying difference in means of a normally distributed outcome measure of  $\geq 1.8$  standard deviations at the 5% significance level and 65% power to detect a difference of 1.5 SD. This took into account the cluster randomisation design. Sociodemographic measures suggested that the schools' populations generally reflected the Leeds school aged population.

The intervention schools received the active programme promoting lifestyle education in schools (APPLES). The programme consisted of teacher training, modifications of school meals, and the development and implementation of school action plans designed to promote healthy eating and physical activity over one academic year (September 1996-July 1997). The comparison schools continued with their usual health curriculum, without the intervention. The philosophy and details of the intervention are described in the accompanying paper, together with an evaluation of the implementation and effect of the programme.<sup>9</sup> The study was approved by the local research ethics committee and the schools' governing bodies. Parental consent was also obtained.

## Main outcome measures

We assessed the effect of the intervention on individual behaviour by collecting data on growth, diet, physical activity, and psychological state at baseline and 12 months later. These measures could not be obtained blind to the schools' intervention status.

**Growth measures**—Weights were recorded without shoes, jumpers, or sweatshirts. Heights were measured by a standard method.<sup>3</sup> We then calculated body mass index.

**Dietary information** was assessed by both 24 hour recall and three day food diaries. We analysed the information from both methods for frequency of consumption of foods high in fat, foods and drinks high in sugars, fruit, and vegetables.


**Physical activity**—The frequency of physical activity and sedentary behaviour was measured by questionnaire and was categorised by frequency of sport and physical activity during the past week and frequency of sedentary behaviour in past 24 hours.

**Psychological measures**—We used three validated tools to evaluate the effect of the intervention on psychological well being: the self perception profile for children<sup>10</sup>; a measure of dietary restraint that has been used in children aged 8<sup>11</sup>; and the adapted body shape perception scale, which consists of a series of seven line drawings ranging from extremely thin to obese for self assessment of the individual's perception of their body size.<sup>12</sup>

To determine the effect on children's levels of knowledge and attitudes towards healthy living, 80 focus groups involving 320 children were held at the end of year. Facilitators were blind to the school's allocation status.

## Analysis of data

We applied a multilevel statistical model using STATA<sup>13</sup> to assess changes in body mass index, diet, physical activity, and psychological well being. This statistical model took into account the pairing and lack of independence among subjects within the school, known as the clustering effect.<sup>14</sup> In addition, we adjusted for sex,



Progress of schools and pupils through trial

age, baseline body mass index, and outcome of interest at baseline and the intervention status.

Body mass index, height, and weight were converted to standard deviation scores using the revised 1990 reference standards.<sup>15</sup> We assessed the effect of the intervention by change in score after adjusting for age, sex, initial body mass index SD score, and type of school. All outcome measures were assessed in the whole cohort, in overweight children (body mass index > 1.04, equivalent to > 85th centile) and in obese children (body mass index > 1.64, equivalent to 95th centile on the 1990 body mass index growth charts).

## Results

### Characteristics of sample

All 10 schools completed the project (figure). In all, 636 children (314 intervention, 322 comparison) took part in the evaluation. The mean (SD) age of children in the intervention schools was 8.4 (0.63) years, with 145 (46%) entering school year 4 and 169 (54%) year 5. The mean age of the children in the comparison schools was also 8.4 (0.63) years, with 145 (45%) entering year 4 and 177 (55%) year 5. Baseline characteristics in terms of growth measures, diet, physical activity, and psychological scores were similar in the intervention and comparison groups.

### Growth

Overall, there was no difference in score between the intervention and control children at the end of the year (table 1). There was also no difference for the overweight (weighted mean difference  $-0.07$ , 95% confidence interval  $-0.22$  to  $0.08$ ) or the obese children ( $-0.05$ ,  $-0.22$  to  $0.11$ ).

### Diet

Analysis of 24 hour recall showed that the intervention children had higher vegetable intakes at the end of the study (tables 1 and 2). The weighted mean difference of  $0.3$  (95% confidence interval  $0.2$  to  $0.4$ ) indicates on average one third of a portion more a day. As the mean baseline was only  $0.6$  portions/child/day, this difference is equivalent to 50% of baseline intake. The same difference was seen for the overweight ( $0.3$ ,  $0.1$  to  $0.5$ ) and obese children ( $0.3$ ,  $-0.1$  to  $0.6$ ). The three day diary (which had a quantitatively and qualitatively lower completion rate) did not show these differences.

Other significant differences included a lower fruit intake in obese intervention children by 24 hour recall. According to the three day diaries, there was a higher intake of foods and drinks high in sugar in overweight children in the intervention group.

### Physical activity and psychological measures

We found no significant difference in physical activity or sedentary behaviour for the sample as a whole. The only significant difference in psychological measures was a small increase in global self worth for obese children in the intervention schools ( $0.32$ ,  $0.0$  to  $0.64$ ).

### Focus groups

Compared with children in control schools, children in the intervention schools had higher levels of self reported behaviour change, greater understanding of the health benefits of diet and physical activity, and increased sophistication of ideas and vocabulary,

**Table 1** Weighted mean difference in body mass index standard deviation score and vegetable intake between the five intervention schools and their control schools

	Body mass index SD score		Vegetable intake	
	Weighted mean difference (95% CI)	% weight of school*	Weighted mean difference (95% CI)	% weight of school*
1	0 ( $-0.2$ to $0.1$ )	25.8	0.2 ( $-0.1$ to $0.4$ )	25.5
2	0.1 ( $0$ to $0.2$ )	18.0	0.4 ( $0.2$ to $0.7$ )	18.2
3	0.1 ( $-0.1$ to $0.2$ )	22.5	0.3 ( $0.1$ to $0.5$ )	23.0
4	$-0.1$ ( $-0.3$ to $0$ )	19.8	0.4 ( $0.1$ to $0.7$ )	16.0
5	$-0.2$ ( $-0.3$ to $0$ )	13.9	0.1 ( $-0.1$ to $0.4$ )	17.4
Overall	0 ( $-0.1$ to $0.1$ )		0.3 ( $0.2$ to $0.4$ )	

\*Statistical weighting.

**Table 2** Weighted mean difference (95% confidence interval) in dietary and physical activity levels between intervention and control children according to weight

	All children	Overweight	Obese
24 hr recall:			
Foods high in fat	0.1 ( $-0.2$ to $0.4$ )	0.4 ( $-0.0$ to $0.9$ )	0.1 ( $-0.7$ to $0.8$ )
Foods and drinks high in sugar	$-0.5$ ( $-1.1$ to $0.1$ )	0.0 ( $-0.8$ to $0.8$ )	1.0 ( $-0.5$ to $2.6$ )
Fruit intake	0.0 ( $-0.5$ to $0.5$ )	$-0.2$ ( $-1.0$ to $0.6$ )	$-1.0$ ( $-1.8$ to $-0.2$ )
Vegetable intake	0.3 ( $0.2$ to $0.4$ )	0.3 ( $0.1$ to $0.5$ )	0.3 ( $-0.1$ to $0.6$ )
3 day diary:			
Foods high in fat	$-0.1$ ( $-0.4$ to $0.3$ )	0.2 ( $-0.9$ to $1.2$ )	0.0 ( $-0.8$ to $0.8$ )
Foods and drinks high in sugar	0.1 ( $-0.4$ to $0.5$ )	0.8 ( $0.1$ to $1.6$ )	0.3 ( $-1.8$ to $2.3$ )
Fruit intake	$-0.2$ ( $-0.6$ to $0.2$ )	$-0.2$ ( $-0.7$ to $0.3$ )	$-0.6$ ( $-1.2$ to $0.1$ )
Vegetable intake	0.1 ( $-0.1$ to $0.3$ )	$-0.1$ ( $-0.5$ to $0.2$ )	$-0.1$ ( $-0.9$ to $0.7$ )
Physical activity:			
Physical activity	$-0.2$ ( $-0.4$ to $0.1$ )	$-0.2$ ( $-0.5$ to $0.2$ )	$-0.48$ ( $-1.3$ to $0.3$ )
Sedentary behaviour	0.0 ( $-0.1$ to $0.1$ )	0.3 ( $0.0$ to $0.7$ )	0.29 ( $-0.1$ to $0.7$ )

A positive value for weighted mean difference indicates higher food intake or activity levels in the intervention schools than the control schools. A value of zero indicates no difference.

willingness and confidence to share their ideas, and basic knowledge. They were more able to recollect topics learnt and activities undertaken in school linked to diet and physical activity.

## Discussion

Implementation of this health promotion programme was highly successful,<sup>9</sup> and it was therefore disappointing that the children showed minimal behavioural changes. The only clinically important positive result was a modest increase in consumption of vegetables. Vegetables are likely to be the most challenging food group to change in children, and a rise of one third of a vegetable portion per child is a huge increase in vegetable consumption across the city per day. Nevertheless, it is a small return for the effort made.

One possible reason for the apparent lack of effectiveness of the intervention could be the alarming increase in the prevalence of overweight and obesity in this entire school population.<sup>3</sup> This rise in body mass index was seen as the children grew older as well as over time. In the face of such an epidemic, a programme with limited resources would be unlikely to have demonstrable effects in countering this increase. Clearly, the social and environmental forces at work require much larger public health counter measures.

### Problems of evaluating complex interventions

Although about 600 children participated, the unit size was only five schools in each arm, which is very small. More schools are needed to achieve a clear result. This would have been more costly and difficult to implement. Several larger health promotion trials have also had results below expectations for the same reason.<sup>16-21</sup>

### What is already known on this topic

Obesity is increasing among school children and demands preventive strategies

Randomised controlled trials of school based primary prevention programmes have all used a prescriptive approach

### What this study adds

Despite changes at school level, behavioural changes were disappointing with this programme based on the health promoting schools philosophy

The only positive outcome was a modest increase in vegetable consumption

The discrepancy between changes achieved at the individual and school level raises issues regarding the problems inherent in such trials

The outcome measures that we used also presented difficulties. Body mass index is reliably measurable but could not be expected to change significantly over such a short time. Indeed, had body mass index fallen, we would have been concerned that we had induced inappropriate eating behaviour. The other outcome measures were less easy to measure. Although the focus groups indicated a change in self reported behaviour, we had no quantitative evidence of this. Accurate dietary assessment is difficult in any age group, but particularly in children. Few validated instruments are available, and the two methods that we used could not assess quantities accurately. We therefore relied more on the quality of food reported. The reports of physical activity levels were even harder to quantify.

There is much debate about whether randomised controlled trials are appropriate for evaluating complex interventions, particularly in primary prevention.<sup>22</sup> Our study, with others, raises critical issues for the development of evidence based health promotion and about appropriate measures to determine whether such interventions are effective and worthwhile.

### Conclusions

Since the programme was successful in changing the ethos of the schools and the attitudes of the children,<sup>9</sup> it is premature to conclude that it was unsuccessful in reducing risk factors for obesity. The intervention was designed for one academic year only, recognising funding limitations. The intervention might have been strengthened if the families were targeted more directly,<sup>23</sup> while still maintaining the focus on the school community.

Other studies of this type suggest that behaviour change is possible over time.<sup>24 25</sup> Given the problems of evaluating behavioural intervention trials, it has been recommended that funding agencies should consider supporting such trials in three phases (a development and validation phase, a small feasibility phase, and a larger multicentre trial) instead of looking for immediate positive results.<sup>24</sup> If we view our study as a first and second phase trial, the programme was a success.

We thank the staff, pupils, and parents of Brodetsky, Cookridge, Fir Tree, Horsforth St Mary's, Moortown, Morley The Newlands, Pudsey Lowtown, Shadwell, Springbank, and the Whartons Otley primary schools for their participation, enthusiasm, and support. We also thank Jenny Walker, paediatric endocrine sister for collecting the anthropometric data, and Andy Vail for statistical advice.

Funding: The research was funded by a grant from the Northern and Yorkshire Region Research and Development Unit.

Competing interests: None declared.

- Garrow JS. Importance of obesity. *BMJ* 1991;303:704-6.
- Dietz WH. Childhood obesity. In: Bjorntorp P, Brodoff BN, eds. *Obesity*. Philadelphia: Lipincott, 1992:606-9.
- Rudolf MCJ, Sahota P, Barth JH, Walker J. Increasing prevalence of obesity in primary school children: cohort study. *BMJ* 2001;322:1094-5.
- Parcel GS, Simons-Morton BG, O'Hara NM, Baranowski T. School-based programmes to prevent or reduce obesity. In: Krasnegor NA, ed. *Childhood obesity: a biobehavioral perspective*. Caldwell, NJ: Telford, 1988:143-7.
- Reniscow K. School-based obesity prevention. *Ann N Y Acad Sci* 1993;699:154-6.
- Story M. School-based approaches for preventing and treating obesity. *Int J Obes Metab Dis* 1999;23(suppl 2):S43-51.
- NHS Centre for Reviews and Dissemination. The prevention and treatment of obesity. *Effective Health Care Bull* 1997;3 (No 2).
- Young I, Williams T, eds. *The healthy school*. Edinburgh: Scottish Health Education Group, World Health Organization, 1989.
- Sahota P, Rudolf MCJ, Dixey R, Hill AJ, Barth JH, Cade J. Evaluation of implementation and effect of primary school based intervention to reduce risk factors for obesity. *BMJ* 2001;323:1027-9.
- Harter S. *Manual for the self-perception profile for children*. Denver: University of Denver, 1985.
- Hill AJ, Pallin V. Dieting awareness and low self worth: related issues in 8 year old girls. *Int J Eating Disorders* 1998;24:405-13.
- Hill AJ, Draper E, Stack J. A weight on children's minds: body shape dissatisfaction at 9 years old. *Int J Eating Disorders* 1994;18:383-9.
- StataCorp. *Stata statistical software; release 6.0*. College Station, TX: Stata Corporation, 1999.
- Thompson SG, Pyke SDM, Hardy RJ. The design and analysis of paired cluster randomised trials: an appreciation of meta-analysis techniques. *Stat Med* 1997;16:2063-79.
- Cole TJ. *Castlemead growth programmes*. Welwyn Garden City: Castlemead Publications, 1993.
- Luepker RV, Perry CL, McKinlay S, Nader PR, Parcel GS, Stone EJ, et al. Outcomes of a field trial to improve children's dietary patterns and physical activity: The child and adolescent trial for cardiovascular health (CATCH). *JAMA* 1996;275:768-76.
- Farquhar JW, Fortmann SP, Flora JA, Taylor CB, Haskell WL, Williams PT, et al. Effect of community-wide education on cardiovascular disease risk factors: The Stanford five-city project. *JAMA* 1990;264:359-65.
- Luepker RV, Murray DM, Jacobs DR Jr, Mittelmark MB, Bracht N, Carlaw R, et al. Community education for cardiovascular disease prevention: risk factor changes in the Minnesota heart health programme. *Am J Public Health* 1994;84:1383-93.
- Carleton RA, Lasaster TM, Assaf AR, Feldman HA, McKinlay S. The Pawtucket heart health program: community changes in cardiovascular risk factors and projected disease risk. *Am J Public Health* 1995;85:777-85.
- COMMITT Research Group. Community intervention trial for smoking cessation (COMMITT): I. Cohort results from a four year community intervention. *Am J Public Health* 1995;85:183-92.
- COMMITT Research Group. Community intervention trial for smoking cessation (COMMITT): II. Changes in adult cigarette smoking prevalence. *Am J Public Health* 1995;85:193-200.
- Green J, Tones K. Towards a secure evidence base for health promotion. *J Public Health Med* 1999;21:133-9.
- Fruhbeck G. Childhood obesity: time for action, not complacency. *BMJ* 2000;320:328-9.
- CART Project Team. Developing methodologies for evaluating community-wide health promotion. *Health Promotion International* 1996;11(3):227-35.
- Contento IR, Balch GI, Bronner YL, Lytle LA, Maloney SK, White SL, et al. Nutrition education for school-aged children. *J Nutr Educ* 1995;27: 298-311.

(Accepted 12 July 2001)

### Endpiece

#### An obstacle course

He had been raised to speak French, which the school regarded less as a key to another culture than as an obstacle course and a brain teaser.

Robertson Davies (1913-95), *What's bred in the bone*, London: Penguin, 1986