

Effect of sublingual misoprostol on severe postpartum haemorrhage in a primary health centre in Guinea-Bissau: randomised double blind clinical trial

Lars Høj, Placido Cardoso, Birgitte Bruun Nielsen, Lone Hvidman, Jens Nielsen, Peter Aaby

Abstract

Objective To evaluate whether routine administration of sublingual misoprostol 600 µg after delivery reduces postpartum haemorrhage.

Design Randomised double blind placebo controlled trial.

Setting Primary health centre in Bissau, Guinea-Bissau, West Africa.

Participants 661 women undergoing vaginal delivery.

Intervention Misoprostol 600 µg or placebo administered sublingually immediately after delivery.

Main outcome measures Postpartum haemorrhage, defined as a loss of ≥ 500 ml and decrease in haemoglobin concentration after delivery.

Results The incidence of postpartum haemorrhage was not significantly different between the two groups, the relative risk being 0.89 (95% confidence interval 0.76 to 1.04) in the misoprostol group compared with the placebo group. Mean blood loss was 10.5% (-0.5% to 20.4%) lower in the misoprostol group than in the control group. Severe postpartum haemorrhage of ≥ 1000 ml or ≥ 1500 ml occurred in 17% (56) and 8% (25) in the placebo group and 11% (37) and 2% (7) in the misoprostol group. Significantly fewer women in the misoprostol group experienced a loss of ≥ 1000 ml (0.66, 0.45 to 0.98) or ≥ 1500 ml (0.28, 0.12 to 0.64). The decrease in haemoglobin concentration tended to be less in the misoprostol group, the mean difference between the two groups being 0.16 mmol/l (-0.01 mmol/l to 0.32 mmol/l).

Conclusion Sublingual misoprostol reduces the frequency of severe postpartum haemorrhage.

Introduction

In low income countries many pregnant women are anaemic.¹ Maternal mortality is four times higher in severely anaemic women than in those who are not anaemic.² In Guinea-Bissau, West Africa, 31% of pregnant women are anaemic³ and the maternal mortality is more than 8 per 1000 live births, postpartum haemorrhage being the most common cause of death.⁴ The primary cause of postpartum haemorrhage is uterine atony, which accounts for 70% of cases.⁵

Several drugs reduce postpartum haemorrhage by causing the uterus to contract. Ergot derivatives have

been used for decades, although oxytocin is the drug of choice in some centres. Several prostaglandins are used as second or third line agents. These drugs, however, must be refrigerated to remain effective. Moreover, most uterotonics must be administered by injection, which requires sterile equipment and training in safe administration, prerequisites unavailable for most women delivering in low income countries.

Misoprostol, a prostaglandin E1 analogue, is heat stable and can be administered orally, rectally, or sublingually. A multicentre study sponsored by the World Health Organization found that misoprostol was less effective for prophylaxis than intravenous or intramuscular injections of oxytocin but did not investigate the possible benefit of misoprostol to the large number of women who give birth outside a health facility with electricity.⁶ Community based distribution of misoprostol in Indonesia reduced the perceived frequency of excessive bleeding and the need for emergency referral for postpartum haemorrhage compared with data from a control area where misoprostol was not available.⁷

To date all randomised studies of prophylactic misoprostol have used oral and rectal administration, though a recent pharmacokinetic study showed that sublingual administration secures the highest peak concentration and the best bioavailability.⁸ A recent pilot study found that sublingual misoprostol and intravenous syntometrine have comparable effects on blood loss in the third stage of labour.⁹ None of the above clinical studies, however, was sufficiently blinded. We tested whether routine administration of sublingual misoprostol could reduce the incidence of postpartum haemorrhage in a resource poor, primary healthcare setting in Guinea-Bissau, West Africa, when assessed in an adequately blinded manner.

Methods

Setting and participants

In Bissau, the capital of Guinea-Bissau, most women give birth at the national hospital—the Hospital

Department of Obstetrics and Gynaecology, Aarhus University Hospital, 8200 Aarhus N, Denmark

Lars Høj
staff specialist

Birgitte Bruun Nielsen
staff specialist

Lone Hvidman
consultant obstetrician

Projecto de Saúde de Bandim, Apartado 861, Bissau, Guinea-Bissau
Placido Cardoso
researcher

Bandim Health Project, Danish Epidemiology Science Centre, Statens Serum Institut, 2300 Copenhagen S, Denmark

Jens Nielsen
statistician
Peter Aaby
professor

Correspondence to:
L Høj
lars.hoj@dadlnet.dk

BMJ 2005;331:723–7



This is an abridged version. The full version is on bmj.com

National de Simão Mendes (HNSM). A growing number, however, give birth at local health centres such as the Centro de Saúde de Bandim, where four midwives attend about 500 deliveries a year. Midwives may seek support from a general physician, not trained in obstetrics. If complications occur, women are transferred immediately to the national hospital, though this is often delayed because of lack of transport. Public electricity is not usually available and the generator at the health centre works less than half of the nights because of lack of diesel or spare parts. There is no running water at the centre. The local health centre is supplied with essential drugs, but they are often out of stock and relatives are often sent to the nearest pharmacy to buy drugs as necessary.

Treatment protocol and consent

In the first month of the trial, an expatriate midwife with experience in research was posted to the health centre. All four local (auxiliary) midwives and the physician were informed about the project, and all staff members were trained during a two week period. The midwives were taught how to apply controlled cord traction and how to cut and clamp the cord at delivery. Though we attempted to improve all routines at the maternity centre, the only intervention studied was the impact of misoprostol on postpartum haemorrhage.

All women who gave birth at the local centre from March 2003 to August 2004 were asked to take part in the study. When the women arrived at the health centre, the midwife in charge explained the purpose of the study to the women. As most women of child bearing age are illiterate we obtained verbal consent. All the women were given the same clinical care. Before the delivery the midwife let the women choose an envelope containing three tablets of either misoprostol or placebo, and these were administered sublingually as soon as the baby had been delivered. Most women received the tablets within two minutes of delivery.

In cases of severe postpartum haemorrhage, local guidelines for treatment were followed; oxytocin was freely available to the midwives during the trial.

Randomisation and masking

Misoprostol and placebo tablets of identical form, size, colour, and packing were produced by U-liang Pharmaceutical, Taipei, Taiwan. Seven hundred opaque envelopes were consecutively numbered and filled with either three tablets of placebo or three tablets of misoprostol 200 µg, distributed randomly by using a list of random numbers.

Measurement of blood loss

After delivery of the baby and drainage of the amniotic fluid, a clean plastic lined absorbent drape was placed under the woman's buttocks to collect all the blood lost. The drape was changed as many times as needed. The woman stayed on the drape or was asked to wear a pad over the next 60 minutes. In the case of severe haemorrhage, midwives were instructed to follow the usual guidelines for management of postpartum haemorrhage, and the supplemental treatment was registered. All drapes and pads were weighed on an electronic scale and the known dry weight of the linen was subtracted. As 1 ml of blood weighs close to 1 g, the balance in grams was assumed to be the total blood loss in ml.




Fig 1 Distribution of postpartum blood loss in women according to treatment

Haemoglobin concentration was measured by finger prick before and 24 hours after delivery. If the woman left the local centre within 20 hours of delivery, she was visited at home to have her haemoglobin concentration measured.

Outcome measures

The primary outcome measure was postpartum haemorrhage, defined as a blood loss of ≥ 500 ml. We analysed the blood loss, the incidence of different thresholds of blood loss, change in haemoglobin concentration between admission and discharge, and the incidence of a 10% decrease in haemoglobin concentration between admission and discharge.

The midwives documented potential side effects by interviewing the women about nausea, diarrhoea, and vomiting one hour after delivery. The midwife in charge measured rectal temperature and recorded any shivering.

Power calculation and analysis

The average blood loss in the third stage of labour is 250-350 ml, and 12% of women will lose > 500 ml.¹⁰ Use of uterotonics should reduce this proportion to 5%.¹⁰ We calculated that we needed 300 women in each group to have a power of 80% ($1 - \beta$) with a risk of type 2 error (α) of 5%.

We entered data on to dBase V and used Stata version 9.0 for Windows for statistical analysis. Transformed blood loss (fig 1) and decrease in haemoglobin concentration approximated normal distributions (fig 2), so we analysed log (blood loss) and decrease in haemoglobin using linear regression. We used Pearson's χ^2 test to compare categorical variables and risks ratios to express relative risks.




Fig 2 Distribution of decrease in haemoglobin concentration after delivery in women according to treatment

Results

Randomisation and progress through trial

A total of 661 women were enrolled in the study, 330 in the misoprostol group and 331 in the placebo group. All eligible women were included and received the randomised drug; there was no loss to follow-up and no one discontinued the intervention. Hence, all enrolled women were included in the analysis (see bmj.com).

Baseline characteristics

Few women in Bissau possess a reliable birth certificate; the median reported age of enrolled women was 23 years (interquartile range 19-27 years). The median weight was 61 kg (interquartile range 57-69 kg). There were no significant differences between the two groups with respect to duration of the third stage of labour, weight, or Apgar score.

Outcome measures

Of the 661 women, 150 (45%) in the misoprostol group and 170 (51%) in the control group had

postpartum haemorrhage of ≥ 500 ml, the relative risk being 0.89 (0.76 to 1.04). The mean blood loss was 10.5% (-0.5% to 20.4%) lower in the misoprostol group than in the placebo group (table 1). Significantly fewer women in the misoprostol group (11%) suffered from severe postpartum haemorrhage, with a blood loss of ≥ 1000 ml (11% *v* 17%; 0.66, 0.45 to 0.98). For a blood loss of ≥ 1500 ml, the relative risk between the misoprostol group (2%) and the control group (8%) was 0.28 (0.12 to 0.64).

Differences in haemoglobin concentrations at admission and one day postpartum varied considerably (table 1); the haemoglobin concentration increased in 33% (218/661) of the women. The mean decrease in haemoglobin concentration was 0.16 mmol/l (-0.01 mmol/l to 0.32 mmol/l) lower in the misoprostol group than in the placebo group (table 1). The proportion of women with a reduction of more than 10% of the admission value did not differ between the two groups.

Side effects

Significantly more women in the misoprostol group experienced shivering and pyrexia (table 2). There were few complaints about nausea, and few in either group had vomiting or diarrhoea. One woman in the misoprostol group died. She was a 19 year old primigravida who gave birth to a 2630 g boy and had postpartum haemorrhage of 1417 ml. Apparently this was under control, but she suddenly died 90 minutes after delivery. The verbal autopsy audit reached no definitive conclusion as to the cause of death.

Discussion

Sublingual misoprostol is effective in reducing the incidence of severe postpartum haemorrhage. Though we expected 12% of the women in the control group to have a blood loss of >500 ml,¹⁰ our measurement technique showed that as many as 51% of the women in the control group had a loss of ≥ 500 ml. We therefore redefined severe postpartum haemorrhage as a loss of ≥ 1000 ml or ≥ 1500 ml. These levels of bleeding affected 17% and 8% of the women in the control group—that is, proportions exceeding those estimated in our power calculations. Hence, the present randomised double blinded study showed that routine misoprostol had a clear protective effect on severe postpartum haemorrhage in excess of 1000 ml. Also, the reduction in haemoglobin concentrations after delivery was less in the women in the misoprostol group.

Table 1 Blood loss and change in haemoglobin concentration in women according to treatment group. Means are shown with 95% confidence intervals in brackets

| | Misoprostol (n=330) | Control (n=331) | Relative risk or mean difference (95% CI) |
|--|---------------------|---------------------|---|
| No (%) with blood loss ≥ 500 ml | 150 (45) | 170 (51) | 0.89 (0.76 to 1.04) |
| Mean measured blood loss (ml)* | 443 (408 to 481) | 496 (456 to 538) | 10.5 (-0.5 to 20.4)† |
| No (%) with blood loss ≥ 1000 ml | 37 (11) | 56 (17) | 0.66 (0.45 to 0.98) |
| No (%) with blood loss ≥ 1500 ml | 7 (2) | 25 (8) | 0.28 (0.12 to 0.64) |
| Mean decrease in Hb concentration (mmol/l) | 0.29 (0.17 to 0.41) | 0.45 (0.33 to 0.57) | 0.16 (-0.01 to 0.32)† |
| No (%) with 10% fall in Hb concentration | 105 (32) | 115 (35) | 0.92 (0.74 to 1.14) |

HB=haemoglobin.

*Log transformed.

†Mean difference.

Table 2 Side effects after delivery in women who received misoprostol to treat postpartum haemorrhage. Figures are numbers (percentages) of women

| | Misoprostol (n=330) | Control (n=331) | Relative risk (95% CI) |
|---------------------------------------|------------------------|--------------------|------------------------|
| Shivering | 189 (57) | 78 (24) | 2.43 (1.96 to 3.01) |
| Fever ($\geq 38.0^{\circ}\text{C}$) | 78 (24) | 11 (3) | 7.09 (3.84 to 13.1) |
| Nausea | 2 (1) | 4 (1) | 0.50 (0.09 to 2.72) |
| Vomiting | 10 (3) | 4 (1) | 2.51 (0.79 to 7.91) |
| Diarrhoea | 10 (3) | 4 (1) | 2.50 (0.79 to 7.89) |

Comparison with other studies

As in other studies, we found a significant increase in shivering and pyrexia in the misoprostol group.^{6 11 12} According to the midwives, however, these side effects were transient and of little disadvantage to the women.

Four randomised controlled trials tested oral and rectal misoprostol against placebo and found no significant reduction in the number of women experiencing postpartum haemorrhage, if the threshold was 500 ml.^{11 13-15} The explanation for this finding might be found in the study done by Tang et al, which compared the pharmacokinetics of three routes of administration: sublingual, oral, and vaginal.⁸ The time to peak concentration was shortest in the sublingual route, at 26.0 minutes (SD 11.5 minutes). As bleeding in an uncomplicated third stage of labour normally ceases within 10 minutes, we would not expect to see any effect of misoprostol on "normal" bleeding. Continuous bleeding for more than 15-25 minutes, however, should be reduced by its uterotonic properties.

Strengths of study

Most studies on misoprostol as a uterotonic agent in postpartum haemorrhage have been insufficiently blinded because the company that owns the patent has been unwilling to cooperate in studies of obstetric use of misoprostol by manufacturing a placebo. However, the rights have now been sold and we were able to obtain the active and the placebo tablets manufactured and packed so that they were completely identical.

Estimation of blood loss is known to be difficult and inaccurate. In several of the previous studies, the birth attendant estimated blood loss, a method known to underestimate the quantity by about 30%.¹⁶ We tried to estimate blood loss more precisely, though we could have overestimated it because of the addition of amniotic fluid or underestimated it because of loss of blood outside the drapes. As these errors are likely to be distributed equally between the two study groups they are unlikely to have introduced any systematic bias that could have affected the significance of our results.

Conclusion

In rural Guinea-Bissau, 75% of women give birth at home,¹⁷ and worldwide only about 50% of women give birth in health facilities. Therefore, strategies must be identified to increase the safety of deliveries attended by unskilled birth attendants. According to the study from Indonesia,⁷ it might be safe and useful to give misoprostol to women during antenatal care and to teach midwives to use the drug safely. Our randomised trial suggests that misoprostol would play an important part in such a strategy to reduce complications of delivery and maternal mortality.

What is already known on this topic

Conventional uterotonics require refrigeration and equipment for injection and are therefore inappropriate for the control of postpartum haemorrhage in poor areas with little infrastructure

Though misoprostol tablets do not require refrigeration, oral misoprostol 600 μg is less effective than conventional uterotonics in reducing postpartum haemorrhage of ≥ 1000 ml

What this study adds

Sublingual misoprostol 600 μg reduces the incidence of severe haemorrhage (≥ 1000 ml)

Misoprostol tablets may reduce the morbidity and mortality associated with postpartum haemorrhage in the developing world

More studies on the use of misoprostol outside developed health facilities are warranted. If the drug is found to be consistently beneficial and safe, sublingual misoprostol should be offered to labouring women at the start of the third stage of labour if injectable uterotonics are not available. The mothers should be informed that shivering and mild fever can be expected.

Contributors: LH was the principal investigator, contributed to the study design, wrote the draft manuscript, and is guarantor. BBN, LH, and PA contributed to the study design and were involved in the data interpretation and writing of the manuscript. PC supervised the execution of the study, did the data entry, and contributed to the write up. Rosa Amelia Caceres Perez taught the midwives, who were Angelina da Silva, Ermelinda Tchuda, Suzana Rodrigues, and Felismina Ant3nio Gomes. JN conducted the statistical analysis and contributed to the data interpretation.

Funding: The Danish Society of Obstetrics and Gynaecology, the Illum Foundation, and the Danish International Development Agency.

Competing interests: None declared.

Ethical approval: The scientific committee of the Ministry of Health in Guinea-Bissau and the Danish central scientific ethical committee (No 624-02-0036).

- Rush D. Nutrition and maternal mortality in the developing world. *Am J Clin Nutr* 2000;72:212S-40S.
- Brabin BJ, Hakimi M, Pelletier D. An analysis of anemia and pregnancy-related maternal mortality. *J Nutr* 2001;131:604-615S.
- Kaestel P. Micronutrient supplementation and other predictors of birth size and perinatal mortality in Guinea-Bissau [thesis]. Copenhagen, Denmark: Department of Epidemiology Research, Statens Serum Institute, 2004.
- Hoj L, Stensballe J, Aaby P. Maternal mortality in Guinea-Bissau: the use of verbal autopsies in a multi-ethnic population. *Int J Epidemiol* 1999;28:70-6.
- Kwast BE. Postpartum haemorrhage: its contribution to maternal mortality. *Midwifery* 1991;7:64-70.
- Gulmezoglu AM, Villar J, Ngoc NT, Piaggio G, Carroli G, Adetoro L, et al. WHO multicentre randomised trial of misoprostol in the management of the third stage of labour. *Lancet* 2001;358:689-95.
- Maternal and Neonatal Health Program. *Preventing postpartum hemorrhage: a community-based approach proves effective in rural Indonesia*. Baltimore USA: MNH Program, 2005.
- Tang OS, Schweer H, Seyberth HW, Lee SW, Ho PC. Pharmacokinetics of different routes of administration of misoprostol. *Hum Reprod* 2002;17:332-6.
- Lam H, Tang OS, Lee CP, Ho PC. A pilot-randomized comparison of sublingual misoprostol with syntometrine on the blood loss in third stage of labor. *Acta Obstet Gynecol Scand* 2004;83:647-50.
- Prendiville WJ, Elbourne D, McDonald S. Active versus expectant management in the third stage of labour. *Cochrane Database Syst Rev* 2000;(3):CD000007.

- 11 Hofmeyr GJ, Nikodem VC, de Jager M, Drakely A. Side-effects of oral misoprostol in the third stage of labour—a randomised placebo-controlled trial. *S Afr Med J* 2001;91:432-5.
- 12 Lumbiganon P, Villar J, Piaggio G, Gulmezoglu AM, Adetoro L, Carroli G. Side effects of oral misoprostol during the first 24 hours after administration in the third stage of labour. *BJOG* 2002;109:1222-6.
- 13 Bamigboye AA, Hofmeyr GJ, Merrell DA. Rectal misoprostol in the prevention of postpartum hemorrhage: a placebo-controlled trial. *Am J Obstet Gynecol* 1998;179:1043-6.
- 14 Hofmeyr GJ, Nikodem VC, de Jager M, Gelbart BR. A randomised placebo controlled trial of oral misoprostol in the third stage of labour. *Br J Obstet Gynaecol* 1998;105:971-5.
- 15 Surbek DV, Fehr PM, Hosli I, Holzgreve W. Oral misoprostol for third stage of labor: a randomized placebo-controlled trial. *Obstet Gynecol* 1999;94:255-8.
- 16 Prasertcharoensuk W, Swadpanich U, Lumbiganon P. Accuracy of the blood loss estimation in the third stage of labor. *Int J Gynaecol Obstet* 2000;71:69-70.
- 17 Hoj L, da Silva D, Hedegaard K, Sandstrom A, Aaby P. Factors associated with maternal mortality in rural Guinea-Bissau. A longitudinal population-based study. *BJOG* 2002;109:792-9.

Cluster randomised trial of intermittent preventive treatment for malaria in infants in area of high, seasonal transmission in Ghana

Daniel Chandramohan, Seth Owusu-Agyei, Ilona Carneiro, Timothy Awine, Kwame Amponsa-Achiano, Nathan Mensah, Shabbar Jaffar, Rita Baiden, Abraham Hodgson, Fred Binka, Brian Greenwood

Abstract

Objective To evaluate the effects of intermittent preventive treatment for malaria in infants (IPTi) with sulfadoxine-pyrimethamine in an area of intense, seasonal transmission.

Design Cluster randomised placebo controlled trial, with 96 clusters allocated randomly to sulfadoxine-pyrimethamine or placebo in blocks of eight.

Interventions Children received sulfadoxine-pyrimethamine or placebo and one month of iron supplementation when they received DPT-2, DPT-3, or measles vaccinations and at 12 months of age.

Main outcome measures Incidence of malaria and of anaemia determined through passive case detection.

Results 89% (1103/1242) of children in the placebo group and 88% (1088/1243) in the IPTi group completed follow-up to 24 months of age. The protective efficacy of IPTi against all episodes of malaria was 24.8% (95% confidence interval 14.3% to 34.0%) up to 15 months of age. IPTi had no protective effect against malaria between 16 and 24 months of age (protective efficacy -4.9%, -21.3% to 9.3%). The incidence of high parasite density malaria (≥ 5000 parasites/ μ l) was higher in the IPTi group than in the placebo group between 16 and 24 months of age (protective efficacy -19.5%, -39.8% to -2.2%). IPTi reduced hospital admissions with anaemia by 35.1% (10.5% to 52.9%) up to 15 months of age. IPTi had no significant effect on anaemia between 16 and 24 months of age (protective efficacy -6.4%, -76.8% to 35.9%). The relative risk of death up to 15 months of age in the IPTi group was 1.26 (95% confidence interval 0.81 to 1.96; $P=0.31$), and from 16 to 24 months it was 1.28 (0.77 to 2.14; $P=0.35$).

Conclusions Intermittent preventive treatment for malaria with sulfadoxine-pyrimethamine can reduce malaria and anaemia in infants even in seasonal, high transmission areas, but concern exists about possible

rebound in the incidence of malaria in the second year of life.

Introduction

Malaria remains the leading cause of morbidity and mortality in young children in impoverished communities in Africa. It is estimated that 515 (range 300-600) million clinical attacks of falciparum malaria occurred globally in 2002 and that 70% of these cases occurred in Africa.¹ Malaria is one of the major causes of anaemia in children in sub-Saharan Africa, accounting for an estimated 18% of the disability adjusted life years lost because of anaemia.² The number of deaths attributable to malaria associated anaemia is estimated at 190 000 to 974 000 a year.³ In high intensity transmission areas, the highest burden of malaria and malaria associated anaemia is in infants.⁴

Control of malaria and anaemia depends largely on passive case detection and appropriate treatment. However, many children who have malaria parasitaemia, anaemia, or both remain asymptomatic,⁵ and chronic asymptomatic parasitaemia increases the risk of severe anaemia.⁶ Chemoprophylaxis during the first year of life significantly reduced the incidence of malaria and anaemia in a study in Tanzania.⁷ The risk of malaria and anaemia increased in the second year of life after chemoprophylaxis had been stopped, however, suggesting that complete protection against malaria infection during infancy had impaired naturally acquired immunity against the infection. Two recent studies in Tanzania, done in areas with perennial transmission of malaria, showed that administration of a full therapeutic course of an antimalarial at set times during infancy, regardless of whether or not the infant was parasitaemic (intermittent preventive treatment in infants (IPTi)), substantially reduced the incidence of malaria and anaemia.⁸⁻⁹ The antimalarial was given in

London School of Hygiene and Tropical Medicine, London WC1E 7HT

Daniel Chandramohan
clinical senior lecturer
Ilona Carneiro
lecturer
Shabbar Jaffar
senior lecturer
Brian Greenwood
director, Gates
Malaria Partnership

Kintampo Health Research Centre, Kintampo, Ghana
Seth Owusu-Agyei
director

Navrongo Health Research Centre, Navrongo, Ghana

Timothy Awine
research officer

Kwame Amponsa-Achiano
research physician

Nathan Mensah
data manager

Rita Baiden
research physician

Abraham Hodgson
director

INDEPTH-Network, Accra, Ghana
Fred Binka
executive director

Correspondence to:
D Chandramohan
daniel.chandramohan@lshtm.ac.uk

BMJ 2005;331:727-33



A figure and three tables are on bmj.com