

Evidence for risk of bias in cluster randomised trials: review of recent trials published in three general medical journals

Suezann Puffer, David J Torgerson, Judith Watson



This is an abridged version; the full version is on bmj.com

Abstract

Objective To examine the prevalence of a risk of bias associated with the design and conduct of cluster randomised controlled trials among a sample of recently published studies.

Design Retrospective review of cluster randomised trials published in the *BMJ*, *Lancet*, and *New England Journal of Medicine* from January 1997 to October 2002.

Main outcome measures Prevalence of secure randomisation of clusters, identification of participants before randomisation (to avoid foreknowledge of allocation), differential recruitment between treatment arms, differential application of inclusion and exclusion criteria, and differential attrition.

Results Of the 36 trials identified, 24 were published in the *BMJ*, 11 in the *Lancet*, and a single trial in the *New England Journal of Medicine*. At the cluster level, 15 (42%) trials provided evidence for secure allocation and 25 (69%) used stratified allocation. Few trials showed evidence of imbalance at the cluster level. However, some evidence of susceptibility to risk of bias at the individual level existed in 14 (39%) studies.

Conclusions Some recently published cluster randomised trials may not have taken adequate precautions to guard against threats to the internal validity of their design.

Introduction

In most clinical trials participants are randomised as individuals to different treatments. Sometimes individual allocation is not possible or desirable, and groups of individuals are randomised instead: this is known as cluster or group randomisation. Many reasons for using cluster allocation exist. For example, evaluation of clinical guidelines or medical education on patient outcomes almost always requires that healthcare professionals are the “unit” of allocation.

Although randomised trials are the most robust evaluative method, poorly conducted studies are susceptible to different forms of selection bias that can make their results unsound. Methodological reviews of individually randomised trials have shown that rigorously conducted trials produce different effect estimates from poorly conducted studies.^{1,2} Less attention has been paid, however, to cluster trials. Cluster trials are generally more difficult to design and execute than indi-

vidually randomised studies, and some design features of a cluster trial may make it especially vulnerable to a range of threats that can introduce selection bias.

In cluster trials potential bias in the execution of the trial can occur at two levels, the first of which is the cluster level. Randomisation of clusters needs to be undertaken carefully and preferably independently. Otherwise, biased allocation may occur (certain clusters being allocated to a particular arm on the basis of reasons that might affect outcome). It is theoretically possible for allocation of clusters to be subverted, as has happened in individually randomised trials.³ Similarly, once clusters have been allocated it is important, as with individually randomised trials, to try to retain the cluster in its allocated group and avoid the cluster dropping out to avoid the risk of attrition bias.

The second level at which bias can occur in cluster trials is after the clusters have been allocated and when individual participants are recruited into the study. For example, in a trial of the effectiveness of a training package for general practitioners, patients had to be identified prospectively after the general practitioner had been randomised.⁴ The prospective inclusion of participants can potentially lead to selection bias through the recruitment of different types of participant by the researcher or clinician. If the person prospectively recruiting participants has “foreknowledge” of the allocation group then, as shown in individually randomised trials, bias can result.³ In addition to this source of selection bias, another can be introduced by the participant if consent is needed after randomisation. Selection bias can be introduced if consent is withheld for either treatment or data collection.

Several ways of avoiding the biases outlined above exist. One is to try to identify trial participants before randomisation and obtain consent for treatment, data collection, or both before allocation. Use of prior identification and prior consent avoids potential biases occurring through foreknowledge of the allocation schedule, by the researcher and patient. If this is not possible, identification and recruitment of participants should ideally be undertaken by someone blinded to the group allocation.

Another problem that can lead to bias, in both individual and cluster randomised trials, is the differential application of inclusion and exclusion criteria. Again this problem can be reduced if the person applying the criteria is masked to the group allocation.

York Trials Unit,
Department of
Health Sciences,
York University,
York YO10 5DD
Suezann Puffer
research assistant
David Torgerson
director
Judith Watson
research fellow

Correspondence to:
D J Torgerson
djt6@york.ac.uk

BMJ 2003;327:785–7



Additional references appear on bmj.com

In this paper we review some recently published cluster trials to determine the extent of their risk of bias. We also describe the steps that some authors took to reduce these risks.

Methods

We hand searched the *BMJ*, *Lancet*, and *New England Journal of Medicine* for all cluster randomised trials published from January 1997 to October 2002. Selection bias can be introduced into a trial in several different ways. In this paper we sought evidence for the risk of bias from several sources.

Secure cluster allocation—This is where evidence exists that cluster randomisation was securely undertaken.

Cluster attrition—This occurs when clusters are lost to follow up after randomisation.

Cluster imbalance—This is where evidence exists of imbalance in important variables at the cluster level.

Differential individual recruitment or consent—This is when different proportions of participants are recruited to the different arms of the trial. If recruitment rates differ between groups this may lead to the risk of bias.

Differential individual exclusion or inclusion—This can occur when eligibility criteria are applied differentially after randomisation, which can introduce bias.

Results

We included 36 trials, 24 of which were published in the *BMJ*, 11 in the *Lancet*, and one in the *New England Journal of Medicine*. In table 1 we describe the basic characteristics of the trials.

Secure cluster allocation—Fifteen trials seemed to use a secure method of allocating clusters; the remainder did not clearly describe who undertook the allocation or how this was done. Most trials used some form of stratified random allocation to reduce the possibility of “chance bias.”

Cluster attrition—In 10 trials a loss of clusters occurred between randomisation and follow up. Most of the trials lost only a small proportion of their clusters, but one study lost more than half (56%) of all the randomised clusters.^{w2}

Differential consent or recruitment—We found some evidence for differential consent or recruitment in seven of the 23 trials that had not undertaken prior identification of participants. Three trials recruited more participants from one group than the other,^{w11 w16 w31} and the other four studies differentially obtained consent from more participants in one arm than the other.^{w9 w13 w29}

^{w34} One trial, although it seemed to identify all the participants before allocation for the main mortality outcome, seemed to have introduced the risk of

Table 1 Characteristics of included cluster trials

Study	No of clusters	No of participants	Description	Clustering accounted for in sample size estimation?
Aveyard 1999 ^{w3}	52	9 301	Expert system for smoking prevention and cessation in schools	Yes
Bennewith 2002 ^{w4}	98	2 141	Prevention of repeat episodes of deliberate self harm	Yes
Carman 2000 ^{w1}	20	1 437	Influenza vaccination of healthcare workers	Yes
Chapman 2000 ^{w5}	8	346	Educational intervention to prevent dog bites	No
Field 2001 ^{w6}	9	494	Two methods of data collection	Yes
Flottorp 2002 ^{w7}	142	12 369	To improve general practice management of sore throat and urinary tract infections	Yes
Gavvani 2002 ^{w8}	18	4 498	Insecticide impregnated dog collars on incidence of zoonotic visceral leishmaniasis	No
Graham 2002 ^{w9}	24	3 794	Teenagers' knowledge of emergency contraception	Yes
Haider 2000 ^{w10}	40	726	Community based peer counsellors on breast feeding	Not clear
Jolly 1999 ^{w11}	67	686	Programme to coordinate and support follow up care in general practice	Not clear
Jordhoy 2000 ^{w12}	6	707	Palliative care intervention	Not clear
Kannus 2000 ^{w13}	22	1 725	Hip protectors	No
Kendrick 1999 ^{w14}	36	2 152	Prevent unintentional injuries in children	Yes
Kidane 2000 ^{w15}	37	70 506	Maternal education for early treatment of paediatric malaria	Yes
King 2002 ^{w2}	116	410	Behavioural therapy to treat patients with depression	Yes
Kinmonth 1998 ^{w16}	43	360	Patient centred care for diabetes in general practice	Yes
Kroeger 2002 ^{w17}	14	2 913	Insecticide impregnated curtains to control transmission of cutaneous leishmaniasis	No
MacArthur 2002 ^{w18}	37	3 580	Community postnatal care	Yes
McCartney 1997 ^{w19}	28	182 200	General practitioner feedback to increase aspirin use	No
Moher 2001 ^{w20}	21	2 142	Secondary prevention of coronary heart disease	Yes
Montgomery 2000 ^{w21}	27	810	Interventions for management of hypertension	Yes
Morrison 2001 ^{w22}	221	689	Infertility guidelines for general practitioners	Yes
Morrow 1999 ^{w23}	39	130	Home based counselling to promote breast feeding	Yes
O' Cathain 2002 ^{w24}	13	10 327	Leaflets to promote informed choice in maternity care	Yes
Olivarius 2001 ^{w25}	311	1 470	Structured personal care of type 2 diabetes mellitus	No
Premaratne 1999 ^{w26}	40	48 800	Effectiveness of an asthma resource centre	No
Sagliocca 1999 ^{w27}	146	404	Hepatitis A vaccine	No
Sahota 2001 ^{w28}	10	636	School intervention to reduce risk factors for obesity	Yes
Shah 2001 ^{w29}	6	325	Peer led programme for asthma education in adolescents	No
Smeeth 2001 ^{w30}	106	42 278	Methods to administer a screening questionnaire	No
Stepoe 1999 ^{w31}	20	883	Behavioural counselling in general practice	Yes
Thompson 2000 ^{w32}	59	4 192	Detection and outcome of depression in primary care	Yes
Van Eijk 2001 ^{w33}	21	46 078	Academic detailing to reduce antidepressant use	No
Wawer 1999 ^{w34}	10	44 107	Prevention of sexually transmitted disease	No
West 1999 ^{w35}	270	44 646	Supplementation with vitamin A or β carotene on mortality related to pregnancy	No
Wight 2000 ^{w36}	25	8 430	Teacher delivered sex education	Yes

Table 2 Evidence of risk of bias

Study	Potential source of bias	Acknowledgment of risk by authors and steps taken
Carmen 2000 ^{w1}	48% v 33% of patients having influenza vaccination, P<0.01; 69% (258/375) v 78% (269/344) accepted virological screening for secondary outcome assessment, P=0.004	Yes, for cluster imbalance, used adjusted odds ratios
Gavagni 2002 ^{w9}	7.6% (143/1870) v 11.4% (229/2006) attrition in control and intervention groups, P<0.001	No
Graham 2002 ^{w9}	12.2% (216/1768) of control group refused consent v 17% (344/2026) of intervention group, P<0.001	No
Jolly 1999 ^{w11}	Control group recruited 15.5% more than intervention group; practice population not given, so impossible to see if significantly different	Noted recruitment "imbalance" in discussion
Kannus 2000 ^{w13}	31.4% (204/650) of intervention group refused consent v 8.7% (94/1075) of control group, P<0.001	Acknowledged potential for selection bias in discussion
Kinmonth 1998 ^{w16}	0.06% (142/225 015) v 0.047% (108/230 560) of practice populations were recruited for intervention and control groups, P=0.02	No
MacArthur 2002 ^{w18}	4.2% (46/1087) v 2.6% (25/977) of intervention and control groups withdrew or moved away, P=0.04	Commented on loss to follow up in discussion
Moher 2001 ^{w20}	0.32% (2/623) v 2.6% (20/772) and 1.3% (10/747) misdiagnoses for control and two intervention groups, P=0.002	No
Olivarius 2001 ^{w25}	8.7% (67/774) v 4.0% (28/696) excluded owing to illness in intervention group and control group, P<0.01	Noted in results more post-randomisation exclusions
Sagliocca 1999 ^{w27}	3.9% (7/178) v 0% (0/173) lost to follow up in control and intervention groups, P=0.02	No
Shah 2001 ^{w29}	22.3% (148/662) v 17.3% (124/717) included in control and intervention groups, P=0.02	No
Step toe 1999 ^{w31}	Control practices recruited 567 v 316 participants given similar sized populations (P value not calculable)	Differential recruitment rate mentioned in discussion, but not as a potential source of bias
Wawer 1999 ^{w34}	17.5% (4002/22 915) v 14.7% (3125/21 192) refused consent or treatment in intervention and control groups, P<0.001	No
Wight 2002 ^{w36}	30.6% (1070/3493) v 27.5% (1069/3892) attrition in numbers reporting intercourse in intervention and control groups, P=0.003	No

selection bias into the measurement of its secondary outcome.^{w1}

Differential application of inclusion or exclusion criteria—We found two trials that seemed to have applied inclusion or exclusion criteria differentially between groups after randomisation. The authors of a study promoting methods of secondary prevention of coronary disease excluded significantly more participants owing to misdiagnosis in the intervention groups than in the control group.^{w20} Similarly, in another trial twice as many participants were excluded from the intervention group because of illness than from the control arm.^{w25}

Differential attrition—Evidence of differential attrition between the randomised groups existed in four trials.^{w8 w18 w27 w36}

Table 2 summarises the potential sources of bias risk in 14 trials in which we observed differences between the groups that indicate a risk of selection bias. Authors of six studies alerted the reader to the potential risk of bias in their study.

Discussion

Elbourne and Campbell have recently argued for amending the CONSORT statement to allow for the special methodological circumstances of cluster trials.⁴ We would echo this call. We found it very difficult in several of the trials to ascertain whether a risk of bias was likely or not. We would wish the following additions to be made. Firstly, a clear statement as to whether the population was identified before or after the allocation decision had been made. Secondly, was the person who recruited the participants blind to group allocation? Thirdly, what was the size of the population within the clusters? For example, Steptoe et al^{w31} did not state the size of the general practice populations in their trial arms and Kinmonth et al^{w15} gave only means. For the first of these studies we could only assume that the recruitment was significantly different, and for the second study we had to make an estimate. This missing information also meant that for some studies we could not be completely sure if

What is already known on this topic

Reviews of individually randomised trials show that results can differ according to quality of methods

Foreknowledge of allocation and failure to use intention to treat analysis can lead to bias

What this study adds

Cluster randomised trials are susceptible to forms of selection bias

Careful planning and execution of such trials can avoid these biases

recruitment bias had taken place. For example, in Kendrick et al no suggestion of any recruitment bias was apparent; however, we could not be absolutely sure as the authors did not present the practice population sizes.^{w14}

Conclusion

Cluster trials are vulnerable to the risk of bias. Careful planning and execution of such trials can avoid these biases.

We thank Doug Altman, Diana Elbourne, Craig Ramsay, and Trevor Sheldon for their helpful comments on an earlier version of this manuscript.

Contributors: See bmj.com

Funding: JW is trial coordinator of the SAPHIRE trial funded by the Medical Research Council; SP and DJT are funded by the University of York.

Competing interests: None declared.

Ethical approval: Not needed.

- Shulz KF, Chalmers I, Grimes DA, Altman DG. Assessing the quality of randomization from reports of controlled trials in obstetrics and gynaecology journals. *JAMA* 1994;272:125-8.
- Kjaergaard LL, Villumsen J, Cluud C. Reported methodologic quality and discrepancies between large and small randomized trials in meta-analyses. *Ann Intern Med* 2001;135:982-9.
- Schulz KF. Subverting randomisation in controlled trials. *JAMA* 1995;274:1456-8.
- Elbourne DR, Campbell MK. Extending the CONSORT statement to cluster randomized trials: for discussion. *Stat Med* 2001;20:489-96. (Accepted 10 July 2003)