



The full version of this article appears on *bmj.com*

## Randomised controlled trial of clinical medication review by a pharmacist of elderly patients receiving repeat prescriptions in general practice

Arnold G Zermansky, Duncan R Petty, David K Raynor, Nick Freemantle, Andy Vail, Catherine J Lowe

Division of Academic Pharmacy Practice, School of Healthcare Studies, University of Leeds, Leeds LS2 9UT

Arnold G Zermansky  
*honorary senior research fellow*

Duncan R Petty  
*research pharmacist*  
David K Raynor  
*professor of pharmacy practice, medicines, and their users*  
Catherine J Lowe  
*lecturer*

Department of Primary Care and General Practice, University of Birmingham, Birmingham B15 2TT

Nick Freemantle  
*professor of clinical epidemiology and biostatistics*

Biostatistics Group, University of Manchester, Hope Hospital, Salford, Manchester M6 8HD  
Andy Vail  
*lecturer*

Correspondence to: A G Zermansky  
arnoldz@easynet.co.uk

*BMJ* 2001;323:1340-3

### Abstract

**Objective** To determine whether a pharmacist can effectively review repeat prescriptions through consultations with elderly patients in general practice.

**Design** Randomised controlled trial comparing clinical medication review by a pharmacist against normal general practice review.

**Setting** Four general practices.

**Participants** 1188 patients aged 65 or over who were receiving at least one repeat prescription and living in the community.

**Intervention** Patients were invited to a consultation at which the pharmacist reviewed their medical conditions and current treatment.

**Main outcome measures** Number of changes to repeat prescriptions over one year, drug costs, and use of healthcare services.

**Results** 590 (97%) patients in the intervention group were reviewed compared with 233 (44%) in the control group. Patients seen by the pharmacist were more likely to have changes made to their repeat prescriptions (mean number of changes per patient 2.2 *v* 1.9; difference = 0.31, 95% confidence interval 0.06 to 0.57; *P* = 0.02). Monthly drug costs rose in both groups over the year, but the rise was less in the intervention group (mean difference £4.72 per 28 days, -£7.04 to -£2.41); equivalent to £61 per patient a year. Intervention patients had a smaller rise in the number of drugs prescribed (0.2 *v* 0.4; mean difference -0.2, -0.4 to -0.1). There was no evidence that review of treatment by the pharmacist affected practice consultation rates, outpatient consultations, hospital admissions, or death rate.

**Conclusions** A clinical pharmacist can conduct effective consultations with elderly patients in general practice to review their drugs. Such review results in significant changes in patients' drugs and saves more than the cost of the intervention without affecting the workload of general practitioners.

### Introduction

Over 80% of drugs prescribed by general practitioners in the United Kingdom are repeat prescriptions—that is, they are re-prescribed without a consultation between the doctor and the patient.<sup>1</sup> Repeat prescribing

is poorly managed in the United Kingdom.<sup>2</sup> In 1994, the Audit Commission suggested that the review of long term treatment might be inadequate.<sup>3</sup> Zermansky subsequently found that 72% of repeat prescriptions sampled in 50 practices had not been reviewed in the past 15 months.<sup>2</sup> He concluded that this is potentially both wasteful and dangerous. Purves and Kennedy expressed concern about the variation in the quality of review between practices.<sup>4</sup>

The Royal College of Physicians and the recent National Service Framework for Older People emphasise the need for regular review of treatment for elderly patients.<sup>5-6</sup> In view of the increasing workload of general practitioners, it has been proposed that pharmacists should review patients. Several North American trials have shown the benefits of pharmacists reviewing long term prescriptions in community practice.<sup>7-11</sup> In the United Kingdom, two limited randomised controlled trials suggest that review of treatment by pharmacists identifies more drug related problems than normal care.<sup>12-13</sup> We tested whether pharmacists can effectively review the conditions and treatments of elderly patients in consultation with the patient.

### Participants and methods


#### Selection criteria and randomisation

We recruited general practices by randomly selecting them from a list of all practices in Leeds Health Authority with four or more partners, computerised repeat prescribing, no previous or current clinical pharmacist involvement, and prescribing costs close to average.

The participating practices provided complete lists of registered patients aged 65 and over who were receiving at least one drug on repeat prescription on 1 June 1999. We excluded patients in nursing or residential homes, those who had a terminal illness, and those who were in clinical trials. We wrote to eligible patients asking them to participate. Those who consented were randomised to an intervention group (clinical medication review by pharmacist) or control group (normal care) by computer generated random numbers.

#### Intervention

The pharmacist (DRP) invited patients to his clinic when their next review was due. Patients with no review date were invited to attend when convenient. Immobile



Process for reviewing repeat prescriptions

patients were visited at home. Non-attenders were invited once more by telephone.

The intervention has been described previously and is summarised as an algorithm (figure).<sup>4</sup> During the consultation with the patient, the pharmacist discussed each condition being treated and asked about relevant symptoms (such as swollen ankles and breathlessness in patients with heart failure). In conditions for which clinical or pathological monitoring was due, the pharmacist directed the patient to the practice nurse or doctor. The pharmacist did not physically examine the patient, although he noted signs that were obvious during the consultation, such as swollen ankles or rash. Patients with new clinical problems were referred to the doctor.

Treatment recommendations were based on national, local, and (where available) practice guidelines. We agreed with each practice the level of intervention that the pharmacist could make without seeking prior approval.

### Usual care

Patients in the control group continued to receive normal care from their general practitioner and primary healthcare staff. Patients were recalled for review of treatment by the general practitioner according to normal custom in the practice.

### Outcome measures

The primary outcome measure was the number of changes to repeat prescriptions between baseline (June 1999) and the end of the 12 month study (June 2000). The secondary outcome measures were changes in number and cost of medicines and frequency of dose and effect on healthcare workload (general practitioner consultations, hospital outpatient attendances, and acute admissions).

### Clinical medication review

Clinical medication review is the process where a health professional reviews the patient, the illness, and the drug treatment during a consultation. It involves evaluating the therapeutic efficacy of each drug and the progress of the conditions being treated. Other issues, such as compliance, actual and potential adverse effects, interactions, and the patient's understanding of the condition and its treatment are considered when appropriate. The outcome of the review will be a decision about the continuation (or otherwise) of the treatment.

### Collection of data

As well as the age and sex of patients, we recorded number of repeat prescriptions, number of times doses were taken a day, and net ingredient cost of 28 days' supply (based on Drug Tariff and *Monthly Index of Medical Specialties* for December 1998) at baseline and the end of the study. The number of consultations within the practice, outpatient attendances, and acute admissions were recorded for the duration of the study. We recorded drop out due to death, leaving the practice, or going into a residential home. We also collected data for six months before the intervention to allow us to test whether the pharmacist's presence contaminated the control group.

### Results

The four practices had a total list size of 28 202 (individual sizes 6342, 7647, 8759, and 5454) with 3308 patients aged 65 and over. We contacted 2403 patients consecutively until the required number of participants was obtained; 1188 consented and were randomised to either intervention or control groups. At the end of the study, data on 1131 patients (581 intervention and 550 control) were available for analysis. There was even distribution of age, sex, practice, and number of drugs on repeat prescription between the intervention and control groups, and there were no differences in drug costs or number of doses at baseline.

### Outcomes

The mean number of changes per patient was 2.2 in the intervention group and 1.9 in the control group (difference = 0.31, 95% confidence interval 0.06 to 0.57;  $P = 0.02$ ). Table 1 shows the numbers of patients who had at least one change to their treatment during the study. More patients in the control group than the intervention group started taking a new drug. There was no clear difference in the number of other changes.

**Table 1** Numbers (percentages) of patients whose repeat prescriptions were changed during the study. Some patients had more than one change

Type of change	Intervention (n=581)	Control (n=550)	Total (n=1131)
New drug started	265 (46)	270 (49)	535 (47)
Drug stopped	239 (41)	180 (33)	419 (37)
Switched drug	119 (20)	93 (17)	212 (19)
Dose changed	98 (17)	61 (11)	159 (14)
Change to generic	64 (11)	37 (7)	101 (9)
Formulation changed	17 (3)	12 (2)	29 (3)
Frequency changed	6 (1)	0	6 (1)
Any of above	438 (75)	397 (72)	835 (74)

**Table 2** Changes in treatment between start and finish of study

	Intervention			Control			Difference between groups (95%CI)	P value*
	Start (n=596)	Finish (n=576)	Change	Start (n=577)	Finish (n=549)	Change		
Mean No of repeat medicines	4.8	5.0	0.2	4.6	5.0	0.4	-0.2 (-0.4 to -0.1)	0.01
Mean cost over 28 days (£)	29.27	31.07	1.80	28.23	34.85	6.52	-4.72 (-7.04 to -2.41)	0.0001
Mean No of dose times/day	2	1.9	-0.1	2.1	1.9	-0.2	0.1 (0.0 to 0.2)	0.17

\*t test.

Table 2 shows the differences between baseline and follow up in the numbers, costs, and doses of repeat prescriptions. Numbers of drugs and cost rose in both groups, but for each the rise was significantly less in the intervention group. The number of daily doses did not differ significantly. There was no evidence of any adverse health outcome in the intervention group as measured by need for consultation with a general practitioner or hospital treatment. The number of deaths was 15 (2.5%) in the intervention group and 25 (4.3%) in the control group (odds ratio = 0.56, 0.29 to 1.1).

In all, 590 (97%) intervention patients had a consultation with the pharmacist (one was seen twice). In the control group, 233 (44%) patients had a documented review with a doctor. The pharmacist took an average of 20 minutes to conduct a review (excluding collection of research data). The gross cost of the pharmacist was £21 per hour, or £7 per patient reviewed. The average reduction in net cost of drugs per patient per 28 days was £4.72 (£2.41 to £7.04)

## Discussion

We have shown that a trained pharmacist can conduct clinical medication review of elderly patients in the general practice setting. The pharmacist's review resulted in more changes to treatment than normal care and produced an important cost saving, even after the cost of the intervention was deducted.

### Validity

We recruited half of contacted patients. There was concern that the participants might not be typical of the practices' eligible elderly populations. We have shown previously that the participants tended to be younger, male, and taking fewer drugs than non-participants.<sup>15</sup> This suggests that our results may underestimate the effects of the review. Patients taking more drugs are more likely to benefit from the pharmacist's intervention, provided that they can be persuaded to attend a review. Attendance would be more likely in the context of care rather than a clinical trial.

The unit of randomisation was the patient. Thus practices contained both intervention and control patients. We collected data for the six months before the study started in response to concern that contamination could occur as a result of the pharmacist's presence in the practice. Comparison of these data with study data showed no evidence of contamination.

### Reasons for difference between groups

The smaller increase in the mean number of repeat prescriptions in the intervention group was mainly due to these patients being more likely to have drugs stopped. Intervention patients had more changes to treatment in general, perhaps because the pharmacist

did a more detailed review than the general practitioners. This effect could be important because patients' compliance has been shown to decrease with increasing number of drugs.<sup>16</sup> Stopping unnecessary drugs may also reduce the risk of adverse effects and interactions.

Review of drug treatment by pharmacists could have increased general practice consultation rates if patients made appointments to confirm advice given by the pharmacist, to have tests done, or to have treatment recommendations implemented. Consultations did increase immediately after the review, but the total number in the year was not different from that in the control group. The increase in consultations was due to patients requiring tests (usually referred to the practice nurse) or to suspected worsening of an existing condition or a new condition (referral to the general practitioner). The extra workload was therefore appropriate and was balanced by a reduced workload in the subsequent months.

We adopted a clinical patient centred approach rather than relying on technical appraisal of the drugs, as in some other studies.<sup>12 17</sup> This resulted in more clinical interventions such as ensuring treatment was monitored, identifying new health problems, suggesting new interventions, and reinforcing compliance. Our study supports the concept of medication review suggested in the National Service Framework for Older People.<sup>6</sup>

The small scale of this trial, involving only four practices in one city and just one pharmacist, limits the generalisability of the results. Nevertheless, it shows that significant and clinically important results can be achieved by pharmacists reviewing patients and their

### What is already known on this topic

Review of patients on long term drug treatment is important but is done inadequately

Evidence from the United States shows that pharmacists can improve patient care by reviewing drug treatment

### What this study adds

Consultations with a clinical pharmacist are an effective method of reviewing the drug treatment of older patients

Review by a pharmacist results in more drug changes and lower prescribing costs than normal care, plus a much higher review rate

Use of healthcare services by patients is not increased

treatment. A larger scale study with more practices and pharmacists is needed to clarify the practicality, costs, and benefits.

We thank Dr Dowson and partners, Dr Geraghty and partners, Dr Clements and partners, Dr Burkill and partners, and the patients for participating and D Buttress for help in recruiting patients and secretarial support. The views and opinions expressed do not necessarily reflect those of the funding body.

Contributors: see bmj.com

Funding: NHS Research and Development National Coordinating Centre for Health Technology Assessment.

Competing interests: None declared.

- Harris CM, Dajda R. The scale of repeat prescribing. *Br J Gen Pract* 1996;46:649-53.
- Zermansky AG. Who controls repeats? *Br J Gen Pract* 1996;46:643-7.
- Audit Commission. *A prescription for improvement. Towards more rational prescribing in general practice*. London: HMSO, 1994.
- Purves I, Kennedy J. *The quality of general practice repeat prescribing*. Newcastle: University of Newcastle upon Tyne, 1994.
- Royal College of Physicians. *Medication for older people*. 2nd ed. London: RCP, 1997.
- Department of Health. *Medicines and older people: implementing medicines related aspects of the National Service Framework for Older People*. London: DoH, 2001.
- Hanlon JT, Weinberger M, Samsa GP, Schmadler KE, Uttech KM, Lewis IK, et al. A randomised, controlled trial of a clinical pharmacist interven-

tion to improve inappropriate prescribing in elderly outpatients with polypharmacy. *Am J Med* 1996;100:428-37.

- Cassidy IB, Keith MR, Coffey EL, Noyes A. Impact of pharmacist-operated general medicine chronic care refill clinics on practitioner time and quality of care. *Ann Pharmacother* 1996;30:745-51.
- Borgsdorf LR, Miano JS, Knapp KK. Pharmacist-managed medication review in a managed care system. *Am J Hosp Pharm* 1994;51:772-7.
- Deady JE, Lepinski PW, Abramowitz PW. Measuring the ability of clinical pharmacists to effect drug therapy changes in a family practice clinic using prognostic indicators. *Hospital Pharmacy* 1991;26:93-7.
- Lobas NH, Lepinski PW, Abramowitz PW. Effects of pharmaceutical care on medication costs and quality of patient care in an ambulatory-care clinic. *Am J Hosp Pharm* 1992;49:1681-8.
- Granas AG, Bates I. The effect of pharmaceutical review of repeat prescriptions in general practice. *Int J Pharm Pract* 1999;7:264-75.
- Mackie CA, Lawson DH, Campbell A, Maclaren AG, Waigh R. A randomised controlled trial of medication review in patients receiving polypharmacy in general practice. *Pharm J* 1999;263:R7.
- Lowe CJ, Petty DR, Zermansky AG, Raynor DK. Development of a method for clinical medication review by a pharmacist in general practice. *Pharm World Sci* 2000;22(4):121-6.
- Petty DR, Zermansky AG, Raynor DK, Vail A, Lowe CJ, Freemantle N, et al. "No thank you": Why elderly patients declined to participate in a research study. *Pharm World Sci* 2001;23(1):22-7.
- Murray MD, Birt JA, Manatunga AK. Medication compliance in elderly outpatients using twice-daily dosing and unit-of-use packaging. *Ann Pharmacother* 1993;27:616-21.
- Goldstein R, Hulme H, Willits J. Reviewing repeat prescribing—general practitioners and community pharmacists working together. *Int J Pharm Pract* 1998;6:60-6.

(Accepted 12 October 2001)

### *A memorable patient* When guidelines fail

She was a 78 year old widow who lived in sheltered accommodation and, despite being partially sighted because of advancing glaucoma, was still able to get about with her white cane. She rarely visited our surgery, and when she did it was for a flu jab or regarding moderate generalised osteoarthritis. This morning, however, she limped into my consulting room apologetically and said, "The heel of my right foot is hurting, doctor. I'm finding it difficult to walk to the shops. My eyesight doesn't help either."

I set out to examine her foot. There was nothing grossly abnormal. It was well perfused, no swellings, and no evidence of any trauma. Deep palpation over the calcaneum produced tenderness. "Bingo," I thought, "plantar fasciitis, with or without calcaneal spur." I reached for an x ray form when I suddenly remembered the fate of the previous x ray form I had sent for an almost identical situation. A consultant radiologist from one of our teaching hospitals had politely returned my request form with a booklet containing guidelines from the Royal College of Radiologists. This had stated that routine radiography of the heel for calcaneal spur is of no value and not recommended. I didn't want my patient to travel all the way to the radiology department only to be sent back with another booklet of guidelines. I reassured her that this was a self limiting condition, prescribed some local rubefacient, and told her to wear a heel pad and to come back if there was no improvement.

Two weeks later, she asked for a home visit as the pain was no better. I visited her after the morning surgery rather grudgingly—a request for a home visit for such a minor condition? Examination did not reveal anything new, and I discussed the option of a local corticosteroid injection. She gladly accepted, and I arranged for her to visit the surgery in two days time and included her in my minor surgery list.

A carer brought her in, and her limp was more pronounced. I carried out the procedure and reassured her that she would soon improve.

The following week an irate son in law, who I had never met, telephoned the surgery asking that a doctor

visit straight away as his mother in law was no better and in a lot of pain. The receptionist told him that the doctors were in the middle of a busy surgery and would visit around midday. An hour later he rang again to say that he was not prepared to wait and was taking his mother in law to the casualty department.

Ten days later I received a call from the hospital. My patient was to be discharged to her daughter's home address for terminal care. Widespread metastasis, including involvement of her right calcaneum, had been diagnosed, and no attempt was made to identify the primary. Completely shocked, I replaced the telephone receiver. It took several minutes to compose myself and continue the rest of the surgery. She rapidly deteriorated and died a few days later.

I have never heard of secondary deposits in the calcaneum. Would x ray examination have altered her prognosis? Most certainly not, but it would have spared her a painful injection. Pain management and other aspects of care would have been different.

This made me think about guidelines, of which there are so many. When it comes to an individual patient, you have to rely on your clinical sense and skills and not fall prey to generalised guidelines and pathways.

The question that I still cannot answer is how do I persuade a colleague in another speciality, who perhaps has not seen the patient, to overlook guidelines set out by his or her royal college and perform a simple investigation?

John Godfrey *general practitioner, Liverpool*

We welcome articles of up to 600 words on topics such as *A memorable patient*, *A paper that changed my practice*, *My most unfortunate mistake*, or any other piece conveying instruction, pathos, or humour. If possible the article should be supplied on a disk. Permission is needed from the patient or a relative if an identifiable patient is referred to. We also welcome contributions for "Endpieces," consisting of quotations of up to 80 words (but most are considerably shorter) from any source, ancient or modern, which have appealed to the reader.