



This is an abridged version; the full version is on bmj.com

Prevalence of left ventricular systolic dysfunction and heart failure in high risk patients: community based epidemiological study

R C Davis, F D R Hobbs, J E Kenkre, A K Roalfe, R Hare, R J Lancashire, M K Davies

Department of Primary Care and General Practice, University of Birmingham, Birmingham B15 2TT

R C Davis
clinical research fellow

F D R Hobbs
professor of primary care and general practice

J E Kenkre
senior research fellow

A K Roalfe
statistician

R Hare
research associate

Department of Public Health and Epidemiology, University of Birmingham

R J Lancashire
computer officer

Department of Cardiology, Selly Oak Hospital, Birmingham
M K Davies
consultant cardiologist

Correspondence to: R C Davis, Sandwell General Hospital, Lyndon, West Bromwich B71 4HJ
R.C.Davis@bham.ac.uk

BMJ 2002;325:1156-8

Abstract

Objectives To determine the prevalence of left ventricular systolic dysfunction, and of heart failure due to different causes, in patients with risk factors for these conditions.

Design Epidemiological study, including detailed clinical assessment, electrocardiography, and echocardiography.

Setting 16 English general practices, representative for socioeconomic status and practice type.

Participants 1062 patients (66% response rate) with previous myocardial infarction, angina, hypertension, or diabetes.

Main outcome measures Prevalence of systolic dysfunction, both with and without symptoms, and of heart failure, in groups of patients with each of the risk factors.

Results Definite systolic dysfunction (ejection fraction < 40%) was found in 54/244 (22.1%, 95% confidence interval 17.1% to 27.9%) patients with previous myocardial infarction, 26/321 (8.1%, 5.4% to 11.6%) with angina, 7/388 (1.8%, 0.7% to 3.7%) with hypertension, and 12/208 (5.8%, 3.0% to 9.9%) with diabetes. In each group, approximately half of these patients had symptoms of dyspnoea, and therefore had heart failure. Overall rates of heart failure, defined as symptoms of dyspnoea plus objective evidence of cardiac dysfunction (systolic dysfunction, atrial fibrillation, or clinically significant valve disease) were 16.0% (11.6% to 21.2%) in patients with previous myocardial infarction, 8.4% (5.6% to 12.0%) in those with angina, 2.8% (1.4% to 5.0%) in those with hypertension, and 7.7% (4.5% to 12.2%) in those with diabetes.

Conclusion Many people with ischaemic heart disease or diabetes have systolic dysfunction or heart failure. The data support the need for trials of targeted echocardiographic screening, in view of the major benefits of modern treatment. In contrast, patients with uncomplicated hypertension have similar rates to the general population.

Introduction

Heart failure and asymptomatic left ventricular dysfunction are underdiagnosed and undertreated. Many

patients are therefore denied the benefit of strongly evidence based treatments, including angiotensin converting enzyme inhibitors¹⁻³ and β blockers.^{4,5} Systematically screening for heart failure and systolic dysfunction is one method that could improve case identification and thereby potentially improve treatment rates.

The most common risk factors for systolic dysfunction and heart failure, in major trials in heart failure, are myocardial infarction, angina, hypertension, and diabetes mellitus.⁶ A screening programme is most cost effective if it is targeted at the patients at highest risk, so identification of how powerfully each of these risk factors predicts systolic dysfunction is important. We investigated this in a prospective substudy of the community based echocardiographic heart of England screening (ECHOES) study.⁷

Methods

Sixteen general practices in the West Midlands region of England were randomly selected, after practices had been stratified geographically and socioeconomically. In this section of the study, we identified all patients with an electronic practice record of myocardial infarction, angina, hypertension, or diabetes. We excluded registered patients who had died or moved and those with severe psychiatric disorders, immobility, or terminal illness.

We sent invitations to 1617 patients selected randomly from the lists obtained. We based eligibility for inclusion on disease registers and did not validate diagnoses before the study. Patients with more than one risk factor were eligible for inclusion in more than one category. All patients who agreed to participate were assessed in their own general practice by clinical history (including prescribed drugs), determination of New York Heart Association functional class, clinical examination, resting 12 lead electrocardiography, and echocardiography including Doppler studies.⁷

We defined heart failure according to European Society of Cardiology criteria—namely, appropriate symptoms (New York Heart Association functional class II or worse) plus objective evidence of cardiac dysfunction.⁸ We classed left ventricular systolic function as “definitely impaired” (ejection fraction < 40%), “borderline” (40-50%), or “normal” (> 50%).

Owing to a lack of agreed diagnostic criteria, we did not attempt to diagnose diastolic dysfunction.

Results

A total of 1062 patients (66%) attended for screening. Table 1 shows the results for left ventricular systolic dysfunction in patients with the defined risk factors, and table 2 gives the results for symptomatic heart failure, subdivided into differing aetiology.

History of myocardial infarction—We found definite systolic dysfunction in 54 (22.1%, 95% confidence interval 17.1% to 27.9%) of the 244 patients with a history of myocardial infarction (table 1). Of these patients, 31 (57.4%) had symptoms of dyspnoea, 17 were asymptomatic and on no treatment, and six were asymptomatic but taking anti-heart failure treatment. An additional 48 (19.7%) patients had borderline systolic function. A total of 39 patients (16.0%, 11.6% to 21.2%) in this risk group had symptoms of dyspnoea with objective evidence of some form of cardiac dysfunction (ejection fraction <40%, atrial fibrillation, valve disease) and thus had definite heart failure (table 2). If patients with ejection fraction 40-50% and symptoms of dyspnoea are included, along with those with ejection fraction <40% and taking drugs for heart failure, but with no current symptoms, the overall prevalence of heart failure (both definite and probable) was 27.5% (22.0% to 33.5%).

History of angina—We found definite systolic dysfunction in 26 (8.1%, 5.4% to 11.6%) of the 321 patients with a history of angina (table 1), of whom 18 (69.2%) had symptoms of dyspnoea. A further 39 (12.1%) patients had borderline left ventricular dysfunction. A total of 27 patients (8.4%, 5.6% to 12.0%) with a record of symptoms of angina had symptoms of dyspnoea and objective evidence of some form of cardiac dysfunction and thus had definite heart failure (table 2). Including patients with borderline systolic dysfunction and symptoms of dyspnoea, and those with definite systolic dysfunction rendered asymptomatic on treatment, the overall prevalence of heart failure (definite and probable) was 15.9% (12.1% to 20.4%).

History of hypertension—As detailed in tables 1 and 2, we found definite systolic dysfunction in only seven (1.8%, 0.7% to 3.7%) of the 388 patients with a history of hypertension, and four (57.1%) of these had symptoms of dyspnoea. Another 31 (8.0%) patients

Table 1 Prevalence of ejection fraction <40% and 40-50% by sex in the diagnostic groups. Values are numbers (percentages)

	Men	Women	Total
Ejection fraction <40%			
Myocardial infarction	36/153 (23.5)	18/91 (19.8)	54/244 (22.1)
Angina	18/173 (10.4)	8/148 (5.4)	26/321 (8.1)
Hypertension	3/197 (1.5)	4/191 (2.1)	7/388 (1.8)
Diabetes	9/111 (8.1)	3/97 (3.1)	12/208 (5.8)
Ejection fraction 40-50%			
Myocardial infarction	33/153 (21.6)	15/91 (16.5)	48/244 (19.7)
Angina	28/173 (16.2)	11/148 (7.4)	39/321 (12.1)
Hypertension	23/197 (11.7)	8/191 (4.2)	31/388 (8.0)
Diabetes	9/111 (8.1)	10/97 (10.3)	19/208 (9.1)

had an ejection fraction of 40-50%. A total of 11 patients (2.8%, 1.4% to 5.0%) with a record of hypertension had definite heart failure. If the 12 patients with borderline systolic function and symptoms of dyspnoea and the two with definitely impaired systolic function rendered asymptomatic on treatment are included as having heart failure, the overall prevalence was 6.4% (4.2% to 9.4%).

History of diabetes—We found an ejection fraction of <40% in 12 (5.8%, 3.0% to 9.9%) of the 208 patients identified as having a history of diabetes (table 1). Ten (83.3%) of these patients had symptoms of shortness of breath. A further 19 (9.1%) patients had an ejection fraction of 40-50%. A total of 16 patients (7.7%, 4.5% to 12.2%) with diabetes had symptoms of dyspnoea with objective evidence of cardiac dysfunction and thus had definite heart failure (table 2). Including patients with borderline systolic function and symptoms, the overall prevalence of heart failure was 12.0% (7.9% to 17.2%).

Statistical analysis—Multiple logistic regression analysis identified only age, previous myocardial infarction, and current systolic blood pressure below 150 mm Hg as independent predictors of ventricular systolic dysfunction. See bmj.com for details of the analysis.

Discussion

Heart failure is common, and we have shown, in the general population arm of the echocardiographic heart of England screening study,⁷ that most patients with left ventricular systolic dysfunction and heart failure in the community have one or more of the risk factors investigated here. The most powerful predictor of systolic dysfunction was previous myocardial infarction.

Table 2 Heart failure on defined criteria (exertional dyspnoea with objective evidence of cardiac dysfunction) in diagnostic groups. Values are numbers (percentages)

	EF <40% alone*	EF <40% + AF	EF <40% + valve disease†	EF <40% + AF + valve disease†	EF 40-50% + AF	EF 40-50% + valve disease†	EF 40-50% + AF + valve disease†	AF (EF >50%)	Valve disease† (EF >50%)	AF + valve disease† (EF >50%)	Total definite HF	EF 40-50% alone	Total HF (definite + probable)
Previous myocardial infarction (n=244)	28 (11.5)	0	1 (0.4)	2 (0.8)	2 (0.8)	1 (0.4)	0	5 (2.0)	0	0	39 (16)	22 (9.0)	67 (27.5)
Angina (n=321)	13 (4.0)	0	4 (1.2)	1 (0.3)	0	0	0	5 (1.6)	3 (0.9)	1 (0.3)	27 (8.4)	22 (6.9)	51 (15.9)
Hypertension (n=388)	3 (0.8)	1 (0.3)	0	0	1 (0.3)	0	0	2 (0.5)	4 (1.0)	0	11 (2.8)	12 (3.1)	25 (6.4)
Diabetes (n=208)	6 (2.9)	2 (1.0)	0	2 (1.0)	0	0	0	3 (1.4)	2 (1.0)	1 (0.5)	16 (7.7)	9 (4.3)	25 (12.0)

AF=atrial fibrillation; EF=ejection fraction; HF=heart failure.

*In addition, six patients with previous myocardial with EF <40% had no symptoms of dyspnoea at time of assessment but were taking diuretics or angiotensin converting enzyme inhibitors; this applied also to two of hypertension group, two of angina group, and none of diabetes group.

†Of the four patients with previous myocardial infarction and valve disease, one had mitral stenosis (MS) and three had mitral regurgitation (MR). Of the four with hypertension and valve disease, three had MR and one had aortic stenosis (AS). Of the nine with angina and valve disease, seven had MR and two had AS. Of the five with diabetes and valve disease, three had MR and two had AS.

What is already known on this topic

The prognosis and symptoms of patients with left ventricular systolic dysfunction and heart failure can be greatly improved by modern treatments

Many patients with heart failure do not have an assessment of left ventricular function, resulting in undertreatment of the condition

What this study adds

Patients with a history of ischaemic heart disease (especially those with previous myocardial infarction) or diabetes commonly have left ventricular systolic dysfunction

These patients would be candidates for a targeted echocardiographic screening programme

In contrast, the yield from screening patients with uncomplicated hypertension would be low

tion. A high prevalence of systolic dysfunction also occurred in the groups with angina and diabetes, but regression analysis did not show these to be independent risk factors. Confining screening to patients with a history of ischaemic heart disease would give a high yield of candidates for evidence based treatments, which alter prognosis as well as symptoms.

In the Framingham heart study hypertension, alone or associated with ischaemic or rheumatic heart disease, was the most common condition predisposing to heart failure, diagnosed with a clinical and radiological scoring system.⁹ In contrast, our data and those from Glasgow suggest that uncomplicated hypertension is less important in predicting left ventricular dysfunction.¹⁰ Even including all causes of heart failure, using objective criteria for diagnosis, the prevalence was only slightly higher in patients with hypertension than in the general population (2.8% *v* 2.3%). The reasons for the differences are not clear. Patients with hypertension in our study had all been identified and treated, which may have reduced development of heart failure. Indeed, recent Framingham data show greatly improved treatment of hypertension.¹¹ Alternatively, some people diagnosed with heart failure by the Framingham scoring system may not have met the diagnostic criteria used here; hypertension is a common cause of radiological cardiomegaly, a major Framingham criterion. Our finding on regression analysis that systolic blood pressure below 150 mm Hg is a predictor of systolic dysfunction is counterintuitive and is likely to reflect the use of angiotensin converting enzyme inhibitors and other antihypertensive drugs in patients with previously identified systolic dysfunction.

We obtained the clinical and demographic data through investigation, whereas details of which patients had the risk factors were obtained from computerised records. Apparent differences (for example, only 91% of patients with a computerised record of history of myocardial infarction gave this diagnosis on questioning) may be due to patients' uncertainty about the diagnosis, incorrect practice record data, or both. A 98% agreement in the case of diabetes suggests that practice records are on the whole accurate. All but 28 of the patients with a computerised record of myocardial infarction had electrocardiographic abnormalities.

In any event, our method represents "real life," replicating a standard method of selecting patients for screening in routine practice.

We believe that these are the first data on the potential diagnostic yield from a targeted, systematic screening programme for heart failure and systolic dysfunction. Such practice based electronic record searching for the at risk population and subsequent echocardiography are feasible. Formal trials are now justified.

The national service framework for coronary heart disease states that "primary care teams and hospitals should put in place models of care so that they use a systematic approach to identify people at high risk of heart failure (e.g. people who have had an acute myocardial infarction)."¹² Our data support this approach.

S McLeod assisted in practice recruitment and coordination, and T Marshall assisted with the initial sampling frame of practices and patients. D Wosornu contributed to study design and carried out some clinical assessments and reports. The following practices kindly agreed to take part in the study: Albrighton Surgery, Albrighton; Baker St Surgery, Stoke on Trent; Castle Practice, Castle Bromwich; Church St Surgery, Kidderminster; Droitwich Health Centre, Droitwich; Elgar House Surgery, Redditch; Great Barr Group Practice, Birmingham; Hollies Medical Practice, Tamworth; Holmcroft Surgery, Stafford; Laurie Pike Health Centre, Handsworth, Birmingham; Lee Bank Health Centre, Birmingham; Ley Hill Surgery, Sutton Coldfield; Moss Grove Surgery, Kingswinford; Red Roofs Surgery, Nuneaton; Riverbrook Medical Centre, Stirchley, Birmingham; and Shirley Medical Centre, Shirley, Solihull.

Contributors: See bmj.com

Funding: NHS research and development cardiovascular disease and stroke programme.

Competing interests: FDRH is a member of the European Society of Cardiology working party on heart failure, chair of the British Primary Care Cardiovascular Society, and Treasurer of the British Society for Heart Failure. MKD is chairman of the British Society for Heart Failure. FDRH and MKD have received travel sponsorship and honorariums from several biotechnology and pharmaceutical companies with cardiovascular products for plenary talks and attendance at major cardiology scientific congresses and conferences.

- Garg R, Yusuf S for the Collaborative Group on ACE Inhibitor Trials. Overview of randomized trials of angiotensin-converting enzyme inhibitors on mortality and morbidity in patients with heart failure. *JAMA* 1995;273:1450-6.
- The SOLVD Investigators. Effect of enalapril on mortality and the development of heart failure in asymptomatic patients with reduced left ventricular ejection fractions. *N Engl J Med* 1992;327:685-91.
- Pfeffer MA, Braunwald E, Moy LA, Basta L, Brown EJ, Cuddy TE for the SAVE Investigators. Effect of captopril on mortality and morbidity in patients with left ventricular dysfunction after myocardial infarction: results of the survival and ventricular enlargement trial. *N Engl J Med* 1992;327:669-77.
- CIBIS-II Investigators and Committees. The cardiac insufficiency bisoprolol study II. *Lancet* 1999;353:9-13.
- MERIT-HF Study Group. Effect of metoprolol CR/XL in chronic heart failure: metoprolol CR/XL randomised intervention trial in congestive heart failure (MERIT-HF). *Lancet* 1999;353:2001-7.
- The SOLVD Investigators. Effect of enalapril on survival in patients with reduced left ventricular ejection fractions and congestive heart failure. *N Engl J Med* 1991;325:293-302.
- Davies MK, Hobbs FDR, Davis RC, Kenkre JE, Roalfe AK, Hare R, et al. Prevalence of left-ventricular systolic dysfunction and heart failure in the echocardiographic heart of England screening study: a population based study. *Lancet* 2001;358:439-44.
- Task Force on Heart Failure of the European Society of Cardiology. Guidelines for the diagnosis of heart failure. *Eur Heart J* 1995;16:741-51.
- McKee PA, Castelli WP, McNamara P, Kannell WB. The natural history of heart failure: the Framingham study. *N Engl J Med* 1971;285:1441-6.
- McDonagh TA, Morrison CE, Lawrence A, Ford I, Tunstall-Pedoe H, McMurray JJV, et al. Symptomatic and asymptomatic left ventricular systolic dysfunction in an urban population. *Lancet* 1997;350:829-33.
- Mosterd A, D'Agostino RB, Silbershatz H, Sytkowski PA, Kannel WB, Grobbee DE, et al. Trends in the prevalence of hypertension, antihypertensive therapy and left ventricular hypertrophy. *N Engl J Med* 1999;340:1221-7.
- Department of Health. *National service framework for coronary heart disease*. London: Stationery Office, 2000.

(Accepted 15 August 2002)