



This is an abridged version; the full version is on bmj.com

Editorial by Clauw and p 1370

Unit of Chronic Disease Epidemiology, School of Epidemiology and Health Sciences, University of Manchester, Manchester M13 9PT

Gary J Macfarlane *professor of epidemiology*
Anne-Marie Biggs *research associate*

London School of Hygiene and Tropical Medicine, University of London, London WC1E 7HT

Noreen Maconochie *senior lecturer in epidemiology and statistics*
Patricia Doyle *reader in epidemiology*

Gulf War Illnesses Research Unit, Department of Psychological Medicine, Guy's, King's, and St Thomas's School of Medicine, London SE5 8AZ

Matthew Hotopf *reader in psychological medicine*

Arthritis Research Campaign Epidemiology Unit, School of Epidemiology and Health Sciences, University of Manchester
Mark Lunt *research fellow*

Correspondence to: G J Macfarlane
G.Macfarlane@man.ac.uk

BMJ 2003;327:1373-5

Incidence of cancer among UK Gulf war veterans: cohort study

Gary J Macfarlane, Anne-Marie Biggs, Noreen Maconochie, Matthew Hotopf, Patricia Doyle, Mark Lunt

Abstract

Objectives To determine whether incidence rates of cancer are higher in UK service personnel who were deployed in the Gulf war than in those not deployed and whether any increased risk of cancer is related to self reported exposures to potentially hazardous material during the period of deployment.

Design A cohort study with follow up from 1 April 1991 (the end of the Gulf war) to 31 July 2002.

Participants 51 721 Gulf war veterans and 50 755 service personnel matched for age, sex, rank, service, and level of fitness who were not deployed in the Gulf (the Era cohort).

Main outcome measures Incident cancers, identified on the NHS central register.

Results There were 270 incident cancers among the Gulf cohort and 269 among the Era cohort (incidence rate ratio 0.99, 95% confidence interval 0.83 to 1.17). There was no excess in site specific cancers among the Gulf cohort. Adjustment for lifestyle factors (smoking and alcohol consumption) did not alter these results. In the Gulf cohort, risk of cancer was not related to multiple vaccinations or exposure to pesticides or depleted uranium during deployment.

Conclusion There is no current excess risk of cancer overall nor of site specific cancers in Gulf war veterans. Specific exposures during deployment have not resulted in a subsequent increased risk of cancer. The long latent period for cancer, however, necessitates the continued follow up of these cohorts.

Introduction

Studies among UK veterans of the Gulf war have shown an excess of self reported morbidity of all types compared with veterans serving at the same time who were fit for active service but who were not deployed.^{1,2} However, no unusual clusters of symptoms are evident,^{2,3} and the symptoms most commonly reported are non-specific.

There has been concern about a possible link between service in the Gulf and an increased risk of cancer, particularly in those exposed to depleted uranium. A review of the evidence, however, concluded that, except in extreme circumstances, any excess risk is likely to be small.⁴ A US study among a sample of Gulf war veterans found an excess of testicular cancer in the short term,⁵ while a study of admissions to hospital found excess admissions for cancers (mostly benign) during 1991. Mortality studies of US⁶ and UK veterans,⁷ however, found no increased risk of death from cancer.

We examined whether UK Gulf veterans have experienced increased incidence rates of cancer in the 11 years since the end of the war compared with service personnel not deployed; whether any excess risk, if found, could be explained by the possible confounding

effects of lifestyle factors (such as cigarette smoking and alcohol consumption); and whether risk of cancer is related to self reported exposures to potentially hazardous material during the period of deployment.

Methods

We carried out a cohort study of all UK armed forces personnel who served in the Gulf area sometime between September 1990 and June 1991 (the "Gulf" cohort). A comparison group of the same size was randomly selected from members of the armed forces who were in service on 1 January 1991 but who were not deployed to the Gulf area (the "Era" cohort). This comparison group was stratified to match the Gulf cohort on age, sex, service branch and rank, and (for the army and air force) level of fitness for active service. Both cohorts were identified by staff from the Gulf Veterans' Illnesses Unit of the Ministry of Defence.

Over 96% of members of both cohorts were "flagged" on the NHS central register. This register, established in 1939, contains an entry for everyone in the United Kingdom who has ever been registered with a general practitioner and all people born in, or who have immigrated to, the United Kingdom. The register contains information on cancers diagnosed, on deaths, and on people emigrating from the United Kingdom. We obtained information on the first diagnoses of malignant cancer with date of diagnosis before 31 July 2002. We calculated person years at risk for cancer from 1 April 1991 (the nominal end of the Gulf War) until the earliest of either date of emigration from the United Kingdom, date of death, date of first diagnosis of cancer, or 31 July 2002.

We analysed the data with adjustment for sex, age group, service branch and rank, and a comparison between Gulf and Era veterans using an incidence rate ratio with 95% confidence interval. To determine whether the incidence rate ratio changed with time, it was calculated separately for the periods before (inclusive) and after December 1997 (this date was chosen to give the same number of incident cancers in the two time periods).

The Gulf and Era cohorts also acted as a sampling frame for three morbidity surveys conducted between 1997-2001.^{1,2,8} The first two, both surveys of general health, drew random samples from each of the cohorts (using non-overlapping samples) while the third survey, a study of reproductive and child health, included all people in both cohorts. These studies collected information in a consistent way about tobacco smoking, alcohol intake at the time of the survey, and, in the Gulf cohort, exposures to potentially hazardous material during service in the Gulf. We used information from the third study if possible (n = 40 853) and otherwise from studies 1 and 2 (n = 8494). Among people participating in at least one

Table 1 Occurrence of cancer among cohort members according to whether they served in the Gulf: demographic and military characteristics

	Gulf cohort (n=51 721)	Era cohort (n=50 755)	Incidence rate ratio (95% CI)
All cancers	270	269	0.99 (0.83 to 1.17)
Men	257	256	0.99 (0.83 to 1.17)
Women	13	13	0.98 (0.47 to 2.12)
Age (years):			
<20	11	12	0.90 (0.40 to 2.03)
20-24	39	48	0.79 (0.52 to 1.21)
25-29	46	41	1.11 (0.73 to 1.69)
30-34	41	40	1.02 (0.66 to 1.57)
35-39	52	38	1.37 (0.90 to 2.08)
>39	81	90	0.89 (0.66 to 1.20)
Rank:			
Officer	77	64	1.18 (0.84 to 1.64)
Other ranks	193	205	0.93 (0.76 to 1.13)
Service branch:			
Army	170	175	0.95 (0.77 to 1.17)
Navy	38	26	1.46 (0.88 to 2.40)
Air Force	62	68	0.92 (0.65 to 1.29)

of these three morbidity surveys, we compared the incidence of cancer between Gulf and Era cohort members, adjusting for the effects of tobacco smoking and alcohol intake. Secondly, we compared cancer incidence according to certain exposures during the time in the Gulf. Effects are again reported as incidence rate ratios and 95% confidence intervals.

Results

Both cohorts initially comprised 53 462 members. After exclusions we followed up 51 721 members of the Gulf cohort and 50 755 members of the Era cohort. Baseline demographic characteristics of each were similar (see bmj.com).

During follow up cancer was diagnosed in 270 Gulf veterans and 269 Era veterans (table 1). There were no large or significant excess of cancers in any subgroup. The incidence rate ratio did not differ significantly between the period up to and including 1997 (1.04, 0.82 to 1.32) and the later period (0.93, 0.74 to 1.19). After exclusion of non-melanoma skin cancers and

cancers for which site information was unavailable, there were a total of 213 and 215 Gulf and Era veterans, respectively, in whom cancer was diagnosed (0.97, 0.81 to 1.18).

A comparison of incidence rates of cancer among subgroups of cohort members who provided information on smoking and alcohol similarly showed no significant excess of cancers among the Gulf veterans. This incidence rate ratio was slightly, but not significantly, higher than that among cohort members who did not take part in one of the morbidity surveys (0.95, 0.75 to 1.19). Among cohort members who did participate in a morbidity survey, the incidence rate ratio did not change after we adjusted for smoking status (current, past, never smoker) and for the usual intake of alcohol at the time they completed the questionnaire (table 2). This additional adjustment for tobacco and alcohol intake also resulted in little or no change in any of the risks for site specific cancers (table 2). Finally, among the Gulf veterans who participated in at least one of the morbidity surveys we examined the incidence of cancer with respect to some of the exposures reported during deployment to the Gulf. There was no subsequent significant excess risk of cancer, either overall (table 3) or for site specific cancers.

Discussion

Our study has shown that the incidence of cancer in those who served in the Gulf was similar to that in veterans who were fit for active service but were not deployed. This result did not change when we took into account smoking and alcohol consumption among a subgroup who participated in one of the UK morbidity surveys. This subgroup had no significant excess risk of cancer associated with reported exposure to pesticides, multiple vaccination against anthrax, plague and pertussis, or exposure to depleted uranium.

There are several methodological considerations regarding the current study. Firstly, though we excluded the small number of cohort members who could not be traced through the NHS central register, some information was available on diagnoses of cancer in

Table 2 Occurrence of site specific cancer among cohort members according to whether they served in the Gulf: adjusted for smoking and alcohol consumption

Cancer type	ICD-10 code	Gulf (n=28 518)	Era (n=20 829)	Incidence rate ratio	
				Unadjusted	Adjusted
All cancers	C00-97	144	95	1.11 (0.86 to 1.44)	1.12 (0.86 to 1.45)
All excluding non-melanoma skin cancer	All except C44	109	75	1.06 (0.79 to 1.43)	1.07 (0.79 to 1.43)
Oral cavity	C00-14	4	5	0.59 (0.16 to 2.18)	0.58 (0.16 to 2.16)
Upper digestive tract	C15-17	2	2	0.73 (0.10 to 5.19)	0.69 (0.10 to 4.93)
Lower digestive tract	C18-21	6	6	0.73 (0.24 to 2.27)	0.73 (0.24 to 2.27)
Bronchus, lung, and trachea	C33-34	3	5	0.49 (0.10 to 1.84)	0.41 (0.10 to 1.73)
Malignant melanoma skin	C43	10	5	1.46 (0.50 to 4.28)	1.50 (0.51 to 4.40)
Other skin	C44	35	20	1.28 (0.74 to 2.22)	1.30 (0.75 to 2.26)
Breast	C50	5	4	0.91 (0.25 to 3.41)	0.98 (0.26 to 3.64)
Prostate	C61	4	3	0.98 (0.22 to 4.36)	1.03 (0.23 to 4.62)
Testis	C62	24	15	1.17 (0.61 to 2.23)	1.17 (0.61 to 2.23)
Urinary tract	C64-8	7	4	1.28 (0.37 to 4.37)	1.29 (0.38 to 4.41)
Central nervous system	C69-72	12	8	1.10 (0.45 to 2.68)	1.08 (0.44 to 2.65)
Lymphoid and haematopoietic	C81-96	24	11	1.60 (0.78 to 3.26)	1.60 (0.79 to 3.28)
All other sites	Various*	8	7	0.84 (0.30 to 2.31)	0.84 (0.30 to 2.32)
ICD code unavailable		2	0		

*C22-32, 35-42, 45-49, 51-60, 63, 73-80, 97.

Table 3 Risk of any cancer by exposure agent among cohort members who served in the Gulf

Exposure agent	Not exposed		Exposed		Incidence rate ratio (95% CI)
	Total	Cancers	Total	Cancers	
Multiple vaccination (anthrax, plague, pertussis)	21 313	107	7 205	39	1.08 (0.75 to 1.56)
Used personal pesticides or insecticides	13 125	82	15 393	64	0.66 (0.48 to 0.92)
Handled pesticides	26 467	134	2 051	12	1.16 (0.64 to 2.09)
Living quarters treated with pesticides	22 251	115	6 267	31	0.96 (0.64 to 1.42)
Depleted uranium	26 426	139	2 092	7	0.63 (0.30 to 1.36)

these people from alternative sources. Among them the number of cancers was higher in the Era cohort (n = 11) than in the Gulf cohort (n = 3). Secondly, information on lifestyle habits was obtained only at the time of the morbidity surveys, 6-10 years after the Gulf war. These data show that the proportion who were current smokers and the amount of alcohol drunk in a week were similar in the two cohorts,^{1 2} and thus the estimates of rate ratios are little changed with adjustment for these factors. We will have underestimated the relative risk of cancer in Gulf war veterans only if they habitually smoked less or drank less alcohol and had recently increased consumption (or vice versa in relation to Era veterans). Neither seems plausible. Finally, the “exposures” reported at the time of the Gulf are self reported, and people who develop an adverse health outcome may differentially recall past events. However, this would over estimate the observed effect of an exposure on health, and could not explain our failure to detect true adverse effects of any of these exposures if they were to exist.

A mortality study among the same group of UK Gulf and Era veterans showed only a small non-significant increased risk of death from cancer nine years after the end of the conflict.⁷ A mortality study among US veterans also failed to show an increased risk of death from cancer seven years after the end of the conflict⁶ or an increased risk of death from cancer among those potentially exposed to nerve gas. An early report of an increased risk of testicular cancer in the period immediately after deployment was no longer evident on further follow up.⁵

Although this study should provide some reassurance of a lack of association between deployment to the Gulf and increased risk of cancer, the long latent period for cancer requires that these cohorts should continue to be followed up and their experience of cancer monitored.

We acknowledge the contribution of other researchers to the conduct of the United Kingdom morbidity surveys and mortality study: David Baxter, Nicola Cherry, William Coker, Francis Creed, Anthony David, Graham Davies, Graham Dunn, Susan Ferry, Lisa Hull, Khalida Ismail, Samantha Lewis, Ian Palmer, Margo Pelerin, Sue Prior, Patrick Sampson, Alan J Silman, Joanne Smedley, Stewart Taylor, Catherine Unwin, and Simon Wessely. We thank Nick Blatchley (Ministry of Defence), who identified the study cohorts, and staff from National Statistics in Southport, who traced them on the NHS central register. Contributors: See bmj.com

Funding: Ministry of Defence.

Competing interests: None declared.

Ethical approval: University of Manchester Committee on the Ethics of Research on Human Beings.

- Unwin C, Blatchley N, Coker W, Ferry S, Hotopf M, Hull L, et al. Health of UK servicemen who served in Persian Gulf War. *Lancet* 1999;353: 169-78.
- Cherry N, Creed F, Silman A, Dunn G, Baxter D, Smedley J, et al. Health and exposure of United Kingdom Gulf war veterans. Part 1: The pattern and extent of ill-health. *Occup Environ Med* 2001;58:291-8.
- Everitt B, Ismail K, David AS, Wessely S. Searching for a Gulf War syndrome using cluster analysis. *Psychol Med* 2002;32:1371-8.

What is already known on this topic

Veterans of the 1990-1 Gulf war have an excess of self reported symptoms

Their mortality experience (including cancer mortality) is similar to that seen in military personnel who were serving at the same time but who were not deployed in the Gulf

What this study adds

Incidence of and mortality from cancer in Gulf War veterans is almost identical to that seen in veterans who were not deployed in the Gulf

This comparison takes account of lifestyle factors (smoking and alcohol consumption), which are known to influence cancer risk

The risk of cancer was no higher in Gulf war veterans who reported exposure to specific substances such as depleted uranium or pesticides

- Royal Society Report. *The health effects of depleted uranium munitions Part 1*. London: Royal Society, 2002.
- Knoke JD, Gray GC, Garland FC. Testicular cancer and Persian Gulf War service. *Epidemiology* 1998;9:648-53.
- Kang HK, Bullman TA. Mortality among US veterans of the Persian Gulf War: 7-year follow-up. *Am J Epidemiol* 2001;154:399-405.
- Macfarlane GJ, Thomas E, Cherry N. Mortality among UK Gulf War Veterans. *Lancet* 2000;356:17-21.
- Maconochie N, Doyle P, Davies G, Lewis S, Pelerin M, Prior S, et al. The study of reproductive outcome and the health of offspring of UK veterans of the Gulf war: methods and description of the study population. *BMC Public Health* 2003;3:4. (Accepted 7 October 2003)

Endpiece

Reporters and wolves

Recalling his father, a professional photographer and journalist, Joseph Brodsky wrote, “For several years he freelanced all over the country under contract to the All Union Agricultural Exhibit in Moscow ... but the pay was less than meagre and the three of us existed solely on my mother’s salary as a clerk in the borough’s development council ... [It] appears that most of his life was spent on foot (‘reporters, like wolves, live by their paws,’ was his favourite utterance), among ships, sailors, captains, cranes, cargo ... he liked to be near the water, he adored the sea. In that country [the Soviet Union], this is the closest one gets to freedom. Even looking at it is sometimes enough, and he looked at it, and photographed it, for most of his life.” At the end of their lives, the Soviets refused to let Brodsky’s aged parents out to visit their son.

Joseph Brodsky in “In a room and a half”
from his book of selected essays,
Less than one, 1985

Fred Charatan, retired geriatric physician, Florida