

CME

Follow the link from the online version of this article to obtain certified continuing medical education credits

Management of sickle cell disease in the community

Valentine Brousse,¹ Julie Makani,^{2,3} David C Rees⁴

¹Department of Pediatrics, Reference Centre for Sickle Cell Disease, Hôpital Universitaire Necker-Enfants Malades, APHP, Paris, France

²Department of Haematology and Blood Transfusion, Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania

³Nuffield Department of Medicine, Oxford, UK

⁴Department of Paediatric Haematology, King's College Hospital NHS Foundation Trust, King's Health Partners, London SE5 9RS, UK

Correspondence to: D Rees david.rees2@nhs.net

Cite this as: *BMJ* 2014;348:g1765
doi: 10.1136/bmj.g1765

bmj.com

Previous articles in this series

- ▶ Coeliac disease (*BMJ* 2014;348:g1561)
- ▶ Fibromyalgia (*BMJ* 2014;348:g1224)
- ▶ Trigeminal neuralgia (*BMJ* 2014;348:g474)
- ▶ Management of traumatic amputations of the upper limb (*BMJ* 2014;348:g255)
- ▶ Managing wheeze in preschool children (*BMJ* 2014;348:g15)

Sickle cell disease is characterised by unpredictable episodes of acute illness, progressive organ damage, and a lack of effective treatments. It is one of the most common inherited conditions, although its prevalence varies widely. Median life expectancy is currently 40-60 years in high income countries but much less in low income areas.^{1,2} It is associated with protean clinical complications. Patients present to all medical specialties and increasingly to general practitioners. This review aims to provide an evidence based update on how to manage patients with this disease in the community. It does not consider sickle cell trait, which is largely asymptomatic.

What is sickle cell disease?

Sickle cell disease is a recessive condition, caused by a mutation in the β globin gene, which changes the sixth amino acid from glutamic acid to valine. Sickle haemoglobin (HbS) is insoluble when deoxygenated, forming long polymers. These polymers damage the red cell membrane, resulting in rigid sickle shaped cells with a tendency to cause vaso-occlusion and a cascade of pathological events, including infarction, vasculopathy, haemolysis, oxidative stress, hypercoagulability, and inflammation (figure).³

At least 15 different genotypes cause sickle cell disease, although all include the HbS allele. Homozygous disease (HbSS), usually called sickle cell anaemia (SCA), is the most common and severe type of disease in most populations. HbSC disease and HbS/ β thalassaemia are the two other common genotypes.⁴

Where is sickle cell disease common?

About 300 000 babies with sickle cell disease are born worldwide each year, with an estimated 90 000 births in Nigeria and 40 000 in the Democratic Republic of Congo in 2010. Around 40 000 affected children are born in India

SOURCES AND SELECTION CRITERIA

We searched Medline, the Cochrane Database, and ClinicalTrials.gov using the term "sickle" together with certain complications including pain, infection, enuresis, renal failure, spleen, acute chest syndrome, leg ulcers, and stroke. Preference was given to randomised clinical trials, and when these were unavailable large case series were sought. We prioritised newer studies over older ones.

each year, with 10 000 in the Americas, 10 000 in the Eastern Mediterranean, and 2000 in Europe.² Numbers are increasing, and sickle cell disease is thought to be the most common severe genetic disease in the United Kingdom and France, with 10 000-15 000 patients in each country.⁵ The condition was traditionally found in populations of African descent in northern Europe, but this is changing with increasing numbers of mixed race people, particularly in large cities such as London and Paris.

How do patients with sickle cell disease present?

The first symptom in infants is typically dactylitis, or hand-foot syndrome—painful swelling of the hands or feet as a result of vaso-occlusion. This affects 30% of patients in the first year of life and usually resolves in a few days.¹⁰ Treatment with simple analgesia is often sufficient, although severe episodes may require admission to hospital for opiates, particularly if this is the first presentation or the diagnosis is uncertain. Dactylitis is uncommon after 2 years of age.¹⁰

Invasive infection with encapsulated bacteria can be a presenting feature, related to functional hyposplenism, which is seen in 90% of 5 year olds with HbSS disease.¹¹ The relative risk of *Streptococcus pneumoniae* infection in young children with this disease compared with normal controls is 300-600,¹² with a mortality rate of 15%.¹⁰ The relative risk of invasive infection with *Haemophilus influenzae* is 20-100, with a mortality rate of about 20%. Two randomised controlled trials found that penicillin prophylaxis significantly reduces morbidity and mortality associated with pneumococcal infection in children under 5,^{13,14} with oral prophylaxis reducing pneumococcal septicaemia by 84%.¹³

In the absence of neonatal screening, most cases of sickle cell disease present with symptoms in infancy or early childhood. However, patients with milder forms, such as HbSC disease, may be diagnosed as adults with acute pain, or on incidental blood testing, particularly during pregnancy.

What acute complications occur in sickle cell disease?

Here we describe the common acute complications of sickle cell disease and offer advice for GPs on assessing

SUMMARY POINTS

All children with sickle cell disease should take penicillin twice a day until 5 years of age at least

Acute neurological symptoms in children and adults with sickle cell disease necessitate urgent referral to hospital

All patients should receive annual vaccination against influenza and other appropriate vaccinations where available

Patients living in most parts of Africa should use insecticide treated bed nets and take malarial prophylaxis

Children with severe types of sickle cell disease (HbSS and HbS/ β^0 thalassaemia) should be offered primary stroke prevention with annual transcranial Doppler scans and blood transfusion when resources allow this

Children and adults should be offered treatment with hydroxyurea if they have two or more episodes of severe acute pain in a year, or acute chest syndrome

Children, families, and adults should be offered education about sickle cell disease and managing its complications




Diagram showing the pathophysiology and some of the major clinical complications of sickle cell disease. HbS=sickle haemoglobin

Indications for urgent referral to hospital in sickle cell disease

- Severe pain not controlled by simple analgesia or low dose opioids
- Dehydration caused by severe vomiting or diarrhoea
- Severe sepsis: temperature $>38.5^{\circ}\text{C}$ or $>38^{\circ}\text{C}$ if under 2 years old, temperature $<36^{\circ}\text{C}$, or hypotension
- Symptoms or signs of acute chest syndrome including tachypnoea, oxygen saturation more than 5% below steady state, signs of lung consolidation
- New neurological symptoms or signs
- Symptoms or signs of acute fall in haemoglobin
- Acute enlargement of spleen or liver over 24 hours, particularly in young children
- Marked increase in jaundice
- Haematuria
- Fulminant priapism lasting more than two hours or worsening of recurrent episodes

and managing these problems in the community and when to refer. The feasibility of managing acute complications in the community varies geographically, depending on provision of community care and home circumstances. The box lists indications for urgent referral to hospital. In some high prevalence areas, such as London, community nurse specialists with specific knowledge of sickle cell disease can advise families on managing acute illness and may be able to make home visits.

Pain

Acute pain is the most common manifestation of the disease in all age groups. Pain is thought to be caused by vaso-occlusion, which results in ischaemic tissue damage and subsequent inflammation and pain. It can affect any part of the body, although the limbs and back are most commonly involved. The US Cooperative Study of Sickle Cell Disease (CSSCD) found an incidence of 0.8 episodes of acute pain per patient year in HbSS and 0.4 in HbSC disease. Pain was most common in 10-30 year olds, and high pain rates were associated with increased mortality after 20 years of age. Increased pain rates were associated with low levels of fetal haemoglobin (HbF) and high total haemoglobin levels.¹⁵ Initial assessment should include an estimate of pain severity, looking for evidence of infection, severe dehydration, or other precipitating factors that would necessitate referral to hospital. General advice includes keeping warm; drinking plenty; and taking simple analgesia, such as paracetamol, ibuprofen, and weak opioids if pain is severe. There is no good evidence to support any particular form of analgesia, although many patients will prefer a certain drug or combination. Children are more likely to be managed at home, with parents acting as carers. A study of children with sickle cell disease in south London showed an admission rate of 41 per 100 patient years in 2009, compared with 111 per 100 patient years 30 years earlier.¹⁶

Most episodes of acute pain can be managed at home with simple analgesia. Severe pain usually lasts one to five days, with full recovery after one week, but this varies widely. Severe pain is usually managed in hospital with opiates, as recommended by National Institute for Health and Care Excellence guidelines.¹⁷

Acute chest syndrome

Acute chest syndrome is defined as a new pulmonary infiltrate on chest radiography in a patient with sickle cell disease, accompanied by symptoms such as chest pain, cough, dyspnoea, or tachypnoea. It occurs at all ages and has a peak incidence of 34 per 100 patient years at the age of 3 years.¹⁰ The underlying lung disease involves a combination of infection, vaso-occlusion, endothelial dysfunction, and fat embolism in some cases.⁴ A CSSCD study of 671 episodes identified infectious agents in 249 cases. The most common agents were chlamydia, mycoplasma, and respiratory syncytial virus; *S pneumoniae* was relatively uncommon, probably as a result of penicillin prophylaxis and vaccination.¹⁸ Acute chest syndrome is serious because increasing hypoxia creates a vicious circle of increasing vaso-occlusion and worsening hypoxia. Patients usually present to their GP with respiratory tract infections, and most improve with simple outpatient management. Refer patients to hospital if they have a high fever, clinical evidence of lung consolidation, widespread wheezing, signs of respiratory distress, or new hypoxaemia as assessed by pulse oximetry. Patients with coexisting asthma are particularly vulnerable.¹⁹ In the CSSCD study, 13% required mechanical ventilation and 3% died. Hospital management of the syndrome is based on evidence from case series, and involves broad spectrum antibiotics, bronchodilators, ventilatory support, and blood transfusion in severe or deteriorating cases.¹⁸

Neurological problems

Worldwide, HbSS disease is probably the most common cause of stroke in children. In the absence of stroke prevention programmes, these patients have an incidence of infarctive stroke of 0.61 per 100 patient years, with a peak incidence of 1.02 per 100 patient years in children aged 2-5 years.²⁰ Infarctive stroke is caused by vasculopathy of the large arteries, typically the middle cerebral artery.²¹ Stroke can present in many ways including hemiparesis, speech problems, seizures, or altered levels of consciousness. Acute management is largely based on expert opinion and involves urgent transfusion to correct anaemia and reduce the amount of HbS in the blood.²¹ The incidence of intracranial haemorrhage is also increased, mainly in young adults.²⁰ Bleeding occurs from fragile new arteries, which grow in response to ischaemia, and from aneurysms, which result from progressive vasculopathy. Over the past decade, the incidence of stroke in high income countries has fallen owing to primary stroke prevention based on transcranial Doppler scanning to identify children at high risk and selective use of regular blood transfusions.²²⁻²³ Silent cerebral infarcts are seen on magnetic resonance imaging in about 30% of children. Although these are not associated with overt neurological events, they are associated with seizures, cognitive impairment, and headaches.²⁴ Urgently refer any patient with sudden onset headaches, reduced levels of consciousness, limb or facial weakness, or acute cognitive impairment to hospital.

Around 30% of adults and children have neurocognitive and behavioural problems as a result of cerebrovascular disease and need multidisciplinary support,²⁴ including neuropsychology, educational psychology, physiotherapy, occupational therapy, and social services.

Anaemia

Haemoglobin levels in HbSS disease are typically 70-80 g/L, and this chronic anaemia is well tolerated. Although any febrile illness or vaso-occlusive complication can worsen anaemia, the two main causes of an acute fall in haemoglobin of more than 20% below the baseline level are splenic sequestration and parvovirus B19 infection. Splenic sequestration is often triggered by sepsis and occurs when the spleen rapidly enlarges and traps red blood cells. It typically occurs in children under 5 years and is potentially fatal without urgent blood transfusion. If two or more episodes occur, splenectomy is often recommended, on the basis of case series that show a high rate of recurrence.²⁵ Parvovirus B19 infection causes slapped cheek syndrome and also infects erythroid cells, causing transient reticulocytopenia, which last about a week. Because sickled red cells have a shorter survival time than normal ones, reticulocytopenia can result in severe anaemia requiring blood transfusion, with recovery occurring after about seven days.²⁶ Increased anaemia can also be associated with folate and iron deficiency, and renal impairment, although this is usually of gradual onset.

Acutely anaemic patients may present to GPs with increased exertional dyspnoea or other cardiovascular symptoms, although anaemia is often part of a more apparent infective or vaso-occlusive picture. Examination should include assessment of pallor, jaundice, and hepatospleno-

megaly. Urgently refer patients with cardiovascular signs or symptoms, or new enlargement of the liver or spleen, to hospital. Blood tests should include a full blood count, reticulocyte count, haematinics, and parvovirus B19 serology. Patients should be referred to or discussed with a specialist sickle cell disease team in the hospital if haemoglobin is more than 20 g/L below their known steady state.

Infection

Acute infection is one of the most common causes of admission to hospital in patients under 10 years. In a CSSCD study of 694 infants followed prospectively, bacteraemia had a peak incidence of 9.9 per 100 patient years in the first year of life and steadily declined with increasing age.¹⁰ *S pneumoniae* is an important pathogen at all ages, and salmonella infections are particularly associated with osteomyelitis.²⁷ The spectrum of infections changes with age, and Gram negative bacterial and staphylococcal infections are more important in adults.

Suspect infection in the presence of a fever or localising symptom, as in any other patients. Particular concerns when assessing and treating sickle cell disease include increased risk of osteomyelitis and acute chest syndrome. Patients with clinical evidence suggesting either of these conditions should be urgently referred to hospital. Infections of the upper respiratory tract, urinary tract, ear, and skin can usually be managed in the community with oral antibiotics if necessary, unless features in the box are also present.

Many patients are already on regular penicillin prophylaxis; this should be changed to an alternative if antibiotic treatment is needed and subsequently restarted.

Sickle cell disease can affect every organ in the body and a wide range of other acute problems can occur (table).

What chronic complications are common in sickle cell disease and how are they managed in the community?

Chronic complications increase with age and are becoming more common in high income countries as people live longer. Chronic complications are largely managed in the community, through a combination of families, GPs, community paediatrics, nurse specialists, and voluntary agencies. The most common chronic problem is pain, and management relies on physiotherapy and psychological support.²⁸ Pulmonary complications, including restrictive lung disease and pulmonary hypertension, are prevalent in older teenagers and adults. Pulmonary hypertension, confirmed on cardiac catheterisation, affects about 6% of adults and is associated with premature death.²⁹ Management includes regular blood transfusion, hydroxyurea, and drugs used in idiopathic pulmonary hypertension, such as endothelin receptor antagonists and prostaglandins. Community support including home oxygen is necessary with increasing disability.⁴ Locomotor and neurological disabilities are increasingly recognised in older patients and again require multiagency community support.

Psychological problems are also common, as for many chronic diseases, and community support is important, particularly during the transition from paediatric to adult hospital care, when GPs offer continuity.³⁰

Acute complications of sickle cell disease			
Complication	Features	Treatment	Estimated incidence in SCA*
Acute abdominal sickling (girdle syndrome)	Severe pain, abdominal distension, constipation	Analgesia, laxatives, intravenous fluids	Fairly rare: 0.1 episode/patient year ^{10,15}
Cholecystitis	Increased incidence of gallstones from haemolysis	Surgical management with transfusion before surgery	Common: 30-50% of patients by age 30 ¹⁵
Fulminant priapism	Painful, unwanted erection lasting more than two hours	Analgesia, α adrenergic agonists, penile aspiration and irrigation	Common: 30% of men by age 30 ⁴
Macroscopic haematuria	Usually painless; associated with renal papillary necrosis	Conservative management with hydration; transfusion as necessary	Common: 20% adults at some point; also common in sickle cell carriers ³⁶
Acute visual loss	Caused by proliferative retinopathy or retinal artery occlusion	Urgent referral to ophthalmology. Possible exchange transfusion	Fairly rare: 1% of patients by age 30 ⁴⁰

*SCA=sickle cell anaemia (HbSS disease).

Nocturnal enuresis

Glomerular hyperfiltration and reduced sensitivity to desmopressin are detectable at 13 months in children with HbSS disease.³¹ This causes the production of large volumes of dilute urine and a high prevalence of nocturnal enuresis in older children with the disease. A survey of 213 children with HbSS disease in the US found that 42% of 6-8 year olds reported enuresis at least once a month, and 9% over 18 years still reported nocturnal enuresis as a problem.³² There is no good evidence on how to manage this. Advice about not drinking at night and waking the child during the night to pass urine is sometimes helpful. Desmopressin has been used, and a small study of 10 patients reported a 60% response rate,³³ although benefit has not been confirmed in larger trials.

Chronic pain

Pain occurring daily for more than three months is unusual in children with sickle cell disease. However, estimates suggest that chronic pain may occur in 5-10% of adults,³⁴ although accurate figures are not available. Chronic pain can be related to tissue damage, such as leg ulcers, avascular necrosis of femoral or humeral heads, or chronic osteomyelitis, and specific treatment of these problems is important. It is increasingly recognised that some patients report severe pain with no obvious cause. Such pain has some features of neuropathic pain and may be related to increased central sensitisation to painful stimuli. Neuropathic chronic pain is more difficult to manage but drugs such as gabapentin and psychological therapies may be helpful.²⁸ All types of chronic pain in sickle cell disease require multidisciplinary support, with the general approach being similar to other types of chronic pain³⁵; community provision of psychology and physiotherapy are particularly important.

Chronic renal disease

Blood is exposed to hypoxic and acidic conditions in the renal medulla, and in sickle cell disease this promotes HbS polymerisation and vaso-occlusion, which causes ischaemic damage. In addition, glomerular hyperfiltration is present in these patients from infancy, although the glomerular filtration rate falls progressively with age, contributing to renal failure in older adults.³⁶ Renal dysfunction is apparent from an early age—microalbuminuria is common in childhood and 20% of adults develop nephrotic range proteinuria. Chronic renal failure develops in 30% of adults.³⁶ A study of US mortality statistics

from 1999 to 2009 showed that chronic renal failure caused 7% of deaths, making it the third most common cause of death in sickle cell disease.¹ Six year survival was 70% in 106 adults with sickle cell disease undergoing renal transplantation in the US between 2000 and 2011, significantly higher than survival on dialysis.³⁷ Patients are monitored for microalbuminuria yearly and usually treated with a combination of hydroxyurea and an angiotensin converting enzyme inhibitor when this approaches the nephrotic range. Case series suggest that these drugs reduce protein loss, although it is unclear if this approach reduces the number of patients who develop renal failure.³⁸

Chronic leg ulcers

Chronic leg ulcers, typically just above the ankle, affect about 10% of adults with the disease in the US, but 50-75% in Jamaica, and are much less common in children. This difference may reflect the incidence of minor leg trauma. Leg ulcers are often painful. No treatments have a strong evidence base, but small studies suggest that many heal within six months with regular dressings and compression bandages. Other treatments that may accelerate healing include blood transfusion, skin grafting, hyperbaric oxygen, zinc supplements, arginine butyrate, and topical growth factors.³⁹ All these approaches seem promising in case reports or small series, but none has been validated in larger studies.

Proliferative retinopathy

Proliferative retinopathy (stages III-V) occurs in about 20% adults with HbSS disease and 50% of those with HbSC disease.⁴⁰ This can be progressive and lead to vitreous haemorrhage and retinal detachment, although visual loss affects less than 1% patients. Laser treatment may be beneficial for progressive retinopathy, but it is unclear whether routine retinal examination is beneficial. Warn patients about the importance of visual symptoms and refer urgently to ophthalmology if these occur.⁴¹

How can complications be prevented?

Prevention of infections

Penicillin prophylaxis is central to the management of children with the disease. In some countries this is continued throughout life, whereas in others it stops at age 5 years. Routine childhood vaccinations include protection against *H influenzae* type B and conjugated vaccines against *S pneumoniae* in most high income countries.

TIPS FOR NON-SPECIALISTS

Sickle cell disease is very variable, with some patients having severe symptoms every month and others being largely asymptomatic

Encourage adherence to penicillin prophylaxis whenever possible, and facilitate this by providing appropriate access to repeat prescriptions

Ensure that all patients receive recommended vaccines, including annual vaccination against influenza in many countries

Many episodes of uncomplicated acute pain can be managed at home with simple analgesia and community support

Ensure children and adults who were born overseas in high prevalence countries are offered haemoglobinopathy screening (full blood count, haemoglobin analysis)

AREAS FOR FUTURE RESEARCH

The value of aggressively treating all children with sickle cell disease with interventions such as hydroxyurea, regular blood transfusion, or haematopoietic stem cell transplantation to prevent organ damage and premature death

The value of simple commonly used interventions during acute admissions, including intravenous fluids, oxygen, thromboprophylaxis, and antibiotics

The development of specific treatments to shorten the length and severity of acute vaso-occlusive pain

The discovery and development of drugs designed to treat sickle cell disease, including those that promote haemoglobin F synthesis, prevent red cell dehydration, and inhibit HbS polymerisation

The validation of DNA, plasma, and imaging biomarkers that can identify infants at risk of severe complications early

The development of evidence based interventions in low and middle income countries, particularly African and Asian ones

ADDITIONAL EDUCATIONAL RESOURCES**Resources for healthcare professionals**

National Heart, Lung and Blood Institute (www.nhlbi.nih.gov/health/prof/blood/sickle/sc_mngt.pdf)
—American website with information on diagnosis and management of all aspects of sickle cell disease for high-income countries; free resource, no registration

NHS Sickle Cell and Thalassaemia Screening Programme (www.sct.screening.nhs.uk)
—Information on antenatal and neonatal haemoglobinopathy screening in England, with guidelines and standards on diagnosis, clinical management, and patient information leaflets; free, no registration.

Global Sickle Cell Disease Network (www.globalsicklecelldisease.org)
—Information on sickle cell centres and meetings across the world, with names of sickle cell disease specialists across the world; free, registration for full access

South Thames Sickle Cell and Thalassaemia Network (www.ststn.co.uk)
—Information on meetings, and access to guidelines covering diagnosis and management; free, no registration

Resources for patients

Sickle Cell Society (www.sicklecellsociety.org)
—Website of UK Sickle Cell Society with news, information, and patient leaflets; free, no registration

NHS choices (www.nhs.uk/conditions/Sickle-cell-anaemia/Pages/Introduction.aspx)
—Patient information on NHS screening programme for sickle cell disease; free, no registration

Sickle Cell Disease Association of America (www.sicklecelldisease.org)
—American website with news, guidelines, and information; free, no registration

Rofsed (www.rofsed.fr)
—French language website for children, parents, and professionals; particularly good for teenagers; free, no registration

Children should also receive unconjugated pneumococcal vaccine from 2 years of age, repeated every three to five years, and immunisation against meningococcus, influenza, and hepatitis B.⁴²

Prevention of acute vaso-occlusive complications

Many patients report exposure to cold or dehydration as precipitants of acute pain. Studies in tropical countries suggest increased frequency of acute pain during the rainy season, whereas in the northern hemisphere windy weather is associated with increased hospital admissions.⁴³ Advise patients to dress appropriately for the weather and ensure adequate hydration. It is important for children and adults to be physically active, although sports involving prolonged exertion or exposure to cold may provoke vaso-occlusion. Advise patients not to ski as anecdotally this can precipitate acute complications, because of cold temperatures and low oxygen levels at high altitude. Flying on commercial airlines is well tolerated and does not usually require prophylactic blood transfusions or supplemental oxygen, unless the patient has pre-existing hypoxia or cardiopulmonary problems. Patients should not fly if they have acute pain or other acute illness.

Hydroxyurea is the only drug that reduces the frequency of acute vaso-occlusive complications. In a double blind randomised controlled trial, 152 adult patients with HbSS disease were given hydroxyurea and 147 assigned placebo. Hydroxyurea was associated with a significant reduction in episodes of acute pain (median 2.5 v 4.5 episodes/year; $P < 0.001$), acute chest syndrome, and blood transfusion.⁴⁴ Subsequent trials showed similar reductions in pain frequency in children, but provided no evidence that hydroxyurea protects against organ damage, despite observational studies suggesting reduced mortality.⁴⁵ In Europe, hydroxyurea is usually started if more than two episodes of acute pain occur each year, or after a severe episode of acute chest syndrome. Use is limited, partly because of theoretical concerns about the side effects of a cytotoxic drug and the potential increased risk of cancer and subfertility. These risks are unquantified, but seem to be small. In England, 15–30% of patients are currently taking hydroxyurea,⁴ and it is often stated that it is underused. In the US, hydroxyurea is more widely prescribed for a wide range of indications, including hypoxia, severe anaemia, pulmonary hypertension, renal impairment, and cerebrovascular disease.⁴⁶

How should pregnant women with sickle cell disease be managed in the community?

Women with sickle cell disease have near normal fertility. There has been concern that hormonal contraception might increase the risk of acute complications and venous thrombosis in women with sickle cell disease. Although no large studies have looked at this population specifically, a recent systematic review of eight non-randomised and cross sectional studies found no evidence that hormonal contraception was associated with an increased risk of complications in sickle cell disease, although there were no studies of venous thrombosis.⁴⁹ This review found that progestin-only and combined hormonal contraception were safe in women with sickle cell disease.

A large database study of 18 000 deliveries to women with sickle cell disease in the US showed that all complications were increased compared with controls. For example, the odds ratio was 2.5 (1.5 to 4.1) for deep vein thrombosis, 2.2 (1.8 to 2.6) for intrauterine growth retardation, and 6.8 (4.4 to 10.5) for sepsis.⁵⁰ Complications of sickle cell disease are also increased, with 50% of women developing episodes of acute pain.⁵¹

Women with sickle cell disease should receive pre-conceptual counselling about prenatal diagnosis and the risks associated with pregnancy; prenatal diagnosis should also be discussed when they are first seen during pregnancy. Hydroxyurea, which is potentially teratogenic, should ideally be stopped three months before becoming pregnant, and definitely when pregnancy is suspected or confirmed.⁵² Pregnancy outcomes depend greatly on the facilities and expertise available. There is no established evidence base on managing pregnant women with sickle cell disease, although regular blood transfusions are typically used in

those with a severe clinical course, such as those developing frequent episodes of pain, acute chest syndrome, or those with previous complicated pregnancies. Women should be referred to an obstetric team with experience of high risk pregnancies, and followed in a specialist sickle cell disease pregnancy clinic where available.⁵³

Penny Ryba reviewed and commented on the article. Megan K Jones originally drew figure 1.

Contributors: All authors helped plan the article, wrote parts of it, and reviewed and edited the final version. DCR is guarantor.

Competing interests: VB received funding from Novartis for travel to meetings but does not act on behalf of any particular group or lobby. JM has no competing interests. DCR is a consultant for Eli Lilly for a trial of prasugrel in sickle cell disease and principal investigator for this study in the UK. He is also principal investigator for a study of Prevenar in sickle cell disease funded by Pfizer. He has received funding from Novartis for travel to meetings, and has an unrestricted educational grant for £2500. He is an expert witness in one or two medico legal cases a year but does not act on behalf of any particular group or lobby.

Provenance and peer review: Commissioned; externally peer reviewed.

References are in the version on bmj.com.

Online State of the Art reviews: Drug treatment of epilepsy in adults

The second in our series of “State of the Art” reviews is drug treatment of epilepsy in adults (<http://www.bmj.com/content/348/bmj.g254>). About 75% of adults with new onset epilepsy will become seizure free with antiepileptic drugs although around half of them will experience adverse effects. The rest will have drug resistant epilepsy with seizures, adverse effects, increased mortality, and comorbidities.

The review summarises the evidence on when to start treatment, how to select the first antiepileptic

drug, and how to optimise treatment on the basis of each drug’s benefits and harms. The adverse effects of each drug are presented in a colour coded figure making it easy to compare drugs side by side. The drugs’ mechanisms of action are also illustrated (figure).

Management of the 20-30% of people with drug resistant epilepsy is discussed including how to tailor treatment in special patient groups, for example pregnant women and people with comorbidities.

RESPONSE TO ANTIPILEPTIC DRUGS

