either amounting to actual fits, or only to an aura epileptica.

The following are two of some instances which I have seen of epilepsy of the retina.

CASE XI. Stephen P., aged 22, had had two years before I saw him an attack of epilepsy, of which (he said) the effects lasted for three days. On recovering, he found the left eye closed (ptosis), and he could not open it for seven days. From that time, he had had no further fits. He did not remember his sight failing him for any time, or immediately before the fit. What he now complained to me of was, that about once a week or fortnight he would lose his sight for a time, but that he recovered it again. I found that he was amblyopic with the left eye, and appeared to be slightly myopic with both. With the ophthalmoscope, I detected neither a hypermetropic nor a myopic refraction of the eye; the optic discs were reddish, their contours rather ill-defined; the vessels were rather small; there was an appearance of tension in the internal tissues.

CASE XII. A man, aged 21, a year ago, whilst at work, found a "mist" come over his eyes sufficiently dense to obscure objects around him. He has ever since been subject to these attacks every four or five weeks. They last for from five minutes to half an hour at a time; the average being a quarter of an hour. During an attack, he can see his way about, and even read with difficulty. He has never had any fits, but occasionally feels giddy. After one of his attacks, he feels sickish and has a headache.

I am quite aware how imperfect is the above sketch that I have ventured to offer on these obscure functional impairments of the retina. It is necessarily so, from the comparatively slight amount of attention the subject has hitherto commanded. Imperfect as it is, I trust, however, that it may prove sufficiently suggestive to induce others to follow up what I cannot help considering as a most interesting and important study, not only for the ophthalmologist, but for all who are interested in the investigation of disease generally.

CARCINOMA OF THE STOMACH.

By JOHN RICHARD WARDELL, M.D., M.R.C.P., Physician to the Tunbridge Wells Infirmary.

[Concluded from page 639.]

To attempt to assign any real or regular cause of cancer, would be at the best but a vain endeavour; and all that we can assert, in the present state of our knowledge, are mere hypotheses and vague surmises, which are either not borne out, or are absolutely refuted, by the facts which accumulated cases present. We know, however, that some occult cause, whatever it may be, effects a very potent change in the fluids; that it institutes some morbid condition of the blood capable of favouring the genesis of those flagrant cell-growths known as malignant. We see this condition in persons whose bodily conformation and external appearances are most opposite; sometimes in the ruddy and muscular, as well as in the pale and attenuated. The belief of many that it is an hereditary disease is a creed which has long been entertained, and one that is becoming still more prevalent; for certain it is that, in many instances, other members of a family are, upon inquiry, found to have died of the same disease. Nor is the acknowledgment of this transmission by inheritance at all inconsistent with what we know of the vital properties of the organism. Absolute cancerous material cannot be transmitted with the germ; but some hidden, inscrutable impress is transmissible, which impress at a remote period favours the development

of that product; and we must confess that it cannot be entirely divested of any material relations. We know that other diseases, as tuberculosis, gout, insanity, are hereditary; and there is no reason why the cancerous diathesis should not be transmitted in like manner. That constitutional tendency is, however, very greatly promoted or retarded by the operation of external agencies; and there is no doubt that a high state of civilisation conduces to its progress and results. With the exception of the uterus, there is no organ so prone to cancer as the stomach; numerical data in our own and other countries having abundantly attested this fact. Men are more decidedly prone to it than women, probably on account of their more intemperate habits, greater exposure to vicissitudes of temperature, and greater mental anxieties. Some few instances are given of its occurrence before the age of thirty. It is, however, very seldom observed before the age of thirty-five; far more frequently in those approximating fifty, and in still more advanced life. It belongs to those heterogeneous changes in the assimilation of the tissues which proclaim declension of vital power, and are intimately associated with age. There is no reason for believing that one class is more prone to it, or enjoys a greater immunity from it, than another. We meet with it in persons occupying every social position—in the well-nourished, as well as in the half-starved; in those who live in the country, as well as those who live in cities.

In the great majority of examples, it is primary cancer which affects the stomach. This organ may, it is true, be the seat of the secondary form; but such cases are mere exceptions to a great general rule. The lymphatic glands may become contaminated by the presence of the disease in neighbouring viscera; as, for example, when the head of the pancreas, the liver, the spleen, the omentum or mesentery, are affected. Some pathologists have affirmed that it is the colloid variety which is most generally found in the stomach. It would, I think, more correctly express the fact, if we say that it is the scirrhus-colloid which is most frequently met with in that particular situation—the hard, fibred basic substance upon which is superimposed the gelatiniform, locular mass. Colloid may and does coexist with the villous, encephaloid, and melanotic forms; but these combinations are not nearly so often observed. The carcinomatous matter is infiltrated into the areolar tissue; because, in such loose and comparatively unresisting structure, it there finds less opposition to its deposit. In the course of time, the muscular and other tissues become encroached upon, and cancerous cells are formed within the muscular filamentous sheaths, the molecular constituents of the muscles being absolutely displaced and occupied by the new formations. One of the great characteristics of carcinoma is displacement and occupation. The product which is substituted not only does not possess the secretory capabilities of the parts removed, but it gravely interferes with the functions of the organ or organs which it has selected as its *habitat*. Again, the vitiated secretions, which are poured out into a hollow viscus, confer additional disorder. Carcinoma of the stomach fully illustrates this evil. The large jelly-like mass, by mixing its perverted exudations with the gastric juice, so injuriously operates upon the normal qualities of that fluid as to render it quite unequal to the due performance of its office; hence one cause of the sour ejections, the dyspepsia, the occasional attacks of diarrhoea, and that gradual diminution of flesh and strength, which an impaired chyme must inevitably produce. The encephaloid variety grows most rapidly; the scirrhus most slowly. Scirrhus and colloid may exist in the stomach for

years before they destroy; the encephaloid will produce a fatal issue in the course of a few months. The one typifies the acute, the other the chronic form. The encephaloid and scirrhous are considered to give the most pain; the colloid the least. The loculi or alveoli, which are so characteristic of the colloid, are formed simply by irregular infiltration; they vary in size and configuration, according to the circumferential pressure, the amount of fibrous tissue, and their deep or superficial position. In the stomach and peritoneum, the colloid mass is more diffuse and less nodular than when deposited in glandular structures. Microscopic investigation has in latter years cleared up much that was previously obscure respecting the ultimate structure of cancer; and, by this acquired knowledge, diagnosis has been rendered much more certain than it was when mainly based upon clinical observation and empirical practice. Cancer-cells are regarded by Lebert as modified lymph-cells, often monstrously altered in size. Collis says they present more or less resemblance to typical forms; and in any specimens there will be much variation in the size and outline of individual cells, yet with much elementary similarity in the tumour. Lebert describes the colloid cells as being large, pale, oval, round, or tubular, lying in clusters. There are also small granular irregularly shaped corpuscles, which are regarded as cancer-cells hindered in their development. The pyloric end of the stomach is most prone to carcinoma; next in frequency, the cardiac orifice; after that situation, the lesser curvature. The splenic end is least liable to the affection. It is a curious fact—nevertheless, one which is true—that the deposit is rarely or never seen to extend into the duodenum; and the only reason why this abrupt termination thus obtains appears to be, that the duodenum is far more scantily supplied with areolar tissue than the stomach. The mass may present much resemblance to effused lymph; and it should not be forgotten that lymph will sometimes be deposited in the walls of the stomach, especially towards the pylorus, to considerable extent. But the malignant growth is much less uniform in its configuration; and there is not superficial softness in mere lymphic deposit. When the disease pervades the cardiac orifice, the growth will extend into the œsophagus, and cause organic stricture. When the liver becomes secondarily affected, the germs are transferred by the lymphatics and the veins, especially by the latter; because, it should be remembered, the gastric veins run to the liver. The reason why the lungs are not implicated in carcinoma of the stomach, is supposed to be on account of the cancer-cells being too large to pass through the hepatic lobular plexus. When the liver has become carcinomatous, sometimes unevenness and pitting may be felt on its surface. I have known this in marked manner when there has been much emaciation. The stomach is often found agglutinated to the liver, and more especially when the disease is at the pylorus. When the growth at the pyloric orifice becomes considerable, the pylorus will fall into the right iliac fossa, simulating tumour or impaction at the ileo-cæcal junction. I recently saw a case which I believe to be very illustrative of this fact. Rokitsansky has known the pylorus to touch the symphysis pubis. We have seen in Case iv above recorded, that even when the mass hangs from the lesser curvature, the pain may be chiefly referred to the right fossa. This is a fact which, in a practical point of view, is of much importance. This falling down can, of course, only be when there is no attachment to neighbouring viscera. On reference to Case iv, it is stated that the patient was always in the least pain when in the

erect position. The autopsy fully explained this peculiarity. The mass was pressed upon in the sitting or lying postures; but, on the resumption of the erect position, it would hang tolerably free in the gastric cavity.

The symptoms by which this formidable disease can be recognised are, in the earlier date of the affection, often obscure, and by no means easy of interpretation. When it has made progress, the diagnosis becomes comparatively easy. Sometimes it happens that the complaint has considerably advanced before it was even suspected. Again, it occasionally occurs, as Dr. Watson has remarked, without from first to last presenting any pathognomonic symptoms at all. It usually begins with dull, aching pain at the epigastrium, sour eructations, and an uneasy feeling of fullness and distension, which are at first merely attributed to indigestion; afterwards, there are loss of appetite, depression of spirits, and, curiously enough, an unwonted petulancy and irritability of temper. The tongue is comparatively clean; nor is there any symptomatic fever. Costiveness, alternating with occasional attacks of diarrhœa, supervenes, doubtless caused by undigested food and that butyric fermentation which the unhealthy secretions from the cancerous surface are known to produce. No symptom, however, augurs greater import than the loss of flesh; and, if accompanied with the yellowish dusky cachexia, in addition to the more obvious and ordinary symptoms, the prognosis must needs be most unfavourable. When the ordinary stomachic remedies fail, and the wasting goes on, there can be but little doubt of malignancy. As the complaint advances, vomiting generally comes on. In two of the foregoing cases, it came on immediately after eating. In both, the disease was at the cardiac end of the stomach. In Case ii, where there was partial stricture of the œsophagus, the food was instantly rejected. In all the cases, the ejections were invariably sour. Sometimes there will be mixed with the glairy mucus black or coffee-grounds looking flakes, the results of hæmorrhagic exudation, or the erosion of some of the smaller vessels; or there may be active hæmorrhage. This symptom is in the latter stage. At the first, simple ulcer may be simulated; but, in simple ulcer, the pain is not so persistent, nor so sharp and lancing; it occupies but a small area; and the rest of the mucous membrane is healthy, and there is little or no loss of flesh. When the orifices are not obstructed, sickness is less urgent, and the patient lives longer. When the pylorus is the seat, a special train of phenomena supervene. This orifice becomes gradually narrower until positive occlusion well-nigh results; the imperfectly digested food can then but with great difficulty pass through the contracted portals; the stomach labours to overcome the obstruction; and the applied hand can feel a vermicular movement resulting from its efforts. By this excess of action, and the abnormal gases which are generated by perverted secretion, the organ becomes large, and percussion elicits preternatural resonance. When there is constricted pylorus, the intestines are empty, and the belly drawn in—a condition which is seen in some other forms of disease; for example, in typhus, when there is diminished chyle and great deoxidation of the tissues. When the cardiac orifice is the seat of the tumour, the swallowed morsel gives rise to pain immediately after its descent into the œsophagus. There will then be instant rejection, or (Case i) the food will feel to stick as it were at the epigastrium; and the pain to which such obstruction gives rise is excessively severe. When the growth is in the lesser curvature, it very generally extends to one or both orifices. In Case iv, it did not do so. Sickness may

be a distressing symptom even when the inlet and outlet are free from disease. The mental faculties usually keep clear to the last. In all of the above examples, this was the case.

I have said the diagnosis is more difficult at the first than afterwards. When the affection has progressed, palpation will assist the formation of an opinion very materially. Pressure gives pain at some circumscribed spot. This pain often radiates through into the back. Frequently, a large, irregular, hard substance can be felt in the right epigastric region, between the mesial line and the right false ribs. The movement of this substance gives pain. When the cardium is the site of the growth, some thickening can be discerned (after emaciation has progressed) towards the splenic end of the organ. The patient's description of the pain should never be disregarded. Sharp, stabbing, hot, burning, were terms employed by the patients whose cases have been given above. We should bear in mind that the pain may simulate that of renal calculi, of ordinary hepatic pain, of tumour in the posterior mediastinum, of abdominal aneurism, of lumbago. There may, however, be but little pain, as sometimes little vomiting, from first to last. I have known carcinoma of the uterus be attended with scarcely any pain. If the branches of the par vagum enter the mass, there will be great suffering. If the tumour lie in front of the abdominal aorta, it will be lifted up by the pulsations, and simulate aneurism. Negative and stethoscopic facts will, however, guard us against such a mistake. In chronic or acute irritation of the gastric mucous membrane, the pain is diffuse, and referred to the entire epigastrium—not circumscribed as in malignant tumour. The ejected matters will assist in some measure in arriving at a correct diagnosis, not only as regards the true nature of the disease, but as to its particular position in the organ. Under the microscope, cells of a specific character can, on careful examination, be discovered mixed with the vomited fluids. If the growth be at the pylorus, the food will be partially digested; if at the cardium, it will generally be little altered. If there be much blood, we should suspect the encephaloid or melanotic variety. Notwithstanding all the foregoing rules for observation, necroscopy will from time to time reveal conditions unsuspected, and disappoint us by not displaying those which we had anticipated.

Some time ago I was requested by Mr. Hutchinson of Lamberhurst to see with him a patient who had long been affected with persistent stomach-disease. There was no very decided cachectic expression; yet he got thinner, and his condition caused anxiety. The tongue was livid, flabby, smooth, creamy; the appetite impaired; the epigastrium full and resonant. Pressure gave pain, which was *not localised*, but diffused over the whole region. There was pain after meals, but no vomiting. The urine was voided in normal quantity; specific gravity 1015; excess of triple phosphates. I gave it as my opinion, notwithstanding the loss of flesh and other doubtful symptoms, that his case was not malignant. Mineral acids, bitter infusions, morphia at bedtime, and a regulated diet, were followed by gradual improvement; and for more than twelve months he has been quite well. An in-patient was admitted into the infirmary under my care several months ago, in the person of a tall, powerful woman, who had been servant in a gentleman's family. For more than half a year she had had epigastric pain, and more especially after eating. She looked sallow, anæmic; and had become thinner. Her spirits were much depressed, and she believed she laboured under a fatal malady. A medical opinion had been given sus-

pecting cancer. I was written to by the lady with whom she had lived, respecting her going into the Cancer Hospital. The epigastrium was rounded and resonant; no tumour; no hardness; and pressure gave diffuse dull pain, which did not lancinate into the back. Tongue creamy, flabby, not loaded. Right hypochondrium duller than normal. I expressed my decided opinion that her affection was not cancer. A blister to the epigastrium, a mixture with infusion of rhubarb and calumba and the bicarbonate of soda, with small doses of strychnine, mild mercurial alteratives, and an easily digested and nourishing diet, were followed by speedy improvement. She soon left the hospital well, and her health continues good at this date.

In saying a word relative to treatment, there cannot be much discrepancy of opinion upon this head. When we are impressed with the belief of the existence of malignancy, the remedies can only be palliatives, or at most such as are deemed to possess the power of retarding the progress of the complaint. All depressing mental emotions should be avoided, as well as every description of ingesta which might irritate the lining membrane of the stomach. The food should be light, digestible, and nourishing. Solids are to be utterly discarded. The various farinaceous articles of diet are to be taken; and these may be advantageously given in combination with soups, broths, and jellies; new laid eggs, milk (mixed with lime-water when there is acidity), pounded meats, and like bland articles of food. With regard to medicines, the nitrate of bismuth, as a mineral astringent, has been much lauded. It may be given in combination with hydrocyanic acid and some bitter infusion. When there is excess of acidity, the aqua calcis is preferable to the other antacids. If this be not employed, Brandish's alkaline solution, with some aromatic water, will be found serviceable. Morphia, henbane, and conium I have found of great service, even at the earlier stage, in tranquillising the nervous system and giving sleep. When the disease shall have advanced, morphia is the all-important remedy. When sulphuretted hydrogen is developed, the creasote pills produce much benefit. The bowels may be regulated by pills of aqueous extract of aloes, extract of belladonna, and quinine. An opium and belladonna plaster to the stomach has a comforting effect; and towards the close, when the pain becomes excessive, opiate epithems are not to be omitted. When the stomach will not retain food, nutritive eucmata may advantageously be employed.

DANGER OF INJECTIONS INTO ANEURISMS. A man in the Aix Hospital had a traumatic aneurism of the brachial artery at the elbow-joint. Compression was tried for a month, but failed. M. Goyrand therefore injected into the tumour five drops of perchloride of iron. The aneurism at once became firmer, the thrill of it ceased, the pulsation in it was less, but still continued. A second similar injection was therefore tried. Immediately thereupon the whole hand became pale; the sensation in the fingers was most painful; the temperature was lowered. The hand was like that of a dead person. All pulsation of the tumour had ceased. Frictions and hot applications were applied to the hand. On the next day, a blue circle was seen at the finger and gradually increased. Deep pain was felt in the hand. Eventually the hand was separated at the wrist by dry gangrene, and the man left the hospital cured in a month. The accident was accounted for by supposing that a clot of fibrine had been driven into the ulnar and radial arteries.