himself, each one for his brother, must work it out *here*. Duly used, this society is a rock of salvation against those sins of the heart which so easily beset us—mutual distrust and mistrust and jealousy, breeding discord and the strife of tongues; and duly used, the county hospital is a way of salvation from those sins of the head—intellectual apathy, complacent self-sufficiency, and the careless habits into which even the best of us are only too prone to fall.

Clinical Lectures

ON

ANATOMY AND MORBID ANATOMY IN CLINICAL SURGERY.

DELIVERED AT ST. BARTHOLOMEW'S HOSPITAL

BY

C. B. LOCKWOOD, F.R.C.S.,

SURGEON TO THE HOSPITAL.

LECTURE I.

GENTLEMEN,—The subject of my first lecture is anatomy in clinical surgery. Anatomy is the foundation of all surgery, and I shall try to show its bearing upon clinical surgery. It is impossible for me to impress too deeply upon you the importance of a thorough knowledge of human anatomy. If you do not already possess it, I urge each one of you to study a small fragment every day. He who will read, mark, learn, and inwardly digest a single bone in the course of a week will find that, after a while, he has a splendid equipment.

I was led to choose this subject by an incident which happened in the wards but a few days ago. A youth had had his hand crushed by machinery. He had lacerated wounds on the back and in the palm of the hand. The index finger had been seriously damaged. In place of the proximal interphalangeal joint there was a gaping wound, with bone protruding. When this state of affairs was seen I naturally tried to form an estimate of the extent of the injury, and of the subsequent utility of the hand. I therefore asked the dresser in charge of the case—a gentleman who works hard and is most earnest and attentive in the performance of his duties—whether the patient could feel with his finger and thumb. This point had not been investigated, and yet the youth could not feel with either finger or thumb. Consider for a moment what consequences this omission might entail. There is now an Employers' Liability Act, and under it the patient is entitled to compensation for the damage he had sustained. But if the medical man under whose charge he happens to be is not acquainted with the extent of the injury, obviously the sufferer may not obtain a suitable recompense. Then there is the employer. He is entitled to a just and accurate report, for upon that the extent of his liability may depend. But the consequences of this omission may also be very disastrous to the medical man in charge of the case. If as witness in a court of law—and every one of you will have to testify in courts of law—he fails to display adequate knowledge, it will be harmful to his reputation. At the present moment, however, I am more interested in the psychology of this case than in its surgery; I am concerned to know why the dresser's mind did not start a proper train of thought, leading on to correct knowledge of the extent of the injury. He was not thinking anatomically. Had he thought anatomically he would have passed in review each structure. He might have proceeded from the surface of the depths of the hand, or he might have considered the structures in the order given in a familiar textbook, such, for example, as *Gray's Anatomy*. Then he would have begun by asking what injury the bones had sustained. The other day a medical friend had his hand crushed. His case was considered in anatomical order. The question of an injury to the bones was raised. He hardly seemed to appreciate the desirability of an x-ray photograph until one had been taken, and showed that two of the metacarpal bones were broken. He held an accident insurance policy. So the bones will

be passed in review, and photographed on the slightest possibility of damage. Next come the joints. An injured joint should not be overlooked: broken into, full of blood, inflamed, it may be stiff unless the injury has been recognized and movements begun early. Then come the muscles and tendons: divided tendons may require suture. A hand with ununited tendons is not of much use. Continuing the order in Gray, pass in review the arteries, veins, lymphatics, and nerves, and, where the trunk is concerned, the viscera; the cranium, with the brain and special organs; the spinal column and its nervous contents. If, gentlemen, you investigate every injury in due anatomical order, it is not likely that you will overlook any grave defect. Injuries to the limbs sometimes necessitate amputation. In deciding this question proceed again in anatomical order: estimate the amount of destruction to the skin, fascia and muscles, bones and joints, damage to the great vessels and nerves. The leg will gangrene if the superficial femoral vessels be torn. There are hardly any other channels than these to supply the lower limb with blood. I recall a little tragedy which took place very suddenly and very quietly. A man had a popliteal aneurysm. The surgeon exposed the superficial femoral artery, and when the aneurysm needle was passed between it and the vein a thin stream of dark blood ran slowly but steadily from the wound. The artery was tied, but those who saw that stream of blood knew that the vein had been injured and that the leg would gangrene, which it did. To cut off nearly all the arterial supply from the lower limb is a critical thing to do, but to injure the vein at the same time so that it fills with clot, thus obliterating the only channel for the return of the blood, is an overwhelming disaster. Then the great nerves should be passed in review. Complete division of the great sciatic nerve, taken with the other injuries, might lead you to decide upon amputation. But remember that sensation may be absent in a limb whose main artery has been damaged.

The kind of anatomy which is required in clinical surgery is not minute, but at the same time entails a correct knowledge of the limits and extent of the various cavities of the body. Given a swelling in the region of the knee-joint, the first thing to decide is whether it be inside or outside the capsule, or whether both inside and outside, with a communication through an opening in the capsule. A swelling outside obscures the anatomical structures on the outside of the capsule—such, for example, as the tendons on the inner or outer side of the knee, or the patella. But a swelling inside distends the capsule and lifts it up, together with the structures lying upon it. If you are properly acquainted with the limits and attachment of the capsules of the joints, it will help you very much to decide these rather difficult questions. After having made up your mind on anatomical grounds where the swelling is situated, then you will decide whether the swelling is fluid or solid. If fluid, you will try to decide the nature of the fluid; and if solid, whether it is the result of inflammation or of new growth.

Anatomical thinking will help you to avoid the pitfalls which beset injuries to the head. We often see a paragraph headed "Drunk or dying?" The history is usually as follows: Some one is found insensible upon the pavement. It is not known how he got there. He may have fallen, he may have been knocked down. At all events he is insensible and suffering from alcohol, or disease, or may be concussed. Concussion is a state of insensibility accompanied by shock. The boxer who is knocked out is suffering from concussion. Under proper conditions of rest and warmth, the concussed patient recovers consciousness, and proceeds to his home, or it may be to prison. There he gradually becomes drowsy, then unconscious, and dies of compression. Whenever the bony cavity in which the brain resides is suddenly encroached upon by fluids or by solids the brain ceases to perform its vital functions, and unless the pressure of the fluid or solid is taken away death ensues. This is what happens in these cases of "Drunk or dying." The injury was enough to cause concussion, but at the same time lacerated a vessel inside the skull. The blood accumulated within the skull, diminished its capacity, and compressed the brain. Now, such cases as this cannot be overlooked by those who have learnt to think in anatomical order. Every case of concussion calls for the most grave consideration, and my

house-surgeons know that if they like they can fill every one of my beds with trivial cases of concussion rather than send away one with commencing compression. Now, how does anatomical thinking prevent you from falling into error? He who thinks anatomically will investigate the patient's nervous system in order. I suppose that he may not learn much from his investigation of the olfactory nerves. My observations lead me to suppose that this must be a singularly defective sense in most people. But the second cranial nerve, the optic, may throw great light. After an injury to the head it would be, to say the least, somewhat feeble to overlook the fact that the patient had become blind in one eye. Then with regard to the third and other orbital nerves, what shrewd inferences can be drawn from a dilated pupil! Or from a slight squint. The auditory and facial would not be forgotten. The fauces would be examined, and last, the movements of the limbs. I have seen a man lying in bed with a partial paralysis of his left leg. This I suppose had been overlooked because his leg was underneath the bedclothes and because anatomy had not been applied to clinical surgery.

I have already referred to the cavities of the body, and said that it is necessary for you to know their extent and topography. I will try to tell you my own crude way of thinking over the topography of the cavities of the body, such as, for example, the antrum of Highmore, or the tympanum. There used to be a book on anatomy, written by the late Erasmus Wilson, who was such a benefactor to the College of Surgeons. And in this book, in giving the relations of the great arteries to their surroundings, the vessel is drawn with a square outline. I expect your minds are much the same as my own, and if you try to imagine the relation of one circle to another you may find it exceedingly difficult to determine whether it ought to be in front or to one side. But if you have in your mind a square, or a cube, it becomes easy to determine whether a given structure is or is not round the corner. Now let us apply this to a clinical case, such as that of a tumour growing in the antrum of Highmore. Thinking anatomically, and considering the antrum to be a cube, you will say, Does it bulge in front? This will lead you to look inside the mouth, and examine the front of the antrum. Next you will ask, Does it bulge at the back? Well, at the back of the antrum is the sphenomaxillary fossa, with the lower divisions of the fifth nerve. This will lead you to investigate your fifth nerve. Then you will ask, Does it bulge at the outer side? So you look for a swelling in the zygomatic fossa. At the inside you will not forget the nose, which may be stopped up. Below? You will look into the mouth. Above? You will examine the orbit.

Consider the tympanum in the same way. Doubtless you will begin at the outside of the cube, and investigate the external ear and tympanic membrane. Next, doubtless, you will examine inside where the internal ear is. So that you will not forget to find out whether or not the patient is deaf. The roof is a thin layer of bone and dura mater between the tympanum and the inside of the skull, and upon it rests the temporo-sphenoidal lobe of the brain, to which septic infection so often spreads. And, lastly, you will consider the front and back, and remember that at the back of the tympanum, besides the mastoid cells, are the sigmoid sinus and cerebellar fossa of the skull.

I have occupied you for some time with a subject which, although of intense practical importance, is at the same time apt to be dull and tedious. But I will not cease without referring to the anatomy of the trunk. And here I would give you a few words of advice. Learn the broad facts of topographical anatomy before you essay the minute; learn to crawl before you walk, to walk before you run. In some works on topographical anatomy the trunk of the body is mapped out like a chess-board, interesting, perhaps, to those whose memories may surpass their understanding, and whose lives are spent in the dissecting room, but not to those who have so much to learn. Let us suppose the clinical case of one who has a pain beneath the right costal margin at the upper end of the right rectus abdominis muscle, with abdominal tenderness and rigidity, vomiting, and perhaps constipation. When I demonstrated and lectured upon anatomy, I advocated the use of a few surface centres. By that I mean a centre in the surface of the body beneath which important structures are grouped. Beneath the

eighth right costal cartilage is one of the important centres of the abdomen, for its centre marks the situation of the foramen of Winslow. Need I enumerate the structures grouped round the foramen of Winslow? The finger upon the centre of the eighth costal cartilage has above it the transverse fissure of the liver and all the structures in it; it has the pylorus and the head of the pancreas below it. A little to the right is the gall bladder, the gall bladder lying on the first part of the duodenum. When the gall bladder is inflamed the patient often vomits. And next, just below the tip of the gall bladder, is the transverse colon. So when an inflamed and distended gall bladder rests upon the transverse colon, flatulence and constipation ensue. Another important surface centre is a point in the right semilunar line, where it is cut by a line drawn from the umbilicus to the anterior superior spine of the ilium. Here is the ileo-caecal junction with the ileo-caecal valve, the end of the ileum, the lower part of the mesentery, and the structures in it, the caecum, the beginning of the right colon, and the vermiform appendix. I trust we all have our own ways of thinking, but I suggest surface centres to you as one of the ways of bringing your anatomical knowledge to bear upon your clinical surgery. Doubtless you will discover the other surface centres for yourselves. When I taught anatomy about nine sufficed.

Anatomical thinking is of especial usefulness in the investigation of diseases of the alimentary canal, or in connexion with those of the genito-urinary tract. Given a case in which blood or pus is present in the urine, your thinking would begin at the meatus urinarius, and pass along the urethra. I have known an ordinary gonorrhoea mistaken for nephritis because there was albumen in the urine, the urethra not having been examined. Next, the membranous urethra, and the prostatic. Remembering the prostate, doubtless a rectal examination will be made. Remembering the openings into the prostatic urethra, the vesiculæ seminales will be felt, and the mind follow the course of the vasa deferentia, and the testicles will be examined. Otherwise a tuberculous mass in the epididymis might be overlooked. Next the bladder will be considered, the ureters, and lastly the kidneys. A patient with much blood in the urine was supposed to have a growth in the bladder. And so she had, but something else in addition; for, when the genito-urinary tract was examined in anatomical order and the kidneys felt, a large tumour was discovered in the right kidney. It had been overlooked. I have referred to a rectal examination. In carrying out this you must, as Luther Holden used to say, carry your anatomy at the tip of the finger. Examine the anal aperture methodically, so as not to overlook a pile or fissure or fistulous tract. Then the front of the rectum, feeling for the membranous urethra, each lobe of the prostate in turn, the vesiculæ seminales, the trigone of the bladder, and, last of all, at the tip of your finger, the recto-vesical or Douglas's pouch of peritoneum. Anatomical thinking would also lead you to examine the hollow of the sacrum, where enlarged lymphatic glands might be felt.

I shall now endeavour to show you some of the uses of morbid anatomy in clinical surgery. But do not think that there is any hard-and-fast line between anatomy and morbid anatomy.

Here are specimens in bottles, not very satisfactory, because they cannot be taken out and handled. Let us proceed on anatomical lines, beginning with the bones. Fractures near the ends of the long bones will be a source of great trouble and anxiety. Here is a specimen of a fracture called after Colles, an Irish surgeon. Of course you are familiar with Colles's fracture. It is a transverse fracture of the lower end of the radius. Morbid anatomy teaches that the fracture is usually impacted. What ought you to do with that impaction? Are you going to leave the impaction, or are you going to reduce it? I do not know what you are taught, but it is, to say the least, impolitic to leave one of your patients with a deformed wrist. An anaesthetic ought to be given, and the deformity put right. Have you been thinking anatomically? There is something else about Colles's fracture. The various things at the lower end of the radius will have passed through your mind, and the wrist-joint not forgotten. What happens to the wrist-joint when there is a fracture so close to it? Whenever there is a fracture in the vicinity of a joint its cavity may have been broken into. Mr. D'Arcy Power examined a large number of specimens

of Colles's fracture, and found nearly all went into the wrist-joint. A joint with a fracture into it may not only undergo a mechanical alteration but is also filled with blood. Blood in a joint causes an attack of synovitis, and that will persist until all the blood has gone, and perhaps afterwards. When you have made up your mind that the fracture has gone into the joint you will take measures to prevent that joint being stiffened by adhesions, and movements begun as soon as the pain allows. Here is another specimen of a fracture associated with this hospital. It is named after Percival Pott, one of the ablest of our surgeons. Go to the library and read the preface to Pott's works. He asks a rather curious question: "Why are people so fond of going to quacks? And why, even when the quack has maimed them, do they send their friends?" We may suppose that if a person has got well after having been to a quack he boasts about it as a tribute to his intellectual acuteness in choosing the right quack. If maimed, he sends somebody else, perhaps on the principle of the fox without a tail, or as a subterfuge to hide his own misfortune. Pott's fracture is a very grave injury. Look at this admirable specimen: the internal malleolus is broken into the joint, and the joint has been full of blood, and inflamed. If those in charge of such a case did not take care, the joint would become very stiff. When in the habit of thinking in this way you will not leave a Pott's fracture long, but think in terms of morbid anatomy, and take good care to move the ankle-joint soon.

Here is another morbid anatomy specimen, a contraction of the palmar fascia. I once had the privilege of being a colleague and friend of the late William Adams—a man of unusual courage and sagacity—and saw him working at the morbid anatomy of this disease. I commend his methods to your earnest consideration. William Adams was the first to saw through one of the long bones. He sawed through the neck of the femur, and invented for the purpose the instrument which has long gone by his name, Adams's saw. You, gentlemen, can scarcely comprehend what that meant when every one expected the patient to die of blood poisoning. Dupuytren's name was given to this affection because he was a distinguished man, and one who founded his surgery on anatomy and morbid anatomy. Before Dupuytren investigated contraction of the palmar fascia it was considered to be due to shortening of the tendons. I learnt all this from William Adams, and watched him work in the most painstaking manner to confirm Dupuytren's observations and make them his own. First he got a skilled anatomist to dissect the palmar fascia and all its prolongations; then specimens of its contractions. After Adams had made his observations he devised, from his knowledge of anatomy and morbid anatomy, the operation for the subcutaneous division of the fascia. Nowadays we should not mind whether the fascia were divided subcutaneously or by an open wound; but in those days many wounds suppurated, so that surgeons had a great leaning towards subcutaneous operations. The suppuration was thought to be due to the entrance of air. They did not know much about bacteria or of the necessity for boiling instruments. Still, they knew that a small puncture was safer than a wide incision.

When I watched this very able man's methods, it dawned upon me that morbid anatomy was the clue to clinical surgery. Morbid histology, bacteriology, and the other "ologies" are but additions.

Pott's name is also associated with the tuberculous disease of the spine, admirably shown in this specimen. Here we can see the light which morbid anatomy throws upon clinical surgery. You all know how that spine became bent. First, we may suppose that the child had tuberculous milk to drink. As Dr. Walsham has so clearly pointed out, the tubercle bacilli entered through the tonsils and adenoid tissues in the naso-pharynx. Then they passed by the lymphatics or blood stream to various parts of the body—in this instance to the cancellous tissue of the vertebrae. First the cancellous tissue became inflamed. You will have had abundant opportunities of seeing tuberculous glands in the neck. In the beginning these are swollen, tender, painful, but not very hard; and so, doubtless, are the vertebrae. Next the inflammatory products caseate, so that the bodies of the vertebrae become composed of stuff of a cheesy appear-

ance and consistence. Plenty of caseous material is seen in this specimen. Cessation is the naked-eye evidence of the presence of tubercle bacilli. You may be occasionally deceived by other forms of degeneration. Gummata look very like caseous material, also some malignant growths. Then this caseous material softens and an abscess forms. There is the morbid anatomy; what are the clinical events? At first the vertebra is inflamed. That series of joints, the spine, behaves like any other inflamed joint and ceases to move. Next, the muscles in the vicinity of the inflamed joint waste, so do those of the spine. A caseous gland ceases to be painful and becomes very hard. But the next stage—that in which the caseous material softens—has a great bearing on clinical medicine, because now, instead of the bodies of the vertebra, there is an abscess. But weight cannot be supported upon an abscess, so the spine collapses and the Pott's curve appears. One who thinks in terms of morbid anatomy will now try to find out what may have become of the pus. This may never be known; but in the ordinary sequence of events the abscess will reach the surface of the body.

Here anatomy lends us its aid. From between the bodies of the vertebrae come out the nerve trunks. The pus follows their course, and some goes backwards along the posterior division of the spinal nerve, forming a lumbar abscess. Or the pus may track down the anterior divisions of the spinal nerves which enter the psoas muscle. Therefore look for a psoas abscess. The pus may pass along the anterior crural nerve into the iliac fossa. An iliac abscess is easily overlooked unless morbid anatomy be invoked. Finally, the pus may go down the lumbosacral cord, along the sciatic nerve, into the gluteal region, and hence a gluteal abscess.

Next, taking the morbid anatomy of Pott's fracture one step further. Here is a specimen showing how the damage is repaired. If tubercle was a very fatal disease I suppose the inhabitants of this island would have been wiped out long ago. But the human body has great power of resisting tubercle. When the body of a vertebra has nearly all disappeared, then the process of repair begins. The process of repair in these abscess cavities is always the same—they heal by granulation. When the granulations have grown in the gap in the vertebral column they become fibrous or bony. Thus you may have seen people walking about after Pott's disease of the spine and carrying sacks of coal upon their shoulders.

Here is another specimen showing the bearing of morbid anatomy upon clinical surgery. It is a tuberculous knee. This is what happened. Tubercle bacilli entered and were deposited in the cancellous tissue of the bone or in the synovial membrane, and the joint became inflamed, stiff, and flexed, and the muscles wasted. (Note the inevitable flexion. A leg with a bent knee or hip is of little use. The bending must be prevented with splints or weights, and if it has taken place be overcome.) Then the inflammatory material caseated. The caseous material is quiet. At last the caseous material softened. Then an abscess formed, either in the synovial membrane or in the cancellous tissue of the bones. When the caseous material softens the pus may point under the skin. When opened this abscess may lead into the joint, so that in opening it be careful lest a fresh infection be implanted upon the tuberculous. If the patient resists the tubercle, and has not had septic arthritis in addition, the disease follows exactly the same course as in the spine. William Adams often pondered over articular cartilage. Why should the cartilaginous femur ossify all but two little pieces at each end? Articular cartilage never ossifies, although in osteo-arthritis its edge may thicken and ossify. So that as long as articular cartilage is present movement may return, and as long as it is present ankylosis cannot occur. But in the course of tuberculous arthritis the cartilage is removed in the manner seen in this specimen. Granulations grow from the bone, and from the synovial membrane, and eat the articular cartilage. When it has done that, and filled the space between the ends of the bone, it becomes fibrous or bony. Fibrous union is painful. The fibrous ankylosis may be firm and strong, and without movement, but it will be painful. Your object is to attain bony ankylosis, which is strong, and does not hurt. The late Sir William Savory used in sonorous phrase to say, "Fibrous ankylosis is the opprobrium of surgery." But the removal of articular cartilage and the

formation of bone is a long and tedious process. So the surgeon steps in and performs an operation founded upon morbid anatomy. He cuts this process short, and gives the patient in a little time what Nature would give after many months of suffering. Accordingly he removes the whole tuberculous synovial membrane and tuberculous bone. But he does something more: he removes the whole of the articular cartilage, puts the ends of the bone together, and keeps them there until they unite by bone.

A stricture of the urethra is, to my mind, a very good example of the way of thinking from morbid anatomy to clinical surgery. Here is a stricture of the urethra. It is in the membranous urethra, and very narrow. It might have measured 7 mm. in circumference. Do not assume that the membranous urethra is the commonest place for strictures. The commonest place for strictures of the urethra, if you examine the urethra properly with bullet-headed bougies, is in the anterior part. The person with a stricture in the anterior part of the urethra has, as a rule, a reflex spasm of the compressor urethrae. Those who are not aware of this, and who do not use bullet-headed bougies, overlook the anterior stricture, and mistake the contracted compressor urethrae for one. A good deal of what is written about strictures is not founded, as it should be, upon correct morbid anatomy, but upon very incorrect clinical observation. Think what will be the first clinical effect of that 7 mm. stricture—obviously a small stream of urine. Make it your invariable habit, when a patient has anything the matter with the genito-urinary system, to see the urine passed into a glass. That affords the opportunity of seeing the kind of stream and the urine with those significant flakes of lymph in it. Other results may follow. Not only may the stream be altered in shape, but much time be spent in passing it through such a narrow channel. A gentleman told his medical man that it took him a long time, nearly a quarter of an hour, to pass water. It was thought that, because the patient took such a long time to urinate, he must have an abnormally large quantity to pass, and diabetes was diagnosed. As a matter of fact, he had a rather narrow stricture deep down in the urethra. The late Mr. Berkeley Hill operated, and, when told what had been said about diabetes, laughed, and replied, "They often say that." But I do not think it can be a very common mistake, nowadays at least, but it is a mistake which may befall any one unless the urine is passed into a glass. Observe the next thing which has happened in this case of stricture. The urethra behind the stricture is a little dilated. What does that mean? That dilated urethra collects a little urine in it after the patient has finished micturating. That remains, and is a source of discomfort, and additionally so because it runs out afterwards and soils the clothing. And so people with stricture not infrequently smell of stale urine. Notice the next thing which has happened. The bladder is thick and small. When the human muscles, either striped or unstriped, have to do additional work they become bigger and stronger. Think what happens to a man who has a stricture which has resulted in hypertrophy of the bladder. In the first place, the bladder will not hold much. The person with a bladder like the one I hold up will pass at the utmost 4 or 5 oz. of urine at a time. But still his kidneys, we hope, are making the proper quantity, so he has to pass urine frequently. If you have seen him in the act you will know that only 4 or 5 oz. issue at a time. What next? The hypertrophied bladder is enormously strong, and the person who has a hypertrophied bladder, as soon as he has a desire to pass water has to rush off and pass it as quickly as possible, otherwise the strong bladder will shoot the urine out into the garments. But the hypertrophied bladder does not merely expel the urine through the stricture; it drives it into the ureters, which become dilated together with the hilum of the kidney. Thus one with a stricture of the urethra is in a sad plight. Before dividing the stricture learn how the kidneys are excreting. Those with incompetent kidneys oftentimes die. Unfortunately, too, a septic infection may coexist. An unboiled catheter is a dangerous weapon.

I will merely take one more morbid anatomy specimen, which is important because it belongs to a very common and dangerous disease. There is not much to be seen in this inflamed appendix. If you make up your mind that a patient has an inflamed appendix, be sure that he has some septic material in its interior, with ulceration such as this

of the lining membrane, and the passage of the infection into the lymphatics, and on its way to the peritoneum. A dangerous state of affairs. A line cannot be drawn between anatomy and morbid anatomy. This inflamed appendix is situated at the ileo-caecal junction, a very critical part of the alimentary tract. Inflammation at the ileo-caecal junction disturbs the function of the whole alimentary tract. Proceed in anatomical order. Begin at the mouth to learn the clinical history. Consideration of the mouth will lead you to ask what can be eaten. Those with appendicitis commonly say that they cannot eat certain things; they will say they have had to give up fruit or vegetables. Every now and again they say they can eat anything. But most people with appendicitis have to be careful about their diet. From the mouth go on to the oesophagus, and that will remind you to ask whether there is vomiting. If a person with appendicitis vomits it means that the attack is acute. And after the oesophagus comes the stomach. Those with appendicitis often have indigestion and pain after food. Next we come to the intestines. An inflamed appendix causes blocking at the ileo-caecal valve, so that wind and faeces do not escape easily from the small intestine. Thus we find continual rumblings, because the small intestines are trying to get wind past the ileo-caecal valve. They may not be able to get it past, and so distension supervenes. Ladies with this condition may have to undo their stays. Passing onwards to the large intestine, do not fail to inquire about its functions. Those with appendicitis are often constipated. You cannot place implicit reliance on any of these signs, but proceeding anatomically you will find one or other of them is almost sure to be present. No person can have an inflamed structure in the interior of the body without general signs—alteration of temperature, of pulse, a leucocytosis, and rigors when the infection is especially dangerous. A person with appendicitis who has rigors is like one with whitlow, except that the whitlow is in the interior of the abdomen, not at the periphery of the body. Continuing your anatomical train of thought, the pelvic position of the appendix will be remembered. About 20 per cent. of those with appendicitis have the appendix on the brim of the pelvis or hanging into the pelvis, so that the inflammation can spread to the pelvic organs. First, there is the bladder: so inquire for pain and frequency of micturition. When the inflammation reaches the rectum or sigmoid there is pain during defaecation. In females there is a close connexion between the right broad ligament and the appendix, so that inflammation may spread from one to the other. Besides, the appendix may lie in contact with the right ovary and tube, so that not infrequently we meet with salpingitis or pyosalpinx. The ovary seldom becomes inflamed. Thus anatomy and morbid anatomy teach you to consider the pelvic organs in all cases of appendicitis.

In response to an invitation from the Committee and Medical Officer of the St. Marylebone Dispensary for the Prevention of Consumption, an influential and representative gathering visited the dispensary, at 15, Allsop Place, N.W., on January 25th. The principles and methods of work were explained to the visitors, who also inspected the institution. As is well known the chief sources of infection exist in the homes. To the dispensary, which is free, patients who are not already under a medical practitioner may go for examination, diagnosis, and treatment. From the dispensary a physician and nurse visit the homes of the patients, who are thus educated in open-air treatment, and in the protection of others by the observance of a few simple precautions. The medical officer also examines all those who have been in contact with the patients, and thus the early cases are diagnosed and treated. From the dispensary suitable cases are sent to sanatoriums for cure, advanced cases to special hospitals, and children are taught the value of fresh air and sunlight. The dispensary works in intimate association and co-operation with the public health authority, and with the various charitable agencies in the borough. The dispensary is the central bureau for the collection and dissemination of information regarding tuberculosis, and the centre of scientific and philanthropic effort against this disease. The St. Marylebone Dispensary is on the lines of those already established at Edinburgh, in the Borough of Paddington, and in Berlin, Montreal, New York, Boston, and Pittsburg, and is entirely supported by voluntary subscriptions.