Heart disease and Bright's disease are so intimately associated with arterial degeneration that it is often impossible to say in a given case how far the condition is cardiac or renal; the absence of response is to a great extent a measure of the renal element in the case and of value for prognosis, signifying, as a rule, that the condition is terminal.

REFERENCE.

¹ BRITISH MEDICAL JOURNAL, November 26th, 1910, p. 1670.

TWO CASES OF DEATH FROM POST-ANAESTHETIC ACID INTOXICATION.

BY

GILBERT BROWN, M.B., CH.B.,

LATE HOUSE-SURGEON AND SENIOR RESIDENT MEDICAL OFFICER,
CHILDREN'S INFIRMARY, LIVERPOOL.

It has long been known that acetone is present in the urine of patients who have been anaesthetized.

Becker, for instance, states¹ that acetone follows anaesthesia in at least two-thirds of the cases, irrespective of the nature of the anaesthetic or the duration of the anaesthesia, "and that if it be already present it is increased thereby." This has been confirmed by Abram² in a series of cases at the Royal Infirmary, Liverpool.

The importance of this fact has been recognized by all, though some, like Wallace and Gillespie of the Prince of Wales's Hospital, London,³ fully appreciate the dangers and take suitable prophylactic measures.

The two cases I have to record both occurred at the Children's Infirmary, Liverpool, within a few weeks of one another.

CASE I.

The patient was a boy aged 6 years, suffering from a chronic intussusception which had been present for six days. On August 10th, 1910, at 11.30 a.m. he was given ethyl chloride, followed by ether by the open method. A right rectus incision was then made, and a large chronic intussusception found and reduced easily; there were no adhesions and practically no congestion. The abdomen was sutured in three layers.

The anaesthesia lasted about twenty-five minutes, and 3½ oz. of ether were used.

After-progress.

At 8 p.m. the respirations were very rapid (40), though there were no râles in the chest. He was given atropine gr. $\frac{1}{20}$ four-hourly.

At 1.30 a.m. he was worse, expiration being prolonged at both bases.

At 2.30 a.m. he vomited, so was given sod. bicarb. ʒj in ʒij of water; this was retained, and no more vomiting took place.

At 10.0 a.m. he was much worse; pulse 160, but hardly perceptible. Cyanosed; dilated pupils; he was given oxygen and brandy. Retained salines well.

At 1.30 the same day the child was just conscious, but obviously dying; the breath smelled of acetone. Respirations 72; air hunger; pulse hardly perceptible. Cyanosis not present as oxygen was given continuously.

Urine.

A catheter specimen of urine was obtained and presented the following characteristics:

Light colour, slight deposit of mucus, acetone smell just perceptible, acid, sp. gr. 1028. No albumen, bile, pus, or sugar. Phosphates precipitated on boiling. Acetone present in large amount, demonstrated by both nitro-prusside and iodoform tests. Diacetic acid was present. No microscopic examination was made.

The child died before treatment could be commenced, and no post-mortem examination was allowed.

Both the night sister and nurse noticed that the breath had a peculiar odour, but did not report it as they "thought it due to a new kind of chloroform."

CASE II.

This was a child, aged 1 year and 10 months, with inguinal hernia.

On September 17th, 1910, at 5.30 p.m., anaesthesia was commenced with equal parts of chloroform and ether given on a Schimmelbusch mask, but the child stopped breathing twice, although he was not deeply under. Artificial respiration was necessary on both occasions. A change was made to ether by the open method, and this was borne very well.

The anaesthesia lasted about forty minutes; of C.E. mixture ʒij were used, but much was wasted; of ether ʒjss.

After-progress.

Had a comfortable night, and was well till 7 p.m. on the following evening, when vomiting occurred and the pulse became 180, temperature 103.2°; ? smell of acetone on breath; respirations 44. Chest normal, child very quiet. Pupils normal.

Urine.

Specimen of urine obtained was clear and of light colour; a large amount of white urates appeared on cooling; the urine

was acid and smelt of acetone. No albumen, bile, pus, blood of sugar.

Acetone was shown in very great quantity by Legal's nitro-prusside test and Lieben's iodoform test, and diacetic acid was present.

Treatment was at once commenced as follows: (1) Rectum washed out with sod. bicarb. solution; (2) ʒj sod. bicarb. in ʒij water per rectum; (3) sod. bicarb. ʒss in water every hour by mouth.

This had no effect and the child became worse.

1.45 a.m. Much worse, respirations very rapid, pupils dilated, continuous rectal saline commenced.

4.55 a.m. Pulse much weaker, given pituitary extract miiij and sod. bicarb. ʒij in one pint of water, subcutaneously.

7 a.m. Moribund, given oxygen, and pituitary repeated.

8.5 a.m. Death.

An autopsy at 2.30 the same afternoon showed the liver slightly paler than usual, but otherwise everything was normal.

REMARKS.

In neither case was the urine tested for acetone before the operation.

Since the occurrence of these cases the urine of every child in the infirmary is tested for acetone on admission by Legal's test, and if present the operation is delayed, if possible, till the acetone has been eliminated, and treatment is continued after the operation.

Wallace and Gillespie pointed out⁴ that an individual deprived of carbohydrates excretes acetone and diacetic acid, but the exhibition of carbohydrates rapidly causes the disappearance of these substances. They therefore employ glucose as a prophylactic. So, acting on their suggestions, the following scheme has been instituted at the Children's Infirmary, Liverpool:

All patients admitted to surgical wards for operation (usually admitted 2.30 p.m.) have

1. Glucose, ʒj, at 4, 6, and 8 o'clock on day of admission and 4, 6, and 8 o'clock on following morning. If the child is 8 years or older ʒij are given.

2. Sod. bicarb. in the strength of ʒj to ʒiv of water instead of plain "wash-out" before operation.

If acetone is present in urine on admission,

1. Continue glucose two-hourly in daytime.
2. Give sod. bicarb. ʒj in ʒj of water two-hourly in day and four-hourly in night.

Liquid glucose is more satisfactory than the solid, and children take it well as a sweet. Sod. bicarb. solution is frequently only taken by children after a struggle.

Since the initiation of this measure no serious symptoms have developed in any patient; but the treatment has not been in force long enough to give a reliable comparison, though the post-anaesthetic vomiting has been less.

I am indebted to Mr. Dun for permission to publish the first case, and to Dr. McClellan for the second.

REFERENCES.

¹ *Deut. med. Woch.*, Leipzig, No. 19, 1895, 17 and 18, 1894. ² *Liverpool Medico-Chirurgical Journal*. ³ *Practitioner*, February, 1910. ⁴ *Ibid.*

RADIUM IN CANCER.

BY

CHARLES J. MORTON, M.D. EDIN.

SURGICAL RADIOGRAPHER TO GUY'S HOSPITAL

It is now some years since the newer method of using radium, advocated by Wickham and Degrais,¹ of Paris, and others, was introduced, but as yet there are few recorded results of cases treated in this country. I have employed the method in suitable cases during the past two years, and a description of some of the more typical may be of interest.

The method differs from the older one of enclosing the radium in glass or metal tubes. Instead, pure radium in the form of sulphate is spread on a varnished surface of metal or linen, and screens of various densities are interposed between the applicator and the tissues. In this way it is claimed that a much smaller quantity of radium is required, and the amount of the radiation can be graduated and concentrated either on the surface or on the deeper parts. The best results appear to be obtained from applicators containing 2½ mg. of pure radium to each square centimetre. I have used two applicators of this strength, one measuring 4½ and the other 9 square centimetres.

In the majority of my cases the disease has been in a very advanced stage. In two only was the cancer still in an operable condition. These were very similar cases of