It would seem, therefore, that although definite failure in one or more of these tests is not enough, *per se*, on which to reject a candidate, still a combination of three or four failures is often accompanied by evidence of actual or recent impairment of one or other of the various systems of the body. It is noteworthy also that, in practically every case that could not be said to satisfy rigid standards, failure was due to inability to reach the minimum rather than to exceed the maximum limit, there evidently being a decided tendency towards diminished rather than increased labyrinthine reaction sensibility.

4. Relationship between these Tests and others Used by the Aviation Candidates Medical Board, Hampstead.

In order to see if there is any relationship between the results of these tests and those employed routinely by the Aviation Candidates Medical Board a series of 100 cases was taken and a comparison made. The tests selected for comparison were three in number—namely, tremor of eyelids or hands, self-balancing, and the rod and board test. It was found, however, that a pilot might have what are accepted as normal labyrinthine reactions and yet at the same time perform only moderately or badly one, two, or all three of the above-mentioned tests. Of the pilots discussed under (3), only one of the three with all four labyrinthine reactions unsatisfactory was poor in two tests—namely, self-balancing and the rod and board test; the other two performed all the tests satisfactorily. In addition, in the case of the other pilots discussed under (3), abnormal labyrinthine reactions seemed to bear no relationship at all to the successful execution or otherwise of these three tests. This is perhaps not surprising, as the turning reactions only test the integrity and tone of the cerebro-cerebellar-vestibular tracts, whereas it is well known that many factors may enter into the successful execution of the other tests cited.

5. Value of these Tests from the Point of View of the Medical Selection of Pilots.

At present, and until other and perhaps surer standards are laid down, one is inclined to view the utility of these rotation tests, at all events from the point of view of routine examination, as being problematical. It is true that they may draw attention in certain instances to particular defects which might or might not have been otherwise noted, but, on the other hand, their rigid observance might easily exclude men whose past histories and recent performances are those of perfectly successful pilots. Their value, however, and the data they give to anyone capable of interpreting such information is undoubtedly great in certain instances, as the following two cases recently seen by myself will show:

CASE I.

Lieutenant L. complained of deafness in the left ear incurred in May, 1918. The history was that whilst descending, and about 700 feet from the ground, he felt a tightness in the head and a mist before his eyes. He made a bad landing—"pancake"—was so dizzy on attempting to alight that he could not keep his feet, and had to be taken to his quarters on an ambulance and put to bed. He did not vomit, but things seemed to go round him to the right and he slept badly. During the next few days he was very shaky on his feet when attempting to walk. About two weeks later he was permitted to start flying again. He is now unable to stay long in the air and is always sick should he remain up as long as an hour. Before his accident he was a good stunter, but now either cannot or dare not stunt. On examination he was found to be totally deaf in the left ear, and his nystagmus and past-pointing indicated impairment of the left labyrinth; he was actually sick whilst being tested. The caloric test confirmed the view that his left labyrinth was not functioning, so that here was a pilot with a completely destroyed left labyrinth whose defect had unfortunately not been recognized, and who, being allowed to fly was a constant danger to himself and anyone else who happened to be his observer.

CASE II.

Lieutenant G. was, in May, 1917, in a balloon which was struck by lightning; the telephone receivers were in his ears at the time. The balloon was immediately hauled down, but he felt "overpowered by dizziness," which was accentuated when attempting to walk to his hut after reaching the ground. He also noticed that he was quite deaf in the left ear. After a week's leave he was again put on balloon duty and managed to continue for three months. During this time he always felt uncomfortable and dizzy in the air, though never troubled similarly before the accident. At the end of these three months he was, fortunately for himself, transferred to ground duties. On recent examination for assessment of gratuity,

I found him to be completely deaf in the left ear, and the rotation and caloric tests showed a functionless left labyrinth. Like Case I, this was an undiagnosed case of a destroyed labyrinth, which only careful rotation and caloric tests could demonstrate.

CONCLUSIONS.

1. Experienced pilots have, if anything, a slight tendency towards diminished labyrinthine reactions.
2. Disturbance, present or past, of some other system of the body may affect labyrinthine reactions.
3. Where deafness is present no medical examination of a candidate or pilot is complete without a careful investigation of the functions and reactions of the semicircular canals of both sides.
4. In the absence of a discovery of present or past signs or symptoms pointing to an aural affection, routine examination by means of the Bárány tests is superfluous, provided that a thorough general medical examination is made.

THE EFFECTS OF RADIUM TREATMENT ON WAR INJURIES IN THE NEIGHBOURHOOD OF NERVES.*

BY

WALTER C. STEVENSON, M.D.DUB.,

CAPTAIN R.A.M.C., HOME HOSPITALS RESERVE,
SURGEON TO DR. STEVENSON'S HOSPITAL AND TO THE INCORPORATED
ORTHOPAEDIC (CHILDREN'S) HOSPITAL OF IRELAND, DUBLIN.

THAT radium in suitable doses does not injure nerve tissue is shown by the work of Mme. Fabre and Dr. Paul Touchard on five cases of syringomyelia, all of which improved, showing increased mobility of the limbs with diminution of stiffness. One patient's hand had been quite helpless in a *main en griffe* position, but after radium treatment she was able to flex, extend, and separate her fingers. A second was able to resume his work as a draughtsman after six months' disuse of pencil and compasses. Another remarkable fact was the disappearance, in one case, of the trophic disturbances characteristic of Morvan's disease.¹

The conclusion that the powerful physical agent (the gamma rays from radium C) has some effect on injured nerve areas was forced on me, more or less by accident, in the following case:

CASE I.

Pte. S. fell off a transport wagon on March 15th, 1917, fractured the anatomical neck of his left humerus and injured his brachial plexus. When admitted to the Special Military Surgical Hospital, Blackrock, over eight months later, he had limited movement of the shoulder and little power to move his wrist; he was unable to flex his thumb and fingers, especially at the distal joints. Sensation in the hand and wrist, except over the ulnar distribution, was impaired. His first three fingers were discoloured, and there were trophic changes of the nails of the second and third fingers. On December 31st, 1917, graphs of the movements of the wrist and fingers were made with the aid of a strip of lead, as is the routine in this massage department. On January 18th, 1918, with the purpose of increasing the movement in the shoulder-joint, 100 millicuries of radium emanation screened by 2 mm. of lead in a flat applicator, kindly sent me by Dr. Lynham of the London Radium Institute, was applied for two hours in the axilla, and for two hours each on the front and back of the joint. Two days later the masseuse brought the patient to me and asked why he could flex and extend his wrist better and flex his fingers better, and why the discoloration of his fingers was less when radium was only applied to the shoulder. The graphs, which I have before me, taken on January 3rd, 1918, are conclusive evidence of an increase of voluntary movement, which cannot be explained by any change of weather or treatment, nor by suggestion, for it was not expected. The movement in the shoulder was practically unaltered except that voluntary abduction was temporarily reduced from 63 degrees to 59 degrees.

It is difficult to explain the effect of radium in the foregoing and following cases. I can only report the results of careful observations and accurate records in which I have eliminated as far as possible the many sources of error, especially suggestion. The sudden improvement in movements cannot be explained by greater personal attention to a patient, muscle re-education, the recent removal of a splint, or any change of environment or treatment that I am aware of. That radium will within six hours

* Paper read before the Section of Surgery, Royal Academy of Medicine in Ireland, November 21st, 1919.

alter the microscopic appearance of malignant cells has been proved by Morson;² that it will free or loosen scar tissue, and also that in some cases it will act like a charm in removing the pain and tenderness in a scar, I have demonstrated frequently. It has an anaesthetic as well as an analgesic effect.²

Radium cannot be expected to work miracles and materially benefit gross nerve lesions; it would appear, however, to stimulate normal functional activity in nerve tissues which have been injured but are on the road to recovery, either by softening the scar tissue around them, or by some action on the nerve cells themselves, possibly by improving their nutrition. In some cases radiation appears to lessen hyperactivity in a nerve which is being irritated by scar tissue.¹

When one realizes the number of nerve lesions that have occurred in this war, and the long periods during which one has to wait for recovery after nerve suture, any treatment that presents possibilities of shortening the period of incapacity is worth considering. On this account the following cases are, I believe, worth recording. All the patients except the first were treated by radium emanation supplied by the Royal Dublin Society, Radium Institute. I use a surface applicator of 1.5 mm. thickness of lead, as described in detail elsewhere.³

CASE II.

Pensioner A. fell 30 feet on April 28th, 1916, and fractured the lower part of his spine. On February 18th, 1917, his left leg was flaccid, the right leg spastic; he was incontinent, and could move about with difficulty on crutches. On December 28th there was no response to either current in the right extensor longus hallucis, or in dorsiflexors of foot or peronei on left. January 3rd, 1918: Radium applied to spine. January 7th: Patient stated that for the first time he felt that his feet were cold. A good response to galvanism was obtained in the dorsiflexors and evertors of the left foot and a slight response in the right extensor longus hallucis, ACC > KCC in each case. On January 10th, 1918, the patient could get about more easily on his crutches.

CASE III.

This case, Major B., should perhaps not be included here, as the patient's hand and wrist were radiated as well as the wounded area in the upper arm. He was wounded on February 10th, 1917; the musculo-spiral nerve was freed on May 17th, and the median on June 17th, 1917, by Mr. William Taylor at the Castle Hospital, Dublin, where he attended for massage. During the first six months of treatment he improved greatly, but for three months before I saw him the massense informed me there had been little or no improvement in the fingers and wrist. Radium was applied on February 19th, 1918, and I left Dublin a couple of days later. On March 10th the patient wrote: "I believe you have had the diagrams of my hand showing the enormous improvement it made in ten days. I think the results were extraordinarily good." The graphs verify his statements. I applied radium again on April 3rd, 1918, and shortly afterwards he was passed for homeservice.

CASE IV.

Pte. C. The median nerve, partially divided in the lower forearm, was sutured by Major W. S. Haughton on November 12th, 1917. On December 16th radium was applied for prevention of post-operative scar, and on the 17th there was some pain in the palm where it had been anaesthetic; voluntary extension of the wrist and extension and flexion of the fingers, which had been rigid from muscle spasm, now became possible. On January 9th, 1918, there was marked improvement in the movement of the wrist and fingers.

CASE V.

Pte. D. was wounded on the ulnar side of the forearm on December 11th, 1917. Limited voluntary flexion and extension of all joints of the hand. For months any attempt voluntarily to move the fingers had caused violent intention tremors. Radium applied to scar on August 9th, 1918. On the 12th the tremors had markedly diminished, and the patient had a much larger range of voluntary movement.

CASE VI.

Pte. E. was wounded through the left shoulder and chest on October 9th, 1917. He had pain in the scar at the back of the chest, recovering wrist-drop, and hyperextension of the fingers at the metacarpo-phalangeal joints. Radium given on September 21st, 1918, in the front of the shoulder, axilla, and over the scar. Three days later there was some improvement of flexion of the fingers; extension of wrist was much improved. Abduction of the shoulder had increased by 13 degrees, and the scar on the chest was less painful and tender.

CASE VII.

Pte. F., wounded in the left elbow on March 23rd, 1918; brachial artery severed and tied at both ends. Elbow could not be completely extended; hand red and congested, radial artery small. Radium was applied to front of elbow on August 6th, 1918. On the 24th he was seen with Major Haughton, who

agreed that the colour of the hand and the radial pulse were improved; he could extend his elbow completely, and the pronation of the forearm and power of grip were increased as shown by measurements.

CASE VIII.

Pte. G., wounded in right upper arm on July 18th, 1916; trophic changes first and second fingers. Median nerve freed from scar by Lieut.-Colonel W. I. de C. Wheeler on August 9th, 1917. On December 21st, 1917, radium was applied to the operation scar in the upper arm. Next day, opposition of thumb to little finger, which before treatment had been very slowly performed, was now briskly carried out; increase in range of movement of fingers and wrist, except flexion of first interphalangeal joint of first finger, which was limited. A week later further improvement was shown in graphs.

CASE IX.

Pte. H., wounded in left leg, March 16th, 1918; the divided external popliteal nerve was sutured by Major Haughton on August 12th, 1918. *Electric reactions*: Slight response in tibialis anticus to galvanic, none to faradic current; other muscles could not be tested on account of the current causing cramp in the foot. On August 21st radium was applied to the wound and operation scars. Next day the electrical reactions were greatly improved; response to galvanic current and slight flicker of movement to faradic current in the tibialis anticus; no response in other muscles, but no cramp as previously.

CASE X.

L.-Cpl. I., wounded middle left forearm, September 8th, 1916; two inches of median excised and nerve sutured by Major Haughton. On December 15th, 1917, graphs of the finger were taken immediately after massage. On December 16th radium was applied over the sutured nerve to limit the secondary scar of the operation. Next day there was some tenderness in the palm where it had previously been anaesthetic. Graphs taken before massage showed considerable improvement in flexion of fingers, following the application of radium; graphs made immediately after massage showed more than double this range of flexion.

CASE XI.

Pte. J. Wounded in the left shoulder on May 29th, 1917. Operation on median nerve near axilla February 4th, 1918. Hyperaesthesia of hand in region of median nerve distribution. Radium applied March 17th, 1918. Captain Alexander, in charge of the electro-therapeutic department, Alder Hey, reported three days later that there was no improvement in the hyperaesthesia, which had spread to the second finger and also to the radial area. Slight improvement in movement of shoulder.

CASE XII.

Pte. K. Wounded on September 26th, 1917, in the right shoulder, right arm, and immediately above the elbow. Radium was applied in these three localities on April 13th, 1918; the result was diminished tenderness of the scar near the elbow and some improved movement of the fingers. Again treated with radium June 27th, 1918; the movements of fingers before and after were accurately measured by a goniometer. The increase of voluntary and passive movement was as follows:

	Metacarpo-phalangeal Joint.		First Interphalangeal Joint.		Second Interphalangeal Joint.	
	Active.	Passive.	Active.	Passive.	Active.	Passive.
Index finger ...	25°	20°	30°	37°	6°	
Middle finger ...	9°	9°	30°	27°	9°	
Ring finger ...	19°	19°	17°	42°	5°	30°
Little finger ...	20°	13°	16°	20°	5°	

The angles of movement show that after radium treatment the patient could voluntarily flex his fully extended fingers as far as I could force them before radium, while the distance through which I could force them was further increased. For instance, at the index metacarpo-phalangeal joint before radium he could flex voluntarily to 52 degrees and I could force the finger to 72 degrees, while after radium he could voluntarily flex to 77 degrees and I could force it to an angle of 92 degrees. On July 1st, 1918, Captain Lloyd Roberts, neurologist, Alder Hey, examined the hand and noted the improvement in the voluntary movements and pliability of the fingers, though the electrical reactions were not changed since last examined by him on June 22nd; he suggested that the results were due to improved nutrition following radium treatment.

With the exception of a painful bulb in the ulnar nerve, where no improvement was apparent after radium treatment, the above cases include all of this type I can find in my notes of some 300 military patients treated at Black-rock, Alder Hey, and Shepherd's Bush Special Military Surgical Hospitals. Taken separately the cases may not be convincing, but taken collectively the weight of evidence

is that radium treatment serves a very useful purpose in certain types of nerve trauma and in certain stages in the repair of nerve injuries. The conclusions to be drawn from this work with radium in the treatment of war injuries affecting nerves can be summarized as follows:

1. Radium treatment cannot benefit gross nerve lesions; here operative treatment is indicated, to unite severed nerves or free them from dense scar tissue.
2. After a nerve operation, or after milder degrees of nerve trauma, it would appear to aid and to hasten the return of function in a limb.
3. It improves the nutrition in the area supplied by injured nerves.
4. It may be useful as an aid to diagnosis, and in certain cases will indicate or contraindicate the necessity of operation.
5. It is a valuable adjunct to other forms of treatment.

REFERENCES.

¹ *Radium*, by Dawson Turner, 2nd edition, p. 136. ² Radium Treatment of Scars, with Clinical Notes on Military Orthopaedic Cases, by Walter Stevenson, *Lancet*, March 23rd, 1918. ³ A Lecture on the Technique of the After-treatment of War Injuries by Radium, by Walter C. Stevenson, *Medical Press*, January 8th, 1919.

ON POISONING BY ARSENOBENZOL COMPOUNDS USED IN THE TREATMENT OF SYPHILIS.

BY

ALEXANDER G. R. FOULERTON, F.R.C.S.,

LECTURER ON HYGIENE TO THE LONDON SCHOOL OF MEDICINE FOR WOMEN, AND LECTURER IN THE HYGIENE DEPARTMENT OF UNIVERSITY COLLEGE, LONDON; COUNTY MEDICAL OFFICER OF HEALTH FOR EAST SUSSEX.

BEFORE considering the toxic action of arsenobenzol compounds it will be convenient, since the essential pathological processes appear to be closely similar in either case, to refer to a paper² published in 1918 which dealt with the pathology of another group of cases of poisoning in which what is termed toxic jaundice is a characteristic symptom, and in which there are found after death an infiltration of the liver and kidneys with fat, and in some cases an acute atrophy of the former gland. Six poisons, all of which have a special affinity for fats, were referred to as belonging to this group; three of them—chloroform, ether, and tetrachlorethane—are active fat solvents, and the other three—phosphorus, dinitrobenzene (D.N.B.), and trinitrotoluene (T.N.T.)—are readily soluble in fats.

The essential process of poisoning in these cases appears to be as follows: The poison, whether swallowed into the stomach or inhaled into the lungs, or fixed in the fatty constituents of the skin and absorbed slowly therefrom—as may be the case with one or other of the six—becomes fixed in the food fat of the circulating blood. This fat, with the poison fixed in it, is carried to the liver for the purposes of normal metabolism, there to be transformed by a process of desaturation of the fatty acids into the "available organic fat" which is believed to be the principal source of energy for the work of various active tissues. The destructive action of the poison thus concentrated in the liver is such that the functional activity of the gland is impaired, with interruption of the process of desaturation of the food fat and, presumably, disorder of function in other directions. The normal metabolism of food fat being checked, engorgement of the liver with fat and a condition of lipaemia, caused by an accumulation of unchanged food fat in the blood, result. In experimental poisoning of the cat and rabbit with trinitrotoluene it was found that considerable quantities of the surplus fat were excreted by the kidneys, being readily demonstrable in the urine. And so the kidneys become engorged with fat in process of excretion; and, as with the liver, there is disturbance of the functional activity of the gland as a result of the direct action of the concentrated fat-fixed poison. The investigation led to the conclusion, therefore, that the fatty changes which occur in the liver in such cases as those of delayed chloroform poisoning and poisoning by trinitrotoluene represent an infiltration with food fat which the gland is unable to transform, and that the similar condition in the kidney is caused by an infiltration with unchanged and useless fat which is in process of excretion from the body. The changes do not represent, as had been supposed hitherto, a fatty degeneration of the parenchyma cells of the gland. The first effect of the direct action of the concentrated poison on the liver, as seen in the earlier stages of experimental poisoning with trinitrotoluene, is a perivascular round-cell infiltration of the connective tissue along the course of the portal circulation. In the areas occupied by the migrating cells the first microscopic evidence of interrupted metabolism is found in the presence of

minute globules of fat which apparently have passed through from the portal capillaries. At a later stage, in human cases, the gradual accumulation of this fat produces the gross fatty infiltration which is the most obvious feature of the group of cases of poisoning. In the more prolonged poisoning of munition workers with trinitrotoluene the tracts of round-cell infiltration become replaced in places by dense fibrotic tissue, so that the parenchyma cells, already damaged by the direct action of the fat-fixed poison, undergo a further destructive degeneration which is secondary to the cirrhotic change.

The fatty changes in the liver and kidneys in fatal cases of poisoning by arsenobenzol compounds appear to be essentially identical with those just described. There is an infiltration of the liver with unchanged fat, resulting from an interference with fat metabolism caused by the direct action of the poison on the tissue of the gland. With this there is an apparently consequent engorgement of the kidney with surplus unchanged circulating food fat. In poisoning with the arsenical compounds, whilst there is not a ready explanation of the concentration of the poison in the liver which in the other cases is afforded by an obvious special affinity for fats, it appears to be possible that some similar fixation of the poison in the fat does in fact occur.

At a time when an investigation into the pathology of poisoning by T.N.T. amongst munition workers was nearly completed, three deaths amongst patients who had been under treatment with arsenobenzol compounds (novarsenobillon and kharsivan) came under observation towards the end of 1917, and were apparently the first cases of the kind recognized in this country. Many other cases followed, but for various reasons it was inexpedient to publish any details at the time; and in the paper quoted on the action of certain poisons which affect fat metabolism only a very brief allusion was made to the subject. The close resemblance between these cases and cases of delayed poisoning by trinitrotoluene was referred to; and it was stated that in an experiment in which a rabbit had been poisoned with arsenobenzol changes in the liver had been produced which were similar to those found after the experimental administration of trinitrotoluene. It had been found that in the rabbit the first appearance of fat was not in the parenchyma cells of the liver, but in areas of round cell infiltration which appeared along the course of the portal capillaries. Also there was evidence of the elimination of fat by the kidneys.

The following table (I) sets out the facts, so far as they were ascertainable, as to the dosage and time of appearance of toxic symptoms in 7 cases which occurred in the practice of six different hospitals, and in which it was possible to make a microscopic examination of the liver and kidneys after death. The table includes similar details which were obtained in an investigation of the circumstances of 9 cases treated in yet another hospital, in which the appearance of jaundice was followed rapidly by death in patients who were either under treatment with kharsivan or for whom the course had already been completed. Nothing is known as to the actual condition of the organs in these last 9 cases.

For purposes of comparison with these 16 cases two other series of cases recorded respectively by Professor Stuart McDonald⁴ and by Drs. Strathy, Smith, and Hannah⁵ may be referred to. The former series included 5 fatal cases which had been treated with "salvarsan," combined with the usual intramuscular injections of mercury. It is stated that in all of the cases the jaundice, which appeared at first without any serious general symptoms, was followed after an interval of from two to eight days by symptoms of severity, increasing jaundice being accompanied by haematemesis, active delirium, and coma, with death in from one to four days. Dr. Strathy and his colleagues refer to a series of 58 cases treated in a military hospital, in which unfavourable symptoms of one kind or another occurred either during or after a course of treatment with either arsenobillon or kharsivan or galy, combined in each case with intramuscular injections of mercury. Amongst the 58 cases there were 47 in which there was jaundice, and of these 8 died. The number of injections given in the fatal cases varied from four to eleven. The largest amount of the drug given in a fatal case was 6.95 grams and the least amount 2.2 grams. The average interval between the last injection and the onset of symptoms in a fatal case was forty-one days, the longest interval being forty-eight days and the shortest eighteen days. The average interval between the onset of symptoms and death was five days, the longest interval eleven days, and the shortest two days.

Regarded from the clinical point of view the 29 cases, including those recorded by Drs. McDonald and Strathy, which occurred in the practice of nine different hospitals, together with other published cases, appear to fall into