where the dilated tubes are not close to the surface (this is illustrated by experience of less debatable cases in childhood). If a careful history of such a case is taken, with the assistance of an elder relative, a clear record of bronchopneumonia, sometimes repeated, in early life is not infrequently obtained.

It has already been mentioned in dealing with bronchiectasis in childhood that blood is not infrequently coughed up, and that, occasionally, in notable amount. This may occur in adult life also, and I have known it to occur in a patient who had lost all recollection of former chest disease, but who presented a well marked basal bronchiectasis, of which his female relatives were able to supply a perfectly clear account from his early childhood. The case was, of course, under dire suspicion of pulmonary tubercle.

In addition to damage from childhood infections, the lungs of adults are also very apt to suffer scarring from the results of recovered influenzal pneumonia and of tuberculosis. Basal bronchiectasis or bronchiolectasis, not infrequently double, is not uncommonly to be seen since the recent epidemics. Sometimes it gets forgotten until some acute exacerbation brings it once more to the notice of the medical attendant. Occasionally it has been so far overlooked that some fresh disease, generally tuberculosis, comes under suspicion when new symptoms arise.

The lung damage caused by the healing of tuberculous lesions is so large and varied a subject that I shall not attempt to touch more than one aspect of it here. I would only urge that some amount of bronchiectasis is the usual accompaniment of chronic tuberculous lung disease. The point I wish to call attention to is the presence of crepitations over the diseased areas in cases of arrested or healed pulmonary tuberculosis. Such an occurrence is not uncommon, and yet one finds but little reference to it in the literature. I have many cases under observation which have been long arrested, but where crepitations remain, and might cause anxiety in the mind of an uninitiated observer. In some of these cases there are but the dry crackles of expanding lung, but in others the crepitations are somewhat moist and are due, I feel sure, to dilated bronchioles over a contracting fibroid area—generally in the region of a deep contracting cavity. The diagnosis of "arrest" may be difficult where these are present—it is greatly helped, in my experience, by the absence, in these cases, of the "reflex bands of percussion impairment" which I have described elsewhere on more than one occasion.

In conclusion, I should like to insist on the point, which perhaps I have hitherto only implied, that fibrosis and bronchiectasis, and "damaged lung" in general, vary according to the severity of the inflammation which produced the condition. It is attention to the slighter grades for which I here plead—the textbook account of bronchiectasis applies to the few marked and uncommon cases, whereas the slighter forms are not rare and are seen at all ages. They do not provide us with "pints of fetid sputum which separates into three layers," but they provide us with a varying symptomatology, and with problems in diagnosis, prognosis, and treatment, which may be of vital importance to our patients, and not without moment to our own *amour propre*, and, incidentally, our reputation.

REFERENCE.

¹ St. Bartholomew's Hospital Reports, 1905.

AN UNUSUAL CASE OF TYPHOID SPINE WITH SYMPTOMS OF SPINAL CORD AFFECTION.

BY

PROFESSOR H. TURNER,

ORTHOPAEDIC CLINIC OF THE MILITARY ACADEMY OF MEDICINE,
PETROGRAD.

NUMEROUS clinical observations and pathological research of past years have brought convincing evidence of the fact that the infective agent of so-called acute internal disease may settle on different parts of the skeleton, bringing injuries to bone, cartilage, or component parts of the joints (Virchow, Rokitsky, Ponick). It is well known that even such exclusively internal diseases as influenza may involve the osseous tissue. The "amphibious" qualities of the pneumococcus are proved by every day's observation. Its nefarious rôle in the production of osteomyelitis and arthritis has been very often the cause of diagnostic errors. The ranks of specific "surgical"

microbes, as primarily defined by Ogston, W. Cheyne, Rosenbach, and others, have been gradually enlarged by the enlistment of many other pathogenic elements, which have come forward from the domain of acute infectious diseases such as those belonging to the multifarious group of typhus—that is, typhoid, recurrent, and typhus (spotted) fever. The widespread epidemic of these latter diseases observed in Russia during recent years has given birth to vast material concerning this subject, and Russian literature, notwithstanding external difficulties, has been able to bring abundant illustration of the surgical complications of typhus fever in general. Typhoid or enteric fever, with its stubborn bacteria, capable of retaining vitality in the organism for many years after the conclusion of morbid symptoms, stood foremost in this relation, although recurrent and spotted fever did not remain much behind in showing the same quality. Clinical observations on a large scale gave us the opportunity of confirming the findings of Ebermeyer, Fränkel, Quincke, and many others, who discovered typhoid bacilli in the vertebrae, ribs, sternum, and other bones of the dead body.

The special tendency of the spine to become the seat of infection, as explained by Fränkel's demonstrations of living typhoid bacilli in the cancellous tissue of the vertebral body, we have seen represented in the numerous cases of typhoid spine which came under our observation in the orthopaedic hospital of the Medical Academy. Experience has taught us to mark off these forms of spondylitis amidst the greater number of tuberculous cases, and generally there was not much difficulty in avoiding errors of diagnosis.

The involvement of the spine in the typhoid process takes place, with great probability, during the flourishing period of the disease, and must be of not very rare occurrence. The symptoms of this complication may pass unremarked on the background of the grave picture of the illness. A new wave of temperature, accompanied by severe pains in the back, may draw attention to the intrusion of the virus into the spine. The further progress of the complication may be very slow, not taking a malignant course. It seems as if bone tissue, especially the cancellous part of it, is not a suitable pasture for the typhoid microbe, the feeble pyogenic and destructive character of which has been proved by many communications. The symptoms of the infection may in some cases subside without having been clearly understood, natural curative forces having extinguished its continuance. But as in other parts of the body, and especially in the intestinal channel, the virus of typhoid fever, secluded in the vertebrae, may retain its vitality for months and years, only to regain force under convenient circumstances, trauma in any form being the auxiliary agent.

It may be stated that the pathological anatomy of typhoid spine in the different stages of its development is not founded on such abundant data as in the case of Pott's disease. The benign course of typhoid spondylitis allows the patients to evade further clinical observation, and the final changes in the vertebrae have to be studied on occasional findings in the anatomical theatre or on the shelves of museums. The examination of these specimens, and the interpretation of the pathological changes presented by them, are hampered by the want of clinical notes. It is not easy, therefore, to define clearly the type of typhoid or other infectious spondylitis among the multifarious pathological processes to which the spine is so conspicuously liable. Roentgen rays have done very much to help our advance in this difficult field. The findings seen on the plate, elucidating the origin of clinical symptoms, present themselves as useful guides in the examination of dry pathological specimens. Taking this course of study, I have been able to discern with a great degree of positiveness the doubtful cases of infectious spondylitis compiled by me in the surgical museum attached to my clinic. The general impression received from the examination tends to confirm the supposition that the process of typhoid spondylitis is actually a form of subacute inflammation involving the periosteum, ligaments, and vertebral articulations. Gibney's term, "perispondylitis," remains the most suitable for a brief determination of the pathological changes. The posterior lateral parts of the laminae are principally involved, their mobility probably influencing the exacerbation of the process.

This point of view may help to explain the frequent localization of the illness in the lower dorsal and upper lumbar regions. Ossification of the tissues adjacent to the bone brings firm consolidation of the laminae and locking of the joints. The massive layers of newly formed bone on the

outer surface of the posterior wall of the spinal channel are sometimes exceedingly imposing. Continuous or interrupted bridges are to be seen between the vertebral bodies. The intervertebral discs seem to undergo a process of atrophy, the intervals between the adjacent surfaces of the bodies becoming narrower. In some cases there is to be seen complete synostosis of laminae, articulations, and bodies. Two or more vertebral bodies become firmly and inseparably united. One of my specimens coming from a juvenile subject clearly represents the result. The vertebrae have been united in three pairs, beginning from the lower lumbar and ending by the twelfth dorsal. The next three dorsal vertebrae (9-11) are also firmly glued together, showing traces of heavy destruction of the bodies. And yet there is no kyphotic curve of the spinal process line. The fact of the vertebral bodies having been impaired in this case emphasizes the common assumption that the anterior cancellous part of the column may not be exempt from the destruction. It seems to confirm the investigations of E. Fränkel, who was able to cultivate typhoid bacilli procured from bone marrow squeezed out of the spongy tissue of the vertebrae after death. The paradoxical absence of kyphosis in such cases is easily explained by the effect of previous ankylosing periostitis binding together the posterior halves of the laminae and locking the articulations. The same interpretation may be accepted for the rare occurrence of grave spinal cord symptoms in typhoid affections; paralysis from cord compression is not spoken of.

The picture of typhoid spondylitis as represented by the Roentgen plates is very characteristic. Attention is drawn to the narrowing or partial disappearance of the intervertebral spaces and to the lateral protrusion of the contiguous edges of the bodies. Dense vertical shades on one or both sides of the middle line reproduce the massive bridges of newly formed bone by which the posterior parts of the column are soldered together. A scoliotic curve may be occasionally seen as the result of asymmetrical disposition of the disease.

As regards the clinical symptoms of infectious spondylitis, it may be said that they differ very much from those in Pott's disease. Acute excruciating pain in the back is the chief complaint. Every least movement exaggerates the sufferings; the undulations of a spring mattress may bring them on. The exact situation of the process may be easily determined by manual examination, although kyphotic projection may be absent. Digital pressure reveals intense pain in the lateral parts of the spine. The soft coverings of the region involved may be thickened and oedematous. In some cases several foci of the disease are to be found in different levels of the spine. Fever of an intermittent type generally accompanies the process in its initial stage. Tendency to suppuration or the formation of fistulae is very rarely observed, the diagnosis thereby being deprived of bacteriological proof. According to our clinical experience and the bacteriological findings of Quinke, E. Fränkel, and others, one must allow the assumption that the involvement of the spine in the typhoid process is a frequent integrant part of the disease as a whole, the infection continuing to exist in a latent state. The awakening of the process is generally brought about by trauma, be it in the form of an early attempt to leave the bed or to resume physical work. Therein lies the explanation of the development of typhoid spondylitis some months or even years after recovery from the initial disease. The symptoms and clinical course of spondylitis produced by spotted or recurrent fever are similar to those above described.

The following case may be offered as an interesting addition to the general statistics of the subject:

L. M., aged 23, Red Army, infantry, admitted to the orthopaedic department May 21st, 1921. In November, 1919, he developed typhoid. In February, 1920, he began to suffer from pains in the back. Having re-entered service on August 12th after two months' leave he continued feeling the same pains. In December, 1920, he went through an attack of recurrent fever. Called up again soon after, he began to lose power in his legs in March, 1921. After a certain term of treatment in a military hospital he came under my care.

On admission his state was as follows: General health good. Robust complexion. Standing or walking absolutely impossible. Distinct signs of partial spinal cord compression in the form of ankle and patella clonus; Babinski's phenomenon; rigidity of muscles and weakness of the lower limbs; loss of muscular feeling in the toes; skin hyperaesthesia to pain on the level of the seventh to eighth dorsal segments, and hyperaesthesia to all kinds of sensation lower down. State of atactic paresis of the lower limbs. No trouble of the bladder or rectum. Examination of the spine shows a not clearly defined swelling between the shoulder blades corresponding to the fifth to seventh spinous processes. The integu-

ments of the same region oedematous and reddened. Fluctuation distinct. Palpation of the diseased area of the spine inflicts great pain, especially in the lateral parts of the vertebrae. Vidal's reaction of the blood positive. Temperature between 38° and 39° C.

Patient complains of great pain in the back, especially with every movement. On opening the abscess under general anaesthesia, on May 25th, slightly odorous grey-green pus was found, from which a pure culture of typhoid bacilli was obtained in the bacteriological laboratory of the Academy. There was no doubt about the identification of the micro-organism, as it showed all the usual tests. The cavity of the abscess was found to be located to the right of the spinous processes, leading deeply towards the corresponding laminae. No denuded bone could be felt during the scooping out of its contents.

The subsequent course of the case was favourable, although of long duration, spinal cord symptoms subsiding very slowly. The fistula, periodically closing and reopening, continued to produce pus containing the specific bacteria. Voluntary movements of the legs reappeared in July. The hyperaesthesia disappeared. Babinski's sign and clonus still persisted. By December, 1921, the patient was able to walk freely. At times he felt acute pains in the region of the operation. A probe periodically introduced into the fistula met with something hard resembling denuded bone. The attempts to extract the "sequestrum" were futile, although it seemed to be movable. On June 2nd chloroform was administered and the fistula cut open. With a strong forceps I extracted a foreign body, which proved to be the middle part of a penknife blade about 1½ cm. square, thus astonishing my staff and myself. The patient being eager, still in a drowsy condition, to see the obnoxious bone which had troubled him for so long, the piece of metal was shown to him. His look seemed to wander as if trying to recollect some distant event. "Yes," said he, after a pause, "there has been an affair." The details of the adventure were obtained on the next morning. In the winter of 1918, before he entered service, he took part in a sleigh trip in company of his equals containing representatives of his parish. Having the advantage of being able to play the harmonica, he was stabbed from behind by a jealous village comrade. The blade of the penknife broke into pieces on striking the vertebral laminae; its point probably penetrated the cortical layer of the bone and broke off. The small wound healed without any further trouble in a few days, and no subsequent attention was paid to the tragic event.

In need of a justification of my publishing the above case, I may point to the fact that it is the only positive one, according to my knowledge, in which typhoid spondylitis was complicated by suppuration. The bacteriological verification of the origin of the disease is also of singular interest. The previous trauma, which may have stimulated the onset of the suppuration, is also to be underlined. Symptoms of cord affection are a not very frequent concomitant in typhoid spondylitis.

The wound having healed in two weeks, the patient walked out of the clinic on June 29th, 1922, with some traces of nervous disorder in his legs.

TREATMENT BY DIATHERMY.*

BY

W. J. TURRELL, M.A., M.D. OXON.,

ELECTROTHERAPEUTIC PHYSICIAN, RADCLIFFE INFIRMARY, OXFORD.

THE title of this communication does not necessarily imply the adoption of any electrical method. It simply refers to treatment by penetrating heat. The subject consequently admits of division into two parts: (1) the therapeutic effects of heat penetration; (2) the most effective method of securing this penetration, having due regard to its safety, the ease of its regulation, and its capability of application to the body as a whole or any organ or part of the body.

As to the first part there can, I think, be no difference of opinion in regard to the great importance of heat as a therapeutic agent. Its value has, in fact, been admitted from the earliest times to which medical history extends, and probably no method of therapy has been more unquestionably or more universally adopted.

A variety of methods have been utilized for applying heat treatment—hot sand, hot bricks, poultices, radiant-heat baths, sun baths, paraffin baths, and many others too numerous to mention; but the therapeutic value of all these methods is distinctly and definitely limited, for the simple reason that their action is restricted to the surface of the body and the superficial tissues, and cannot be brought to bear directly upon the deeper organs and structures. They are consequently useless as agents for treatment by diathermy or heat penetration.

There is only one known method by which heat can be

* A paper read at the annual meeting of the Oxford and Reading Branch of the British Medical Association, July 15th, 1922.