The children given live vaccine alone, however, gave significantly higher antibody responses than when live vaccine was preceded by a single dose of inactivated vaccine.

This preliminary trial has not given any information on protection against the disease; this is the aim of a second study now in progress.

We are grateful to the following general practitioners for finding time in their busy practices to take part in the study:

Dr. D. M. Baker (Northwood), Dr. A. P. Barter (Bridport), Dr. S. Bedford (Grimsby), Dr. D. Budge (Leighton Buzzard), Dr. S. Carne (London W.12), Dr. G. Edelsten (Sutton Scotney), Dr. E. Gancz (Dartford), Dr. M. Goodman (Liverpool 6), Dr. R. P. C. Handfield-Jones (Haddenham), Dr. D. W. Hendry (Fife), Dr. E. R. Herst (Woodford Green), Dr. M. E. Holzer (London W.1), Dr. P. Hopkins (London N.W.3), Dr. S. G. Jeffs (Manchester), Dr. R. D. Jones (Godalming), Dr. H. Kidd (Buxton), Dr. K. Lane (London W.C.1), Dr. A. B. Lowther (Porthcawl), Dr. J. K. H. McCullough (Kings Norton), Dr. G. MacNaughton (Edinburgh 4), Dr. R. L. Meyrick (London S.E.6), Dr. J. H. Owen (Porthcawl), Dr. A. Wilson Robertson (Leven, Fifehire), Dr. Watson Rogers (Sherborne), Dr. H. S. Samuel (Theale), Dr. M. Schapira (Keighley), Dr. F. S. Shepherd (Southall), Dr. S. J. Silvey (Coombe Dingle, Bristol 9), Dr. E. Sklar (London N.W.10), Dr. G. R. Staley (Hull), Dr. J. M. Stuart (Coleshill), Dr. M. T. Sweetnam (Stoke-on-Trent), Dr. B. A. Thompson (Chester), Dr. M. K. Thompson (Addiscombe), Dr. T. J. C. Warriner (Chessington), Dr. G. I. Watson (Guildford), Dr. J. McA. Williams (Hartington, nr. Buxton), Dr. W. O. Williams (Swansea).

We are also grateful to Dr. M. T. Paterson, Dr. S. King, and Dr. W. H. S. Wallace, of London County Council Divisions 2, 4, and 8, and to the matrons and nursing staff of the day nurseries in these divisions for their help.

Our thanks are also due to Dr. J. A. Scott (since deceased), who at the time of the trial was Medical Officer of Health for the London

County Council, and to Dr. Ian Taylor, of the London County Council Department of Public Health.

Finally we would like to acknowledge the assistance of Dr. P. Benson, who obtained blood samples from the children in the day nurseries.

We are pleased to add our thanks to Glaxo Laboratories, Pfizer Ltd., and Wellcome Research Laboratories for the vaccines which they supplied free of charge.

REFERENCES

- Andelman, S. L., Schwarz, A., Andelman, M. B., and Zackler, J. (1963). *J. Amer. med. Ass.*, **184**, 721.
 Benson, P. F., Butler, N. R., Goffe, A. P., Knight, G. J., Laurence, G. D., Miller, C. L., and Pollock, T. M. (1964). *Brit. med. J.*, **2**, 851.
 Carter, C. H., Conway, T. J., Cornfeld, D., Iezonni, D. G., Kempe, C. H., Moscovici, C., Rauh, L. W., Vignec, A. J., and Warren, J. (1962). *J. Amer. med. Ass.*, **179**, 848.
 Enders, J. F., Katz, S. L., Milovanović, M. V., and Holloway, Ann (1960). *New Engl. J. Med.*, **263**, 153.
 — and Peebles, T. C. (1954). *Proc. Soc. exp. Biol. (N.Y.)*, **86**, 277.
 Fulginiti, V. A., Leland, O. S., and Kempe, C. H. (1963). *Amer. J. Dis. Child.*, **105**, 5.
 Guinee, V. F., et al. (1963). *Amer. J. publ. Hlth*, **53**, 645.
 Hendrickse, R. G., Montefiore, D., Sherman, P. M., and van der Wall, H. M. (1964). *Brit. med. J.*, **1**, 470.
 Katz, S. L., and Enders, J. F. (1959). *Amer. J. Dis. Child.*, **98**, 605.
 — Milovanović, M. V., and Enders, J. F. (1958). *Proc. Soc. exp. Biol. (N.Y.)*, **97**, 23.
 Krugman, S., Giles, J. P., Jacobs, A. M., and Friedman, H. (1962). *Amer. J. Dis. Child.*, **103**, 353.
 — — — (1963). *J. Pediat.*, **31**, 919.
 — Friedman, H., and Stone, S. (1965). *J. Pediat.* In press.
 Miller, D. L. (1964). *Brit. med. J.*, **2**, 75.
 Milovanović, M. V., Enders, J. F., and Mitus, A. (1957). *Proc. Soc. exp. Biol. (N.Y.)*, **95**, 120.
 Moncrieff, A., and Hussey, B. J. (1948). *Brit. med. J.*, **2**, 972.
 Rosen, L. (1960). *Amer. J. Hyg.*, **71**, 120.
 — (1961). *Virology*, **13**, 139.
 Schwarz, A. J. F. (1962). *Amer. J. Dis. Child.*, **103**, 386.
 Stokes, J., Hilleman, M. R., Weibel, R. E., Buynak, E. B., Halenda, R., and Goldner, H. (1961). *New Engl. J. Med.*, **265**, 507.
 Warren, J., and Gallian, M. J. (1962). *Amer. J. Dis. Child.*, **103**, 418.
 W.H.O. Expert Committee on Biological Standardization (1964). *Wld Hlth Org. tech. rep. Ser.*, No. 293.

A Laser Ophthalmoscope for Retinal Phototherapy

H. VERNON INGRAM,* O.B.E., T.D., M.B., F.R.C.S., D.O.M.S.; NEIL MANSON,* M.B., B.S., D.O.;
 DESMOND SMART,† M.Sc.

[WITH SPECIAL PLATE]

Brit. med. J., 1965, **1**, 823-827

The rapid advancement of maser and laser technology has provided medical science, and in particular ophthalmology, with a therapeutic coherent light source.

Maser and laser stand for microwave and light amplification by the stimulated emission of radiation. The maser works in the centimetre or submillimetre microwave region, and the laser in the near optical (almost to ultra-violet) frequency range. Since the media of the eye transmit the radiations in the range from 4,000 to 9,000 Å (Fig. 1) the energy from a low-power laser working in this range can be focused on the retina without producing any molecular disturbance of the transparent media of the eye.

Stimulated emission from an atomic assembly was first suggested by Einstein in 1917 from considerations of thermodynamic equilibrium. In 1953 and subsequently, the theoretical possibilities of microwave amplification were further considered by Weber (1953), Townes, Basov, and Prokhorov. Gordon,

Zeiger, and Townes (1954) produced an ammonia gas maser. The investigation of solid-state masers by Combrisson, Honig, and Townes (1956) resulted in the production of some discontinuously operating devices. Bloembergen (1956) postulated the continuous emission maser, and Scovil, Feher, and Seidel

FIG. 1.—Spectral transmission curve of the refractive media of the human eye.

* Department of Ophthalmology, University of Newcastle upon Tyne and Royal Victoria Infirmary, Newcastle upon Tyne.

† Head of Applied Physics Section, International Research and Development Company Limited, Fossway, Newcastle upon Tyne.

(1957) constructed such a device. The first light-amplifier appeared as the result of Maiman's (1960a, 1960b) work, and in the same year a coherent optical oscillator was produced by Collins, Nelson, Schawlow, Bond, Garrett, and Kaiser (1960).

A laser is therefore a device which can amplify a light wave, and, like all other amplifiers, the application of a suitable positive feedback makes it function as an oscillator. There are many types of laser—gaseous, liquid, and solid state. The laser with which we are concerned is the solid-state ruby laser. This is a sapphire crystal (Al_2O_3) in which 0.05% of the aluminium ions are substitutionally replaced by chromium ions. It is these chromium ions which give the red colour and which are responsible for the laser action in a ruby. They are excited to a higher energy level by the absorption of the blue-green light from a xenon flash tube, but they are very unstable at this level. Consequently they relax rapidly to a somewhat lower, more stable state, releasing a small amount of energy in the process and producing heat in the crystal. From this intermediate state the chromium ions eventually relax to their original ground state emitting a quantum or photon of red light, so producing the normal fluorescence. On the other hand, this photon may stimulate another excited chromium ion to emit a photon with the same phase, direction, and plane of polarization as the first. In this way a light wavefront can be built up, and this is the process of light-amplification by the stimulated emission of radiation.

To prevent the emission being multidirectional the shape of the crystal is made cylindrical, with flat parallel ends on which reflecting films have been placed. Any wavefronts which are not travelling exactly perpendicular to the end mirrors are soon lost out of the crystal, but those which are are amplified by being reflected back and forth through the crystal. Thus a very intense wavefront can be built up between the end reflecting surfaces, and by making one of them only partially reflecting some of the light can emerge from the crystal. The light emitted from a ruby laser is an intense parallel beam of red light of 6,943 Å at room temperature.

Material

In our experiments we used a ruby rod 2 in. (5 cm.) long and $\frac{1}{4}$ in. (0.6 cm.) in diameter mounted at one line focus of an elliptical cylinder and a xenon-filled flash tube mounted at the other line focus. This arrangement ensures good optical coupling between the ruby and the flash tube, as one property of an ellipse is that a ray of light leaving one focus must, after reflection at the ellipse, arrive at the other focus. The energy for the xenon flash tube is obtained from a bank of capacitors in a power supply unit. To produce a discharge from the flash tube the xenon must be ionized enough to allow the bank of capacitors, which are charged to 1,700 volts, to discharge through it. This is brought about by a pulsed trigger voltage of 7,000 volts applied to a wire wrapped round the tube (Fig. II).

FIG. II.—The ruby crystal and xenon flash tube in its elliptical cavity.

By this means a train of very intense pulses of red light are produced from the ruby, and these can be focused on the surface of, say, a metal to produce a very high electromagnetic field. The effect can vary from minor structural modifications to explosive volatilization depending on the energy in the beam.

Our experiments to develop a laser ophthalmoscope were initially undertaken with a standard Nelas ruby laser measuring 8 in. (20 cm.) long and 4 in. (10 cm.) in diameter, and weighing 13 lb. (5.9 kg.). To this a rather crude ophthalmoscope head was attached. This instrument was far too cumbersome to use clinically and was modified several times until the present model was designed. Our requisites were: (1) the instrument must be as easily handled as, and comparable in size to, an ordinary ophthalmoscope; (2) the illumination of the fundus must be adequate; (3) a localizing device must be projected on to the exact spot on the fundus on which the laser beam will be focused; (4) the laser beam must be in focus on the retina when the localizing device is in focus; (5) the diameter of the laser beam entering the eye should be as small as possible to avoid cut-off by the iris, particularly if the pupil cannot be maximally dilated; and (6) there must be no possibility of the laser beam or any reflection from it entering the eye of the operator or his assistant. This is achieved by the arrangement of the mirrors directing the laser beam into the patient's eye.

Fig. III shows the present prototype laser ophthalmoscope and power pack. The ophthalmoscope is of similar dimensions and weight to a standard battery-operated instrument. The ophthalmoscope head incorporates accessory lenses to correct both the patient's and the operator's refractive error. Variable reflex-free illumination is provided, and the laser is fired by

FIG. III.—The present model of the laser ophthalmoscope and power pack.

means of a microswitch on the handle in conjunction with a foot-switch. The threshold of the ruby crystal has been successively reduced from 290 joules in the first model to 120 joules in the latest. This gives a correspondingly higher maximum output without any increase in the power pack. Air cooling of the crystal has been incorporated so that the instrument can be fired repeatedly every three seconds. The power pack, which measures about 20 by 12 by 14 in. (51 by 30 by 35 cm.) and weighs 80 lb. (35 kg.), is set upon a small trolley and is easily moved about.

We have avoided using dichroic mirrors in the optical system because up to 8% of the beam may be transmitted instead of being reflected and may cause damage. For similar reasons we avoided shutters because they may fail mechanically and must inevitably interfere momentarily with the view of the fundus.

H. V. INGRAM *ET AL.*: A LASER OPHTHALMOSCOPE FOR RETINAL PHOTOTHERAPY

FIG. 1.—Lesion in rabbit's retina produced by 0.11 joule photographed immediately after exposure.

FIG. 2. — Rabbit lesion produced by 0.39 joule immediately after treatment. Note the small bubble in the centre of each lesion.

FIG. 3.—Human lesion. Arruga encirclage suture *in situ* with laser applications (0.2 joule) applied within the line of the suture.

FIG. 4.—Human lesion. Cystic detachment in upper temporal quadrant with line of laser applications along its lower edge. This was treated in two sessions and the pigmented lesions in the top right are four days old. The remainder were done the day of the photograph.

FIG. 5.—Human lesion. An area of peripheral retinal degeneration is walled off by laser applications. These are five days old and are well pigmented.

FIG. 6.—Ruby laser lesion produced by 10 joules, photographed a few seconds after application. Note the bubble of gas and debris and the haemorrhage.

H. V. INGRAM *ET AL.*: A LASER OPHTHALMOSCOPE FOR RETINAL PHOTOTHERAPY

FIG. 7.—Section of a 3-joule lesion, showing extensive disruption of retinal tissue with hole-formation.

FIG. 9.—Oscilloscope trace of a Q-spoiled laser application—a 12-megawatt pulse in 100 nanoseconds (one nanosecond = 1/1,000 of a millionth of a second).

FIG. 8.—Oscilloscope trace of a laser application lasting 800 microseconds (with a mean power of 1 kilowatt).

FIG. 10.—Section of a 0.19-joule lesion after 1 week. The diameter of the lesion was 0.5 mm.

G. W. BEVERIDGE AND A. A. H. LAWSON: BULLOUS LESIONS IN ACUTE BARBITURATE INTOXICATION

FIG. 1.—Section of an early bulla. (Haemalum and eosin. $\times 150$.)

FIG. 2.—Section from margin of an intact bulla. (Haemalum and eosin. $\times 200$.)

Methods

In our animal experiments we have exposed the rabbit's retina to various energies from 0.09 to 0.49 joule. The best results were produced by energies between 0.19 and 0.39 joule. The value at which a lesion could be detected clinically was 0.008 joule. A beam of 0.11 joule produced discrete areas about 0.1 mm. in diameter (see Special Plate, Fig. 1). Higher energies produce larger areas of coagulation about 0.5 mm. in diameter (Special Plate, Fig. 2).

Immediately after exposure a white area appeared at the spot treated. It is sometimes surrounded by a pinkish flush and the changes become more marked up to about three minutes after exposure. With the higher energies a pearly bubble appears in the centre but disappears rapidly—within five minutes. It is probably steam and carbon dioxide due to the localized disruption of organic tissue. Only once in these experiments, with the highest energy, have we seen a haemorrhage. This was quite small but was pre-retinal, and was reported by one of us elsewhere (Ingram, 1964). In none of our experiments with the ophthalmoscope have we found any damage to the transparent media of the eye either clinically or histologically. We have measured the temperature in the retina and found no significant rise after treatment. A 36 S.W.G. chromel alumel thermocouple was inserted under the retina through an opening in the sclera of the eye of a rabbit. The thermocouple had a normal spot-welded junction and was insulated up to the junction. The output of the thermocouple was monitored, using a Cambridge portable workshop potentiometer. The overall response time of the measuring system was between $1\frac{1}{2}$ and 2 seconds. Although this system does not permit measurement of very rapid changes in temperature, the temperatures recorded give a very good indication that the heating with a laser is consider-

FIG. IV.—Retinal temperature changes produced by 0.11 joule laser applications.

FIG. V.—Retinal temperature changes produced by varying laser energy output.

ably less than with the xenon arc coagulator, which uses up to 20 times as much energy. The laser beam, using various energies, was focused on the retina at various distances from the thermocouple junction. The results obtained are shown in Figs. IV and V.

The exposure time to the laser beam is very short—only 0.8 of a thousandth of a second—so that if the localizing device is in the right place on the fundus when the beam is released the patient cannot move his eye in time to do any damage to, say, the macula. It is quite painless, so that a retrobulbar injection is unnecessary and the patient can co-operate by looking in the appropriate direction.

The pupil should be as widely dilated as possible to avoid its cutting off part of the beam and thereby reducing the effect on the retina. We use 2% Cyclogyl (cyclopentolate) and 10% Neophryn (phenylephrine hydrochloride) to dilate the pupil, and to lessen still further this liability we have reduced the diameter of the laser beam to 2 mm. by inserting a telescopic lens system in the path of the beam.

Application to the Human Eye

The laser beam of the latest model of the ophthalmoscope has been applied successfully to the human eye. The areas of coagulation are smaller than those produced by either the xenon arc coagulator or by surface diathermy and vary from 0.1 to 0.5 mm. in diameter, but we think this is an advantage rather than a disadvantage, and will seal a hole or a tear in the retina with less disturbance to the function. It certainly means more individual applications, but the overall operating-time is not appreciably increased, and in many cases is reduced because the instrument is so much easier to use.

We have now come to the conclusion after treating 18 human cases that the effect of the laser beam in rabbits and man is different. In man a higher energy is needed to give a comparable effect. The clinical appearance of the lesion is a much better guide to the dosage required than the measure of the recorded output from the machine. Our present opinion is that the minimum energy which produces a small pearly bubble in the centre of the lesion is the ideal. This pearly bubble disappears in a second or two. We have also found that a higher energy or dose is required the farther out to the periphery we are treating.

The early appearance of the lesion also shows some differences from those in the rabbit. As soon as the central pearly bubble disappears, the circular lesion looks pale with a faint dark ring round it. This, we think, is due to the disruption of the pigment epithelium and an almost immediate pushing aside of the pigment granules. The lesion becomes pigmented within four days and is well marked in seven days.

Fig. 3 (Special Plate) shows a line of 0.2 joule laser applications to a human retina, placed inside the line of reaction caused by an encircling suture. In this case phototherapy was applied to limit an area of retinal detachment which was recurring across the suture line.

Fig. 4 (Special Plate) shows the fundus of a patient who had a cystic detachment of the upper temporal quadrant of the retina of the right eye. The laser applications can be seen all along the lower edge of the detachment, having been so placed as to limit further extension of the cyst while it is being treated by nascent hydrogen from multiple catholysis punctures.

Fig. 5 (Special Plate) illustrates the fundus of a female patient who has an almost total detachment in the right eye with vision of hand movements only. Operation elsewhere on two occasions failed to replace the retina. As she had degenerative changes in the periphery of the left retina, and it was considered that these were a potential cause of detachment, prophylactic treatment was undertaken. She was treated with the laser ophthalmoscope and adequate scarring has been produced to seal off the affected area.

Hazards in Industry

To investigate the hazards to workers with lasers in industry we have experimented with various types giving output energies varying from 1 to 10 joules, some being Q-spoiled.

Application of these higher dosages causes an immediate explosive disruption of the tissues in a rabbit eye unlike anything seen with clinical doses. The appearance is of a punched-out hole in the retina, the destroyed tissue being blown into the vitreous by the gas formed in the layers of the retina (probably steam and carbon dioxide). The gas is at first trapped in the debris, and as a result the vitreous opacity floats upwards (Special Plate, Fig. 6). The bubbles absorb and the debris sinks slowly downwards. If the explosion occurs in the vicinity of a vessel it gives way and produces a large vitreous haemorrhage. If not, choroidal haemorrhages are nearly always present. The majority of the vitreous haemorrhages we have produced under these circumstances have absorbed quite readily.

Sub-explosive yet nevertheless excessive dosage to the retina results in a central hole where the retina has been disrupted, and the base of the hole appears white owing to the ischaemic choroid and underlying sclera. The central hole is fringed by the damaged edge of the retina and choroid. These damaged tissues round the hole may swell during the first few days, and this tends to accentuate the depth of the central depression. By the third and fourth days pigmentary changes appear in the walls of the crater, and at the end of one week any haemorrhages produced are beginning to absorb (Special Plate, Fig. 7).

If the wavelength of these industrial lasers is such that the laser beam is largely transmitted, the clear media of the eye have mostly remained unchanged in short-term experiments. However, low energy but very high power outputs obtained by Q-spoiling cause a disruptive track through the vitreous with very extensive tissue damage and massive immediate haemorrhage. Normally a solid-state laser produces a train of about 800 or 900 relatively low-power output pulses (~ 50 kW). Q-spoiling is produced by removing the reflector from the lasing position at one end of the laser and replacing it when energy has been built up in the laser. A single giant pulse about 1,000 times more intense is then produced (~ 50 MW) (Special Plate, Figs. 8 and 9).

An increasing number of people are being employed in using such devices and could consequently be exposed to this new industrial hazard. Long-term experiments are necessary and are being performed to determine the effects of both pulsed and continuous lasers on the transparent media of the eye and the retina.

Histology of Laser Chorioretinal Lesions

With 6,943 Å energy outputs of from 0.11 to 0.29 joule good chorioretinal adhesion is produced in rabbits. These lesions are less than 0.5 mm. in diameter, being mostly of 0.1 mm. in diameter. Chorioretinal lesions of this order allow a uniform adhesion with the smallest possible area of tissue damage and functional loss. Higher doses are required to produce comparable lesions in man.

The two most important stages of a laser lesion histologically appear to be immediately after application and from six days onwards.

An Immediate Laser Lesion

The area of application macroscopically presents a pimple-like swelling. There is separation of the retinal layers and swelling of the tissue fibres. Cellular disorganization without cellular destruction occurs throughout the retinal layers and the innermost part of the choroid. Neither the adjacent vitreous nor the remaining choroid or sclera shows any change.

Laser Lesions at the Stage of Good Chorioretinal Adhesion (Special Plate, Fig. 10)

Chorioretinal adhesions can be recognized histologically in some cases as early as four days and in most cases are fairly well developed by seven days. Such a lesion shows pitting of the retina over the site of the adhesion. This pitting is enhanced by any intraretinal separation produced by handling and sectioning, but the firmness of an adhesion under these conditions cannot be doubted. Macroscopically the orientation and architecture of the retina appear surprisingly undisturbed. The *sclera* shows no histological change. The terminal capillaries in the *choroid* beneath the adhesion show dilatation but no vessel proliferation or increase in intravascular cell content. The tissue fibres show some separation, but there is no increase in cellularity. The *retinal* lesion does not involve the subjacent vitreous "hyaloid" condensation. The nerve-fibre, ganglion-cell, and inner plexiform layers remain correctly orientated but there is infiltration from the molecular cell layers. The molecular layers show pyknotic cells indicative of cell destruction. The pigment epithelium shows clumping and disruption, with some migration of pigment into the other retinal layers. The loss of cellularity, degree of disorganization, and increase in pigmentation vary from lesion to lesion and with the degree of pigmentation which pre-existed.

A comparison of the histological changes produced by the laser and xenon arc photocoagulator shows some differences. The xenon arc lesions show involvement of the *sclera* and internal hyaloid membrane, choroidal intravascular clumping, and eventually a more extensive fibrous reaction.

Surface diathermy compared with the laser shows even more marked differences, with greater involvement of *sclera* and vitreous, and the chorioretinal adhesions take up to 10 days to develop.

The tissue separation in the early lesions and the development of a firm adhesion by seven days suggests that patients even after prophylactic laser treatment should be allowed only restricted exercise for the first week. Failure to do this may result in retinal detachment, as has already been noted after prophylactic xenon arc photocoagulation.

The histological difference between xenon arc photocoagulation and laser lesions may be accounted for by the laser lesion being the result of a molecular disruption in a very high electromagnetic field strictly limited to the focused area of the beam on the retina. With high laser energies the molecular disruption can be so gross as to be explosive.

The premise that the changes are due to a molecular disruption in the area of an induced electromagnetic field is supported by the wedge shape of the lower-energy lesions. Similar shaped changes occur in metallurgical experiments with lasers. With higher laser energies the molecular disruption can be so gross as to be explosive, and the histology of the lesions produced is very different from those of the lower-power clinical applications. In the case of lesions produced by energies of 3 to 10 joules from ruby and neodymium lasers, there is a central area of complete retinal destruction down to, and in some cases including, the choroid. Where there is no hole formation a marked fibrous reaction occurs in the outer part of the lesion, and contraction of this produces corrugation of the vitreous surface of the plaque. There is disruption of the pigment epithelium and migration of pigment, most pronounced throughout the retinal layers in the periphery of the lesion. In most cases, but not all of the sections examined, adhesion of the retina to the underlying choroid and sclera occurs. The diameter of these lesions produced by the industrial type of laser which we have sectioned so far have measured up to 1.4 mm. in diameter.

Conclusion and Summary

A ruby laser may be used to produce good chorioretinal adhesion of limited dimensions and with minimum structural

damage to the tissues. The application of this form of therapy by an ophthalmoscope produces new technical problems, but when these are adequately solved it could lead to a wider use of phototherapy.

The history of laser development is reviewed very briefly. The laser ophthalmoscope designed in Newcastle upon Tyne is described. The present model is very similar, both in size and in weight, to a standard battery-operated ophthalmoscope. The ruby crystal is contained in the handle, which is connected to the power pack by a flexible lead.

Preliminary clinical experiments which were carried out on rabbits are described and the results are compared with those obtained in the limited number of human beings so far treated with the instrument.

The differences between laser lesions, diathermy lesions, and lesions produced by the Zeiss xenon arc coagulator are discussed.

The use of various lasers in industry are referred to briefly, and some effects of high doses on rabbits' eyes are shown to illustrate possible hazards which may be encountered as the use of lasers becomes more widespread.

The histology of therapeutic laser lesions is described and a brief reference is made to some lesions produced by excessive dosage.

We wish to acknowledge the help we have received from Professor A. G. Heppleston and his staff, of the department of pathology.

We also wish to record our gratitude to Dr. H. Rose, of the International Research and Development Company, Limited, for his enthusiastic collaboration, and to Mr. W. Hetherington for all his help with the practical work of constructing the laser. We are also indebted to Thermal Syndicate Limited, Wallsend on Tyne, for supplying the laser crystals, and to Messrs. Keeler Limited, of London, for their collaboration in producing the ophthalmoscope.

The cost of the research was borne by the International Research and Development Company, Limited, and the United Newcastle upon Tyne Hospitals.

BIBLIOGRAPHY

- Alsop, L. E., Giordmaine, J. A., Townes, C. H., and Wang, T. C. (1957). *Phys. Rev.*, **107**, 1450.
 Basov, N. G., and Prokhorov, A. M. (1954). *J. exp. theor. Phys. (U.S.S.R.)*, **27**, 431.
 Bloembergen, N. (1956). *Phys. Rev.*, **104**, 324.
 Collins, R. J., Nelson, D. F., Schawlow, A. L., Bond, W., Garrett, C. G. B., and Kaiser, W. (1960). *Phys. Rev. Lett.*, **5**, 303.
 Combrisson, J., Honig, A., and Townes, C. H. (1956). *C.R. Acad. Sci. (Paris)*, **242**, 2451; *Onde élect.*, **36**, 989.
 Einstein, A. (1917). *Zur quantentheorie der Strahlung*.
 Gordon, J. P., Zeiger, H. Z., and Townes, C. H. (1954). *Phys. Rev.*, **95**, 282.
 ——— (1955). *Ibid.*, **99**, 1264.
 Ingram, H. V. (1964). *Trans. ophthal. Soc. U.K.* In press.
 Maiman, T. H. (1960a). *Nature (Lond.)*, **187**, 493.
 ——— (1960b). *Brit. Commun. Electron.*, **7**, 674.
 Meyer-Schwickerath, G. (1960). *Light Coagulation*, p. 49. Kimpton, London.
 Scovil, H. E. D., Feher, G., and Seidel, H. (1957). *Phys. Rev.*, **105**, 762.
 Weber, J. (1953). *Trans. I.R.E., Prof. Group on Electron Devices*. PGED-3 (June, 1953).

Small-intestinal and Gastric Abnormalities in Ulcerative Colitis

S. N. SALEM,*† M.B., CH.B., M.R.C.P.ED.; S. C. TRUELOVE,* M.D., F.R.C.P.

Brit. med. J., 1965, **1**, 827-831

In a previous article (Salem et al., 1964) we reported the results of a pilot study of the state of the small intestine and stomach in ulcerative colitis. It was found that during an actual attack of ulcerative colitis the small-intestinal mucosa frequently showed villous abnormalities in biopsy specimens and the faecal fat excretion was often abnormally high; and, correspondingly, the stomach often showed superficial gastritis. Among the patients studied in remission the stomach and small intestine were usually normal as judged by these same criteria.

Since then the study has been considerably enlarged. First, a bigger group of patients with ulcerative colitis has been studied, so that some of our tentative conclusions of the pilot study can be re-examined with the aid of more extensive data. Secondly, some of the patients have been reinvestigated to see if any abnormalities in their small intestine and stomach fluctuate in unison with changes in the severity of their ulcerative colitis. Thirdly, because some of the villous abnormalities of the small intestine were comparatively minor, various control subjects have been examined to provide a yardstick of normality—a necessary step in view of the limited biopsy observations which have been made upon the healthy human small intestine.

Methods

Small-intestinal Biopsy.—We have continued to use a Crosby capsule, but one modified by the fitting of a latex sheath to avoid the occasional loss of the dome of the capsule and by the use of radio-opaque tubing to facilitate radiological localization.

* Nuffield Department of Clinical Medicine, University of Oxford.
 † Physician in the Kuwait Government Medical Service on study leave.

These modifications are described in detail elsewhere (Salem et al., 1965). The biopsy specimen was fixed and stained as in the pilot study.

Gastric Biopsy.—This also was performed by means of the Crosby capsule.

Faecal Fat.—As in the pilot study, the faecal fat excretion has been based on a four-day collection of stools with the patient on a standard hospital diet (or on his ordinary diet if an out-patient), using the method of van der Kamer et al. (1949) to estimate total lipids. A daily faecal excretion of 6 g. or more of total fat has been taken as abnormal.

Vitamin-B₁₂ Absorption.—No further observations have been made since the pilot study, when it was found that most patients with ulcerative colitis showed normal absorption.

D-Xylose Excretion Test.—In this pilot study it was found that the standard test employing 25 g. of D-xylose gave rise to abdominal pain and increased diarrhoea in patients with ulcerative colitis, so that the test was unsuitable in that form. The modified test of Santini et al. (1961), which employs 5 g. of D-xylose, has proved to be suitable for use in ulcerative colitis and has been utilized. Santini et al. (1961) have provided evidence of the reliability of this test with the smaller dose, a finding confirmed by others (Sammons et al., 1962).

Electron Microscopy.—Some of the small-intestinal biopsy specimens have been examined under the electron microscope by Dr. J. C. F. Poole, of the Sir William Dunn School of Pathology, Oxford University. For this purpose the biopsy specimen was immediately cut into two halves on recovery from the capsule. One half was immediately impregnated with osmium tetroxide and divided into small portions for fixation