- Foggie, A. (1962). *Symp. zool. Soc. Lond.*, 6, 51.
 Fulton, F., and Dumbell, K. R. (1949). *J. gen. Microbiol.*, 3, 97.
 Habel, K. (1962). *J. exp. Med.*, 115, 181.
 — (1963). *Ann. Rev. Microbiol.*, 17, 167.
 Harnden, D. G. (1964). *Amer. J. hum. Genet.*, 16, 204.
 Heath, R. B., Fairley, G. H., and Malpas, J. S. (1964). *Brit. J. Haemat.*, 10, 365.
 Huebner, R. J., Rowe, W. P., Turner, H. C., and Lane, W. T. (1963). *Proc. nat. Acad. Sci. (Wash.)*, 50, 379.
 Koprowski, H., Love, R., and Koprowska, I. (1957). *Tex. Rep. Biol. Med.*, 15, 559.
 Lindenmann, J. (1963). *Proc. Soc. exp. Biol. (N.Y.)*, 113, 85.
 — and Klein, P. A. (1965). *J. Immunol.*, 94, 461.
 Massimo, L., Vianello, M. G., Dagna-Bricarelli, F., and Tortorolo, G. (1965). *Brit. med. J.*, 2, 172.
 Moore, A. E. (1949). *Cancer (Philad.)*, 2, 525.
 — (1951). *Ibid.*, 4, 375.
 Moorhead, P. S., Nowell, P. C., Mellman, W. J., Battips, D. M., and Hungerford, D. A. (1960). *Exp. Cell Res.*, 20, 613.
 Nichols, W. W., Leven, A., Hall, B., and Ostergren, G. (1962). *Hereditas (Lund.)*, 48, 367.
 O'Reilly, K. J., Smith, C. E. G., McMahon, D. A., Wilson, A. L., and Robertson, J. M. (1965). *J. Hyg. (Lond.)*, 63, 213.
 Perner, L., Fowler, G. A., and Nauts, H. C. (1958). *Acta med. scand.*, 162, Suppl. No. 338, p. 1.
 Porter, G. H. (1963). *Arch. intern. Med.*, 3, 572.
 Price, W. H., Lee, R. W., Gunkel, W. F., and O'Leary, W. (1961). *Amer. J. trop. Med. Hyg.*, 10, 403.
 Sever, J. L. (1962). *J. Immunol.*, 88, 320.
 Shah, K. V., Cole, G. A., Russ, S. B., Needy, C. L., and Buescher, E. L. (1962). *Symposium C&V. Biology of Viruses of the Tick-borne Encephalitis Complex, Praha*, p. 303.
 Shein, H. M., and Enders, J. F. (1962). *Proc. Soc. exp. Biol. (N.Y.)*, 109, 495.
 Smith, C. E. G. (1956). *Nature (Lond.)*, 178, 581.
 — Turner, L. H., and Armitage, P. (1962). *Bull. Wld Hlth Org.*, 27, 717.
 Smorodintsev, A. A. (1963). *Proc. 7th int. Congr. trop. Med. Malar., Rio de Janeiro*.
 Southam, C. M. (1963). *J. Pediat.*, 63, 138.
 — and Moore, A. E. (1951). *Amer. J. trop. Med.*, 31, 724.
 — (1952). *Cancer (Philad.)*, 5, 1025.
 Thompson, W. R. (1947). *Bact. Rev.*, 11, 115.
 Webb, H. E., and Burston, J. B. (1966). *Trans. roy. Soc. Med.* (in press).
 — and Chatterjee, J. B. (1962). *Brit. J. Haemat.*, 8, 401.
 Wheelock, E. F., and Dingle, J. H. (1964). *New Engl. J. Med.*, 271, 645.
 Wiedemann, H.-R. (1964). *Lancet*, 1, 721.
 Yabe, Y., Samper, L., Bryan, E., Taylor, G., and Trentin, J. J. (1964). *Science*, 143, 46.
 Zilber, L. A. (1962). *Acta Un. int. Cancr.*, 18, 127.

Observations on Infective Drug Resistance in Britain

H. WILLIAMS SMITH,* D.SC., PH.D., F.R.C.V.S., DIP.BACT., M.C.PATH.; SHEILA HALLS,* B.SC.

Brit. med. J., 1966, 1, 266–269

Infective drug resistance is now known to be important in the spread of drug resistance among members of the Enterobacteriaceae. It has been studied extensively in Japan, where it was first discovered in 1959; the results have been reviewed by Watanabe (1963). By comparison little has been published in other countries. Observations on the situation in Britain have been made by Datta (1962), Anderson and Datta (1965), and Anderson and Lewis (1965). These have dealt mainly with *Salmonella typhimurium*.

It was felt that an impression of the importance of infective resistance as a cause of drug resistance in the Enterobacteriaceae in Britain might be obtained by assessing its incidence in *Escherichia coli*, since these organisms form the bulk of the enterobacteriaceal flora of the alimentary tract of man and domestic animals. The results of such a survey are reported in this paper. So are studies involving the transfer, *in vitro*, of resistance from non-pathogenic *E. coli* isolated from the faeces of man, calves, pigs, and fowls to some of the principal *Salmonella* and *E. coli* serotypes that cause disease in these species. The experiments included the use, as donors, of *E. coli* that were tetracycline-resistant only, organisms that are often found in the faeces of pigs and fowls fed diets containing tetracyclines (Smith and Crabb, 1957). The results of transferring resistance among *Salmonella* serotypes and from them to pathogenic serotypes of *E. coli* are also described.

Methods

Transfer of Resistance

Nutrient broth (Oxoid No. 2) in 10-ml. amounts was seeded with 0.02 ml. of a 24-hour broth culture of the "donor" strain and 0.1 ml. of a similar culture of the "recipient" strain. The mixed culture was incubated at 37° C. for 24 hours and centrifuged. The deposit was inoculated heavily on to a Petri plate of solid medium containing a drug to which the donor was resistant and the recipient was initially sensitive (referred to as selection medium). The plate was then incubated at 37° C. for 24 hours and examined for colonies of the recipient strain. If any were found a few were

picked, purified by plating on solid medium, and a single colony was then picked into broth and the drug resistance pattern of the resulting culture determined. Controls consisting of broth that had been inoculated with the recipient culture only were always kept.

When the recipient was a *Salmonella* sp., deoxycholate citrate agar was used as the basal selection medium, the inoculum sometimes first being incubated in selenite F medium at 37° C. for 18 hours in the hope of suppressing the *E. coli* if they were of the kind that multiplied on deoxycholate citrate agar. When the recipient was *E. coli*, MacConkey agar was employed as the basal selection medium, except when the donor was also a strain of *E. coli*. In the latter event nutrient agar containing washed erythrocytes was used if either donor or recipient was haemolytic. When both were non-haemolytic differentiation was usually an impossible task. When the donor was streptomycin-sensitive the situation was overcome by preparing a chromosome mutant from the recipient that was streptomycin-resistant and incorporating streptomycin as an additional drug in the selection medium. On a few occasions a strain of *E. coli* naturally resistant to furazolidone only was used in a similar manner, the additional drug added to the basal selection medium in this case being furazolidone. Normally the drug employed in the selection medium was streptomycin (40–80 µg./ml.), oxytetracycline (20–40 µg./ml.), chloramphenicol (20 µg./ml.), ampicillin (20–30 µg./ml.), neomycin (20 µg./ml.), or furazolidone (10–20 µg./ml.).

Sensitivity Tests

Approximately 0.15 ml. of a 1 in 50 dilution of a 24-hour broth culture of the strain to be tested was spread evenly over the surface of a dried Petri plate of MacConkey agar, and disks containing different drugs were then applied at approximately equal distances apart. The amounts of the drugs used were 50 µg. of streptomycin (S), oxytetracycline (T), chlortetracycline (T), chloramphenicol (C), and neomycin (N), and 25 µg. of ampicillin (A) and furazolidone (F), and 150 µg. of sulphadimidine (Su). The plates were incubated at 37° C. for 24 hours and read. Except in the case of sulphadimidine no difficulty was experienced in making the readings; the zones of inhibition surrounding the disks were either wide or very narrow or non-existent. All cultures that appeared to be sulphadimidine-resistant were retested on nutrient agar in which sulphonamide inhibitors had been neutralized by the addition of lysed horse

* Animal Health Trust, Stock, Essex.

Apart from those included in Table II another 10 strains of *E. coli* that were resistant to the tetracyclines only—five from pigs and five from fowls—were examined to determine whether their resistance was of the infective type. This was done because these *E. coli* strains are the kind that constitute the bulk of the *E. coli* flora of the alimentary tract of pigs and fowls that are fed commercially on diets containing low levels of tetracyclines in the hope of increasing their growth rate (Smith and Crabb, 1957). With the *S. dublin* strain and the streptomycin-resistant mutant of the enteropathogenic pig *E. coli* 0141:K85ab→85ac as recipients the resistance of four of the five pig strains was shown to be infective, tetracycline resistance being transferred from three of the strains to *S. dublin*, and from three to the enteropathogenic *E. coli*. By a similar procedure, but substituting a streptomycin-resistant mutant of a fowl pathogenic *E. coli*, 02:K1:H6, for the enteropathogenic pig strain, the resistance of all of the five fowl strains was shown to be infective, tetracycline resistance being transferred from four of the strains to *S. dublin* and from three to the pathogenic fowl strain.

The details of donor and recipient cultures employed in attempts to transfer drug resistance to the principal *Salmonella*

and *E. coli* serotypes that cause disease in man, calves, pigs, and fowls from non-pathogenic *E. coli* isolated from the faeces of these species are given in Table III. Apart from strains H1, B258, P14, and C9 not being used in mixed culture experiments with the *E. coli* pathogenic to man, calf, and fowl, attempts were made to transfer resistance from all the strains listed as donors to all those listed as recipients. The reason for the omission of these *E. coli* from mixed culture experiments was the impossibility of devising a selection medium to distinguish donor from recipient. Owing to their haemolytic nature this was overcome in the case of the *E. coli* pathogenic to pigs by using blood agar as the basis of the selection medium; all the donors were non-haemolytic. It was overcome in the experiment employing C12 and P17 as donors by using as recipients chromosome mutants of the pathogenic *E. coli* that were streptomycin-resistant, and including streptomycin as well as oxytetracycline in the selection medium; the mutants were prepared in the laboratory. C12 and P17, being tetracycline-resistant only, were included in these experiments because they were representative of the *E. coli* found in pigs and fowls fed diets containing tetracyclines.

TABLE III.—Details of Experiments Using Non-pathogenic *E. coli* as Donors and *Salmonella* and Pathogenic *E. coli* Serotypes as Recipients

Donor Non-pathogenic <i>E. coli</i> Strains			Recipient Strains	
Identification	Origin	Resistance Pattern	Identification	Principal Host Pathogen
H1	Faeces, man	S Su T C	<i>S. typhi</i>	Man
B258	" calf	A S Su T C	<i>S. paratyphi B</i>	"
P14	" pig	S Su T C	<i>S. typhimurium</i>	All species
C9	" fowl	S Su T	<i>S. dublin</i>	Calf
P17	" pig	T	<i>S. cholerae-suis</i>	Pig
C12	" fowl	T	<i>S. gallinarum</i>	Fowl
			<i>S. pullorum</i>	"
			<i>E. coli</i> 0128: K67: H2	Man
			<i>E. coli</i> 055: K59: H7	"
			<i>E. coli</i> 0141: K85ab; K88ab	Pig
			<i>E. coli</i> 0141: K85ab→K85ac	"
			<i>E. coli</i> 02: K1: H6	Fowl
			<i>E. coli</i> 011: K?	"
			<i>E. coli</i> 078: K80 (2 strains)	Calf

Apart from the inability to transfer resistance from P17 to *S. typhi* and *S. pullorum* all the transfer attempts were successful. Several attempts were often necessary before success was achieved, for the same reasons that applied in the studies on the incidence of infective resistance in *E. coli* (see above).

With some exceptions the complete resistance pattern was always found to have been transferred intact. The exceptions were: (1) When P14 was donor the resistance pattern found in the recipient *S. typhi*, *S. paratyphi B*, *S. typhimurium*, *S. cholerae-suis*, and *S. gallinarum* cultures was always S Su C; T resistance was never detected in these recipients irrespective of the drug used in the selection medium. (2) Only the resistance pattern S Su T C was transferred from B258 when either S T or C was used in the selection medium. When A was used instead the recipients acquired the resistance pattern A, A S, or A S Su.

In other studies using multiple-resistant strains of *E. coli* as donors it was not unusual to isolate recipients that possessed different combinations of the complete resistance pattern of the donor. These different combinations were usually obtained when different drugs were used in the selection medium. Sometimes a variety were obtained when the same drug was used. One extreme example of this was noted in mixed-culture experiments employing an *E. coli* strain to which the resistance pattern A S Su T C N had been transferred in the laboratory as donor and 12 *Salmonella* strains as recipients. Of 24 colonies picked from the 12 plates of selection medium containing ampicillin only two had the complete resistance pattern A S Su T C N. The other combinations noted were A Su N C (5), A N S (1), A S Su C (1), A N (5), A S (4), and A (6).

Two strains of *E. coli* whose resistance was of the infective type were of special interest because they had been isolated as long ago as March 1956, freeze-dried soon afterwards, and reconstituted for testing in 1965. They had been isolated from two epidemiologically unrelated outbreaks of diarrhoea in calves very shortly after chloramphenicol became available for use in calves. It was possible to transfer their complete resistance pattern S Su T C to *S. dublin* and *S. cholerae-suis*, the only two cultures with which they were tested.

Infective resistance was also demonstrated in three pathogenic strains of *E. coli*, the only ones tested. These were a human pathogen, 0111: K58: H2 (resistance pattern S Su T), and two pig pathogens, 0147: K89: K88ac (resistance pattern S Su T) and 0141: K85ab→K85ac (resistance pattern T).

Transfer of Resistance from *S. typhimurium* to other *Salmonella* Serotypes and to Pathogenic Serotypes of *E. coli*

All attempts to transfer resistance from three unselected resistant strains of *S. typhimurium* to 16 pathogenic strains of *E. coli* were successful. Only 24 different combinations of donor and recipient were studied.

The *E. coli* strains belonged to the following serotypes: 0128: K67: H2, 055: K59: H7, and 0119: K69 (man); 0141: K85ab; K88ab, 0141: K85ab→K85ac, 0147: K89: K88ac, 0138: K81, and 08: K87: K88ab (pig); 02: K1: H6, 078: K80, and 011: K? (fowl); and 078: K80 (calf).

In the experiments with one of the *S. typhimurium* strains the complete resistance pattern A Su T was transferred intact. In those with the other two all the resistance factors were shown to be transferable, but no recipient was isolated that possessed the complete resistance pattern of the donor. The recipients from one of these donors, whose resistance pattern was A S Su T N F, were either A S Su, A S Su N, S Su N, A N, A, T, N, or F resistant. Those from the other, whose resistance pattern was S Su T F, were either S Su, S Su T, S Su F, or F resistant.

A small number of attempts were made to transfer resistance from two of the resistant *S. typhimurium* strains to strains of other *Salmonella* serotypes. Recipient colonies on the selection medium could only be identified by slide-agglutination tests. Even so, it was possible to detect resistance transference in eight experiments, indicating a high transfer frequency. From the strain with the resistance pattern S Su T F resistance was transferred to a strain of *S. dublin* (A) and *S. typhimurium* (A T). From the strain with the resistance pattern A S Su T N F resistance was transferred to a strain of *S. cholerae-suis* (A S), *S. dublin* (A and A S Su), and *S. gallinarum* (A S).

Stability of Infective Resistance

Twenty-two infectively resistant strains were passaged daily for 10 days in 20-ml. amounts of nutrient broth at 37° C., the necessary subcultures being made with the tip of a platinum wire. Sensitivity tests were performed after each passage. The results are given in Table IV. Loss of resistance by some of the organisms in the culture, as evidenced by the presence around a drug-containing disk of a clearly defined zone of bacterial growth which was less plentiful than the rest of the growth on the plate of medium used for sensitivity-testing, occurred during passage of four of the strains made infectively resistant in the laboratory and four of the strains that had acquired infective resistance naturally. This loss of resistance was first noted at the third to the seventh passage, the proportion of organisms sensitive to one or more of the drugs to which they were initially resistant increasing progressively with passage until at the final passage the change was almost complete. The numbers of different kinds of resistant organisms

In culture No. 9 (Table IV) at the tenth passage were estimated by means of solid media containing different drugs. The total viable count and that of ampicillin-resistant organisms was $1,000 \times 10^6$ per ml. The counts of neomycin-resistant, chloramphenicol-resistant, and streptomycin-resistant organisms were only 50×10^6 , 40×10^6 , and 3×10^6 per ml. respectively. Tests on a small number of colonies from the drug-containing plates of media used for counting revealed that they were either A, A C, A C S, A S, or A N resistant.

TABLE IV.—Effect of Passage in Nutrient Broth on Stability of Infective Resistance

Culture No.	Identification	Infective Resistance Pattern	Origin of Infective Resistance	Resistance Lost during Passage
1	<i>S. cholerae-suis</i>	A S Su T C	Transferred in the laboratory from wild strains	None
2	"	A Su T		"
3	<i>S. dublin</i>	S Su T		"
4	<i>E. coli</i>	A Su T		"
5	0141: K85ab; K88ab	A S Su T N C		"
6	0147: K89: K88ac	S Su T C		S Su C
7	<i>S. typhimurium</i>	S Su C		All
8	<i>S. cholerae-suis</i>	A S Su N C		S Su C
9	<i>S. typhimurium</i>	A S Su N C		S Su N C
10	"	A S Su T N F	Naturally occurring	None
11	"	S Su T F		"
12	"	A Su T		"
13	<i>E. coli</i> untyped	A S Su T C		"
14	"	A S Su T		"
15	"	S Su T C		"
16	"	S Su T		"
17	"	S Su T		"
18	"	S T		"
19	"	A S Su T C		A C
20	"	S Su T		S Su
21	"	Su T C		Su C
22	"	A S Su T C		S Su C

Passage experiments were also performed with 10 strains of non-pathogenic *E. coli*, tetracycline-resistant, and of pig and fowl origin. No loss of resistance was noted. Neither was it noted in 12 similar experiments with *S. dublin* and *E. coli*, pathogenic for pigs and fowls, to which the resistance of some of the 10 strains of non-pathogenic *E. coli* was transferred in the laboratory.

Discussion

Though our survey was limited in size, the results indicate that infective resistance is probably the most common form of drug resistance among the *E. coli* that inhabit the alimentary tract of human beings, calves, pigs, and fowls in Britain. The fact that it has become so widespread among the human and animal *E. coli* populations during the relatively short antibiotic era, and that so many drugs are involved, makes it unlikely that the *de novo* emergence of infectively resistant organisms is a rare event. Rather does the reverse seem more probable. The demonstration of infective resistance to streptomycin, sulphonamides, tetracyclines, and chloramphenicol in two unrelated strains of *E. coli* isolated from calves as long ago as 1956 does not support a view that infective resistance is a phenomenon having risen relatively late in the antibiotic era. It is noteworthy that the calves from which the two strains were isolated were among the first in Britain to be treated with chloramphenicol.

The case against the overuse of drugs is now greatly strengthened, because it is apparent that the resistant non-pathogenic *E. coli* which frequently become dominant in the alimentary tract during drug administration are most likely to be infectively resistant. This resistance might be transferred to other non-pathogenic strains of *E. coli* that may be better adapted, in the absence of the selection pressure provided by the presence of the drug, at colonizing the alimentary tract. So long as infectively resistant strains of *E. coli* persist in the alimentary tract there is a danger of their resistance being transferred to any pathogenic member of the Enterobacteriaceae that may enter it.

Since the drug resistance of the non-pathogenic strains of *E. coli* isolated from the alimentary tract of man, calf, pig, and

fowl was transferred to the principal *Salmonella* serotypes and to some of the *E. coli* serotypes pathogenic for these four species of animal, the colonization of the alimentary tract of any one of these species with drug-resistant *E. coli* from any of the others might be considered an undesirable event. Judging from the survey this would appear to be of limited importance at the present time because there was no great difference between the incidence of resistant strains in the various species. It might well be important in the future if a variety of drugs were used excessively in one of the species. The situation developing in calves in intensive units requires watching in this respect. It is noteworthy that the ampicillin-resistant strains of *E. coli* isolated from human beings (Table I) originated from human beings who looked after intensively reared calves; the faeces of many of these calves contained ampicillin-resistant *E. coli*.

The passage experiments revealed a great difference in stability between multiple-resistant strains. It is apparent that under natural conditions in a drug-free environment some of them would soon lose at least a part of their resistance pattern. All the non-pathogenic *E. coli* strains tested that were tetracycline-resistant only, and that are representative of the *E. coli* found in pigs and fowls fed diets containing tetracyclines, showed no loss of resistance on passage. This finding was not unexpected, because previous observations (Smith and Crabb, 1957) had shown that tetracycline-resistant organisms still dominated the *E. coli* flora of the faeces of pigs three to seven months after the practice of feeding tetracyclines was withdrawn. It is noteworthy that in the present work it was possible to transfer resistance from this kind of *E. coli* to most of the *Salmonella* and *E. coli* serotypes pathogenic for man, calves, pigs, and fowls.

Summary

In a small survey conducted in 1965 drug-resistant strains of *E. coli* were fairly often found in the faeces of human beings, calves, pigs, and fowls. The resistance of most of these strains, and of two isolated in 1956, was infective.

The resistance of four non-pathogenic strains of *E. coli* isolated from the faeces of a man, a calf, a pig, and a fowl was transferred to the principal *Salmonella* serotypes, and to some of the *E. coli* serotypes that cause disease in these four species. Similar transfers were performed using as donor *E. coli* strains that were tetracycline-resistant only—strains that are typical of those that dominate the *E. coli* flora of pigs and chickens that are fed commercially on diets containing tetracyclines. The resistance of most of the strains of this type examined was infective.

The multiple resistance of three strains of *S. typhimurium* was transferred to other *Salmonella* serotypes and to pathogenic serotypes of *E. coli*.

A minority of multiple-resistant strains lost a part or the whole of their resistance pattern during serial passage in broth; this was not observed in strains of *E. coli* resistant to tetracyclines only.

Ampicillin, streptomycin, tetracyclines, chloramphenicol, neomycin, sulphonamides, and furazolidone were all found to be involved in infective drug resistance.

We wish to thank Mrs. L. Stubbs, Miss S. Harding, and Mr. A. Miller for their capable technical help, and Mrs. E. Johnson, Dr. K. C. Sellers, Dr. Joan Taylor, and Mr. J. Walton for advice and assistance in various ways.

REFERENCES

- Anderson, E. S., and Datta, N. (1965). *Lancet*, **1**, 407.
— and Lewis, M. J. (1965). *Nature (Lond.)*, **206**, 579.
Datta, N. (1962). *J. Hyg. (Lond.)*, **60**, 301.
Smith, H. W., and Crabb, W. E. (1957). *Vet. Rec.*, **69**, 24.
Watanabe, T. (1963). *Bact. Rev.*, **27**, 87.