prophylaxis is all-important, and here doctors and nurses should obviously occupy leading roles. The finding that so many members of the medical and nursing professions themselves become victims of drug-dependence and alcoholism once more illustrates the urgent need for much better coverage of these subjects in the medical and nursing curriculum.—I am, etc.,

St. Bernard's Hospital,
Southall, Middlesex.

M. M. GLATT.

REFERENCES

- Glatt, M. M., Pittman, D. J., Gillespie, D. G., and Hills, D. R., *The Drug Scene in Great Britain*, London, 1967.
- Lewin, L., *Phantastica, Narcotic and Stimulating Drugs*, London, 1964, p. 54.
- Carver, A. E., *Brit. J. Ineb.*, 1934, 31, 73.
- Brook, M. F., Hailstone, J. D., and McLaughlan, I. E. J., *Brit. J. Psychiat.*, 1967, 113, 1013.
- General register office, Dec. Suppl. Engl. and Wales, 1951, 1957. Occup. Mortal. Part II. Tables H.M.S.O. London.*
- Glatt, M. M., in *Alcohol and Road Traffic*, Proc. 3rd Internat. Confer., ed. J. D. J. Havard, B.M.A., London, 1963, p. 99.
- "Lyn" *Brit. med. J.*, 1956, 2, 101.

Psychotropic Drugs

SIR,—Recently I was called urgently to see a young woman complaining of severe headache. She had been on a monoamine oxidase inhibitor drug for a considerable time, and was under consultant psychiatric supervision for a moderately severe depression.

During the previous night she had been unable to sleep as a result of an irritating cough. In scientific parlance, she took 30 ml. of a proprietary children's linctus, freely available over the counter, and thereby ingested a total of 9 mg. of the antihistaminic agent promethazine hydrochloride. Shortly afterwards she suffered the typical pounding occipital headache of the M.A.O.I. adverse reaction.

In lay language, here was a young woman who while taking a drug for her depression at the same time took a moderate dose of a children's cough mixture, as a result of which she could in theory have died from a brain haemorrhage, leaving a motherless young family. I make no apology for the dramatic nature of the last sentence. Fortunately, she recovered fully, and I should also point out that she had been well warned about the usual agents likely to cause this reaction. I would be the first person to agree with Dr. William Sargant (27 January, p. 249) that the antidepressants have a specific action, and are not mere placebos supported by the therapeutic personality of the physician. However, from the above experience, I would suggest two things.

First of all there would appear to be a case for regarding the M.A.O.I. group as potentially very dangerous drugs, to be used solely on consultant psychiatric advice. I would suggest that the vast majority of depressive states met with in general practice can be successfully and safely treated with the tricyclic compounds such as amitriptyline and imipramine.

Next, it should behave all practitioners prescribing the M.A.O.I. drugs to impress upon their patients the need not only to avoid the well-known adverse reactors, such as cheese, broad-beans, and Marmite, but to eschew every drug and proprietary medicine

unless previously approved by their physician.—I am, etc.,

Leicester.

LOGAN MITCHELL.

SIR,—Psychotherapists should be able to take heart from Dr. William Sargant's letter (27 January, p. 249). The value of their work has often been criticized because they failed to supply statistics convincing enough to support their claims. But now we are told that even the "most refined statistics" can equally prove or disprove the alleged efficacy of a particular therapy. This being so, psychotherapists like pharmacotherapists can argue their case on the strength of bedside experience, and similarly claim gratifying results in selected patients, without bothering too much about statistics.—I am, etc.,

Weybridge,
Surrey.

I. ATKIN.

Diphenhydramine in the Treatment of Oculogyric Crisis

SIR,—The recent interest in your correspondence columns (2 December, p. 554) in the bizarre manifestations of oculogyric crises following phenothiazine therapy prompts us to record the following unusual instance in a child, and its treatment with the antihistamine diphenhydramine.

The patient, a boy aged 2½ years, was seen two hours after the ingestion of about 6 ml. of a film cement containing as a solvent 80% ethylene chlorhydrin and 20% acetone. His stomach was washed, but he continued to vomit and became increasingly drowsy and shocked. In addition to the usual resuscitative therapy amphetamine sulphate 2.5 mg. was administered orally as a stimulant, followed by a similar dose four hours later, together with an intramuscular injection of 1 mg. because of profuse vomiting. There followed an increase in respiratory and pulse rates, and a rise in blood pressure, which were attributed to the effect of amphetamine. Twenty-four hours after ingestion episodes of upward rolling of the eyes with a fixed gaze,

opisthotonos, and gustatory movements in the mouth were noticed. Neurological examination, liver-function tests, serum electrolytes, and blood urea were normal. The attacks were initially thought to be a form of petit mal (there was no previous or family history of this), and two doses of phenobarbitone 15 mg. each were administered at 12-hourly intervals. The patient had a total of six attacks and a few milder ones in the next 24 hours, when it was realized that they were in fact "oculogyric crises." Oral diphenhydramine 10 mg. six-hourly was then started, and no significant attacks followed, other than two mild ones, within the following 24 hours.

No similar neurological disturbance is known to follow amphetamine¹ or ethylene chlorhydrin. Toxic effects of the latter² include central nervous system depression, peripheral circulatory and respiratory failure, and impairment of liver and kidney functions. There was no evidence that the small dose ingested resulted in any residual deleterious effects, but its association with the oculogyric crisis is very strong. This was also the case with the striking improvement following the administration of diphenhydramine.

A similarly dramatic response was witnessed by one of us (J. A.) in a 10-year-old girl with Riley-Day syndrome (familial dysautonomia) who developed oculogyric crisis following a relatively small dose of chlorpromazine. Besides the well-known effect of the solanaceous drugs in the treatment of oculogyric crisis attention is drawn here to diphenhydramine, which is also useful in the treatment of other extrapyramidal manifestations of phenothiazine toxicity.³—We are, etc.,

DON HILSON.

J. M. ABRAHAM.

The General Hospital,
Ashton under Lyne, Lancs.

REFERENCES

- Wallace, H. E., Neumayer, F., and Gutch, C. F., *Amer. J. Dis. Child.*, 1964, 108, 657.
- Dreisbach, R. H., *Handbook of Poisoning*, 1966, 5th ed. Blackwell Scientific Publications, Oxford.
- Martindale's Extra Pharmacopoeia*, edited by R. G. Todd, 1967, 25th ed. The Pharmaceutical Press, London.

Gynaecomastia in Renal Failure

SIR,—We quite agree with the conclusions of Dr. R. M. Lindsay and others that "gynaecomastia may occur not uncommonly in male patients with chronic renal failure during the early stages of regular dialysis treatment" (30 December, p. 779).

Gynaecomastia has been documented in eight of our 22 patients receiving maintenance haemodialysis. On occasions, breast enlarge-

ment may be extreme, as shown below. Since reporting this "complication" of haemodialysis to the American College of Physicians in April 1967¹ we have learned of over 25 additional cases. In addition, lactation in women has apparently been noted following haemodialysis.²

We are also unaware of any reports specifically dealing with gynaecomastia in