rehabilitation situation that the more severely incapacitated are treated alongside the less handicapped, as this not only constitutes a stimulus to the patients but also to the staff engaged in this work.

This review of the present situation is undoubtedly timely, and one must hope sincerely that the obvious points emerging from it will neither be ignored nor be given the implication that a rash of rehabilitation centres is required. Such rapid expansion is neither economically possible nor entirely desirable. The growth of rehabilitation requires to be carefully planned not only on a regional basis but as part of a carefully designed national policy.—I am, etc.,

ALAN RICHARDSON.

Wolfson Medical
Rehabilitation Centre,
Atkinson Morley's Hospital,
London S.W.20.

Constrictive Pericarditis

SIR,—Your leading article entitled "Non-tuberculous Constrictive Pericarditis" (3 February, p. 265) states that pericardial rheumatoid nodules are pathological findings without clinical manifestations, and that histological examination in cases of rheumatoid constrictive pericarditis consistently fails to show lesions characteristic of rheumatoid arthritis.

Perhaps you will permit me to draw attention to our recently reported¹ patient with acute rheumatoid arthritis in whom cardiac tamponade necessitated thoracotomy. At operation there was a tense pericardial effusion, associated with pericardial thickening and adhesions. Histological examination of the operative parietal pericardial biopsy showed an appearance similar to that of the rheumatoid nodule. It may be that we were able to identify the characteristic histology in our patient because, in contrast with Dr. B. P. Harrold's two cases (3 February, p. 290) which provoked your leading article, our patient's pericardial disease was relatively acute, and he had not received corticosteroid therapy preoperatively.—I am, etc.,

ROGER A. L. SUTTON.

University College Hospital,
London W.C.1.

REFERENCE

- ¹ Sutton, R. A. L., *Proc. roy. Soc. Med.*, 1967, 60, 339.

Coal Gas and the Brain

SIR,—The leading article, "Coal Gas and the Brain" (17 February, p. 398), states that where it is available "exposure to hyperbaric oxygen at 2–2½ atmospheres pressure is the treatment of choice" for carbon monoxide poisoning and quotes the 100% recovery rate achieved by Smith *et al.*¹ in support of this contention. An annotation on a similar theme in the *Lancet* (13 January, p. 77) also gives this reference, and concludes that hypertonic oxygen therapy "may be vital." Garland and Pearce² quote the work of Smith *et al.*¹ as the sole evidence in strong support of this form of treatment.

We feel that these reviews give the impression that only hyperbaric oxygen will achieve a 100% recovery. The excellent results of Smith and his colleagues were obtained when "the average time elapsing between finding

the patient and his admission to hospital has been about 30 minutes," and it should be noted that the death of a patient admitted indirectly to their hospital was excluded from the results. After another three years' experience Smith¹ reported that the number of carbon monoxide poisonings treated with hyperbaric oxygen had risen to 70, with two deaths. This later work is not quoted in any of the three authoritative articles already mentioned. Moreover, "the use of hyperbaric oxygen is considerably limited by the occurrence of oxygen poisoning."³ It is difficult to compare the results of these series with others, as the "respiratory and cardiac depression" accepted by Smith *et al.* as criteria for treatment with hyperbaric oxygen are not sufficiently defined.

Lest it be thought that only hyperbaric oxygen therapy with its attendant risks of oxygen poisoning to doctors and nurses will achieve a very high recovery rate, we report the results in our last 100 patients suffering from carbon monoxide poisoning treated without hyperbaric oxygen. Ninety-nine patients survived with no after effects, and one died.—We are, etc.,

HENRY MATTHEW.

A. T. PROUDFOOT.

Regional Poisoning Treatment Centre,
The Royal Infirmary,
Edinburgh.

REFERENCES

- ¹ Smith, G., Ledingham, I. McA., Sharp, G. R., Norman, J. N., and Bates, E. H., *Lancet*, 1962, 1, 816.
² Garland, H., and Pearce, J., *Quart. J. Med.*, 1967, 36, 445.
³ Smith, G., *Practitioner*, 1966, 197, 553.
⁴ Norman, J. N., and Smith, G., *Postgrad. med. J.*, 1966, 42, 449.

Acute Abdomen

SIR,—Professor Harold Ellis in his interesting and succinct paper on the diagnosis of the acute abdomen (24 February, p. 491) is concerned with the paramount importance of reliance on clinical features, immediate decision, and accurate diagnosis. I question the wisdom of his remark: "One should be suspicious of the diagnosis of acute appendicitis without pyrexia; it does occur but it is unusual." This is dangerous teaching, for the absence of fever, and indeed of tachycardia, should never prevent a diagnosis of acute appendicitis or make for delay in reaching the important decision when to admit to hospital.

Only recently I saw a 45-year-old man with an eight-hour history of abdominal pain. Temperature was normal and pulse rate 80. I was in no doubt about the diagnosis, and a gangrenous appendix was removed the same evening. And this was not the first time I had made the diagnosis with reasonable certainty in the absence of fever.

I cannot agree that it is unusual, or so unusual as to be remarkable enough to raise doubts and suspicions of the diagnosis. For it is just these doubts and suspicions that may tempt one to temporize; and in those few hours a gangrenous appendix may perforate. I would rather advise the student or tiro to note the temperature and pulse rate, but, even if one or both are normal, in the presence of other signs and symptoms of acute appendicitis on no account to be deterred from making the diagnosis and dispatching the patient to hospital with all speed.—I am, etc.,

Sonning Common,
Nr. Reading, Berks.

PETER HEMPHILL.

Suicide in Pregnancy

SIR,—I must support Dr. C. B. Goodhart's view (3 February, p. 318) that suicide does occasionally occur in pregnancy. Although these deaths have not been specifically mentioned in the Ministry of Health's *Reports on Confidential Enquiries into Maternal Deaths in England and Wales*, reports were in fact received on 14 deaths from suicide during the twelve years 1952–63. Furthermore, it is probable that in some cases of suicide associated with pregnancy and childbirth the fact of pregnancy is not disclosed on the death certificate, and so a confidential report is not initiated.

Thanks to the co-operation of coroners and the Home Office confidential reports on these cases are now received by the Ministry of Health, and in future *Reports on Confidential Enquiries* these deaths and other deaths due to poisoning or violence will be recorded, and a more accurate estimate of the risk of suicide in pregnancy will be obtained.—I am, etc.,

London W.1.

HUMPHREY ARTHURE.

Measles Immunization

SIR,—In your leading article (17 February, p. 395) you state that the Ministry of Health has recommended that measles vaccination should be offered to all children who have not been protected by either previous immunization or an attack of the disease. I believe children are immunized in other ways—that is, by having a subclinical attack or possibly having a permanently inherited immunity. If children have passed through two epidemics they can be considered immune. The Table below will demonstrate this.

	1st Epidemic	2nd Epidemic
Number in community	580	587
„ known to have had measles	555	482
Number known not to have had clinical attack	25	105
Number known to have been exposed twice before and considered immune	11	24
Number assumed non-immune	14	81
Number attacked in epidemic	13	78
„ not attacked	1	3

This refers to boys of 14 to 18 in a public school. Of the 580 at the time of the first epidemic there were 11 who were known to have been exposed on two previous occasions, and of the 587 in the second epidemic 24 had had two previous exposures. I am sure we should study those who have been exposed as well as those who get a genuine attack, and this will be even more important when, in the foreseeable future, mumps vaccine may be recommended. The figures also show the infectivity of measles. Only one out of 580 escaped if those who have had previous attacks and those who have been twice exposed are included, and this despite the fact that the number assumed non-immune was only 14 in the first epidemic. The figures for the second epidemic are 3 out of 587. Among those who have been exposed twice I discovered that four belonged to four families totalling 18 siblings, none of whom had had measles, and, furthermore, none of their mothers had had measles. This therefore is another very small group where measles vaccination would be superfluous.