rehabilitation situation that the more severely incapacitated are treated alongside the less handicapped, as this not only constitutes a stimulus to the patients but also to the staff engaged in this work.

This review of the present situation is undoubtedly timely, and one must hope sincerely that the obvious points emerging from it will neither be ignored nor be given the implication that a rash of rehabilitation centres is required. Such rapid expansion is neither economically possible nor entirely desirable. The growth of rehabilitation requires to be carefully planned not only on a regional basis but as part of a carefully designed national policy.—I am, etc.,

ALAN RICHARDSON.

Wolfson Medical
Rehabilitation Centre,
Atkinson Morley's Hospital,
London S.W.20.

Constrictive Pericarditis

SIR,—Your leading article entitled "Non-tuberculous Constrictive Pericarditis" (3 February, p. 265) states that pericardial rheumatoid nodules are pathological findings without clinical manifestations, and that histological examination in cases of rheumatoid constrictive pericarditis consistently fails to show lesions characteristic of rheumatoid arthritis.

Perhaps you will permit me to draw attention to our recently reported¹ patient with acute rheumatoid arthritis in whom cardiac tamponade necessitated thoracotomy. At operation there was a tense pericardial effusion, associated with pericardial thickening and adhesions. Histological examination of the operative parietal pericardial biopsy showed an appearance similar to that of the rheumatoid nodule. It may be that we were able to identify the characteristic histology in our patient because, in contrast with Dr. B. P. Harrold's two cases (3 February, p. 290) which provoked your leading article, our patient's pericardial disease was relatively acute, and he had not received corticosteroid therapy preoperatively.—I am, etc.,

ROGER A. L. SUTTON.

University College Hospital,
London W.C.1.

REFERENCE

- ¹ Sutton, R. A. L., *Proc. roy. Soc. Med.*, 1967, 60, 339.

Coal Gas and the Brain

SIR,—The leading article, "Coal Gas and the Brain" (17 February, p. 398), states that where it is available "exposure to hyperbaric oxygen at 2–2½ atmospheres pressure is the treatment of choice" for carbon monoxide poisoning and quotes the 100% recovery rate achieved by Smith *et al.*¹ in support of this contention. An annotation on a similar theme in the *Lancet* (13 January, p. 77) also gives this reference, and concludes that hypertonic oxygen therapy "may be vital." Garland and Pearce² quote the work of Smith *et al.*¹ as the sole evidence in strong support of this form of treatment.

We feel that these reviews give the impression that only hyperbaric oxygen will achieve a 100% recovery. The excellent results of Smith and his colleagues were obtained when "the average time elapsing between finding

the patient and his admission to hospital has been about 30 minutes," and it should be noted that the death of a patient admitted indirectly to their hospital was excluded from the results. After another three years' experience Smith¹ reported that the number of carbon monoxide poisonings treated with hyperbaric oxygen had risen to 70, with two deaths. This later work is not quoted in any of the three authoritative articles already mentioned. Moreover, "the use of hyperbaric oxygen is considerably limited by the occurrence of oxygen poisoning."⁴ It is difficult to compare the results of these series with others, as the "respiratory and cardiac depression" accepted by Smith *et al.* as criteria for treatment with hyperbaric oxygen are not sufficiently defined.

Lest it be thought that only hyperbaric oxygen therapy with its attendant risks of oxygen poisoning to doctors and nurses will achieve a very high recovery rate, we report the results in our last 100 patients suffering from carbon monoxide poisoning treated without hyperbaric oxygen. Ninety-nine patients survived with no after effects, and one died.—We are, etc.,

HENRY MATTHEW.

A. T. PROUDFOOT.

Regional Poisoning Treatment Centre,
The Royal Infirmary,
Edinburgh.

REFERENCES

- ¹ Smith, G., Ledingham, I. McA., Sharp, G. R., Norman, J. N., and Bates, E. H., *Lancet*, 1962, 1, 816.
² Garland, H., and Pearce, J., *Quart. J. Med.*, 1967, 36, 445.
³ Smith, G., *Practitioner*, 1966, 197, 553.
⁴ Norman, J. N., and Smith, G., *Postgrad. med. J.*, 1966, 42, 449.

Acute Abdomen

SIR,—Professor Harold Ellis in his interesting and succinct paper on the diagnosis of the acute abdomen (24 February, p. 491) is concerned with the paramount importance of reliance on clinical features, immediate decision, and accurate diagnosis. I question the wisdom of his remark: "One should be suspicious of the diagnosis of acute appendicitis without pyrexia; it does occur but it is unusual." This is dangerous teaching, for the absence of fever, and indeed of tachycardia, should never prevent a diagnosis of acute appendicitis or make for delay in reaching the important decision when to admit to hospital.

Only recently I saw a 45-year-old man with an eight-hour history of abdominal pain. Temperature was normal and pulse rate 80. I was in no doubt about the diagnosis, and a gangrenous appendix was removed the same evening. And this was not the first time I had made the diagnosis with reasonable certainty in the absence of fever.

I cannot agree that it is unusual, or so unusual as to be remarkable enough to raise doubts and suspicions of the diagnosis. For it is just these doubts and suspicions that may tempt one to temporize; and in those few hours a gangrenous appendix may perforate. I would rather advise the student or tiro to note the temperature and pulse rate, but, even if one or both are normal, in the presence of other signs and symptoms of acute appendicitis on no account to be deterred from making the diagnosis and dispatching the patient to hospital with all speed.—I am, etc.,

Sonning Common,
Nr. Reading, Berks.

PETER HEMPHILL.

Suicide in Pregnancy

SIR,—I must support Dr. C. B. Goodhart's view (3 February, p. 318) that suicide does occasionally occur in pregnancy. Although these deaths have not been specifically mentioned in the Ministry of Health's *Reports on Confidential Enquiries into Maternal Deaths in England and Wales*, reports were in fact received on 14 deaths from suicide during the twelve years 1952–63. Furthermore, it is probable that in some cases of suicide associated with pregnancy and childbirth the fact of pregnancy is not disclosed on the death certificate, and so a confidential report is not initiated.

Thanks to the co-operation of coroners and the Home Office confidential reports on these cases are now received by the Ministry of Health, and in future *Reports on Confidential Enquiries* these deaths and other deaths due to poisoning or violence will be recorded, and a more accurate estimate of the risk of suicide in pregnancy will be obtained.—I am, etc.,

London W.1.

HUMPHREY ARTHURE.

Measles Immunization

SIR,—In your leading article (17 February, p. 395) you state that the Ministry of Health has recommended that measles vaccination should be offered to all children who have not been protected by either previous immunization or an attack of the disease. I believe children are immunized in other ways—that is, by having a subclinical attack or possibly having a permanently inherited immunity. If children have passed through two epidemics they can be considered immune. The Table below will demonstrate this.

	1st Epidemic	2nd Epidemic
Number in community	580	587
„ known to have had measles	555	482
Number known not to have had clinical attack	25	105
Number known to have been exposed twice before and considered immune	11	24
Number assumed non-immune	14	81
Number attacked in epidemic	13	78
„ not attacked	1	3

This refers to boys of 14 to 18 in a public school. Of the 580 at the time of the first epidemic there were 11 who were known to have been exposed on two previous occasions, and of the 587 in the second epidemic 24 had had two previous exposures. I am sure we should study those who have been exposed as well as those who get a genuine attack, and this will be even more important when, in the foreseeable future, mumps vaccine may be recommended. The figures also show the infectivity of measles. Only one out of 580 escaped if those who have had previous attacks and those who have been twice exposed are included, and this despite the fact that the number assumed non-immune was only 14 in the first epidemic. The figures for the second epidemic are 3 out of 587. Among those who have been exposed twice I discovered that four belonged to four families totalling 18 siblings, none of whom had had measles, and, furthermore, none of their mothers had had measles. This therefore is another very small group where measles vaccination would be superfluous.

I realize that measles immunization will usually be given to those from 1 to 5, but nevertheless strong pressure is being applied to recommend that children who are entering school at age 14 should be immunized.—I am, etc.,

R. E. SMITH.

Board of Graduate Studies,
Faculty of Medicine,
University of Birmingham.

Medindex

SIR,—I hasten to support Dr. J. B. Metcalfe's plea to the publishers of *Medindex* (24 February, p. 515) and join him in requesting them to resort to the old format. I would also like them to include a list of drugs, arranged alphabetically, giving the scientific names first and the trade equivalents (if any) alongside. Many readers of publications such as the *B.M.J.* must spend a considerable time trying to find out what "drug" the authors of an article are writing about. The inclusion of a short list of equivalent names at the end of each article would make life easier for the not so scientifically minded.—I am, etc.,

Skipton, Yorks.

C. H. ROBINSON.

SIR,—We have answered already through the post Dr. J. B. Metcalfe's criticisms (24 February, p. 515), but for the benefit of your other readers will you permit me to summarize briefly the position of *MIMS vis-à-vis Medindex*?

Until early in 1967, when this company took it over, *Medindex* was published quarterly. It was providing very similar information to that given in *MIMS*, which was, and still is, published monthly. We have evidence to show that *MIMS* was greatly preferred to *Medindex*. *MIMS* is intended as an *aide-memoire* to proprietary drugs. It is not a textbook of pharmacology or therapeutics. In the autumn we are to publish the first edition of the *Medindex Compendium*, which will be issued annually. This will include monographs on proprietary drugs, lists of approved names, a colour identification chart for tablets and capsules, and additional information which cannot be included in *MIMS* if it is to preserve its unique character.

We are grateful for the numerous letters of appreciation of *MIMS* and for the helpful comments that we receive every day from doctors and pharmacists in general practice and in the hospital service. The *Compendium* has been planned with the support of the pharmaceutical industry and is expected to be the most comprehensive and detailed publication of its kind.—I am, etc.,

E. N. PULLOM,
Editor, *MIMS*.

London W.1.

Selection of Medical Students

SIR,—My point (20 January, p. 176) is that some able students who would make good doctors are bad at physics and chemistry. Dr. I. M. Richardson (17 February, p. 445) would exclude them because they could not master first M.B. science, whereas I would replace first M.B. science by a general science

course. Doctors need a scientific attitude, but this, far from being a pure scientist's prerogative, can be acquired by any able student.—I am, etc.,

Whickham,
Newcastle upon Tyne.

ANDREW SMITH.

Clinicopathological Conferences

SIR,—Doctors all over the world who are regular readers of the *B.M.J.* look forward to your instructive Clinicopathological Conferences. Apart from their intrinsic merits, they give us the feeling of personally sitting in on a very high-level discussion by experts

who have developed special interests in certain subjects in their subspecialties.

However, the level of their discussion and knowledge is such that it is not always clear what specialties those taking part normally work in. For the benefit of those who are not familiar with the positions held by the staff at Hammersmith Hospital, I believe it would help in following the material presented even better if such indications as "Professor X—professor of pathology" or "Dr. Y—M.R.C. research fellow in clinical medicine" were added at least at one point.—I am, etc.,

Paediatric Department,
Ashkenol General Hospital,
Israel.

AARON BLOCH.

Deaths from Asthma

SIR,—The investigations of Dr. F. E. Speizer and colleagues (10 February, p. 335) have confirmed the increased death rate from asthma in recent years, and they have made a valuable assessment of treatment (10 February, p. 339) in fatal cases. Although the cause of this increase remains quite uncertain, I agree that the fact that death was so often unexpected can be explained in some cases by previous misassessment of the severity of the illness. A contributory factor to this may sometimes have been the still widely held belief, especially in children, that asthma is largely a psychologically determined condition. In this way the main effort of assessment and treatment may have been directed to vague and quite unhelpful psychological factors and away from the critical physical state of the patient. Reluctance to use steroids is a likely consequence of this attitude.

Bronchial lavage, as you suggest, certainly has some place in the treatment of severe asthma unresponsive to routine measures. In acute status asthmaticus it is an entirely rational preliminary to the institution of assisted respiration, and if undertaken before the situation has become too desperate may even eliminate the necessity for this procedure. In the subacute and severe chronic asthmas the results of bronchial lavage are often extremely difficult to assess. In the York-Harrogate areas, in the last two years, 45 bronchial lavages have been carried out in 40 patients by the method described by H. T. Thompson *et al.*¹ The results have not yet been fully analysed, but in a few patients there could be no doubt that the results have been remarkably beneficial. In the majority there has been some improvement, but other factors may have been involved and the improvement has sometimes been brief. In others little or no improvement has been obtained. There have been no serious complications.—I am, etc.,

Scotton Banks Hospital,
Knaresborough, Yorks.

W. H. HELM.

REFERENCE

- 1 Thompson, H. T., Pryor, W. J., and Hill, Jacqueline, *Thorax*, 1966, 21, 557.

SIR,—Your leading article (10 February, p. 329) very rightly stresses the necessity of treating severe asthma with corticosteroids. It continues giving general advice how to treat routinely emergency cases, but this

advice is in contrast to the opinion of most practitioners experienced in this field. You recommend "100 mg. of hydrocortisone should be given intravenously or intramuscularly every quarter of an hour until relief is obtained"—a tall order—to be followed by 40–80 mg. oral prednisolone "rapidly tailed off over one or two weeks." Many would feel that in such general advice intravenous aminophylline ought to have been included as preceding hydrocortisone; they would also object to the rapidly tailing off within one to two weeks, which could be easily understood as taking it off completely. This would be a rather dangerous procedure causing withdrawal symptoms and a possibly fatal outcome which I am sure you would want to avoid.

Less disquieting are some minor faults in your leader—for instance, the recommendation of a recent but not yet fully evaluated drug and the reference to irrelevant animal experiments about possible dangers of isoprenaline. I realize that leaders cannot always be faultless, but the major errors in this leader are potentially harmful and should have been avoided.—I am, etc.,

London N.3.

H. HERXHEIMER.

SIR,—Thank you for publishing the papers on the increase in deaths from asthma by Dr. F. E. Speizer and others (10 February, p. 335). It is a pleasure to read such careful reasoning applied to such well-collected information. We all make mistakes and can therefore only welcome such a clear guide about how to avoid some of them.

I was disappointed therefore to read "your" suggestions (10 February, p. 329) on "What is to be done?" because so many of them are only practical if the patient has a physician in constant attendance. The papers show that any action must be taken quickly, because a third of those who are going to die in an attack of asthma are dead within two hours of the start of it. You comment that "steroids are urgently needed in all severe exacerbations of asthma." Are we then to teach every asthmatic to call us at the start of every attack so that we can start the suggested scheme of injected steroids if we find that the attack is severe? Surely guidance about what constitutes a severe attack would be more useful, so that the patient could take oral steroids and thereby