Correspondence

Letters to the Editor should not exceed 500 words.

Vaccination Against Measles. R. M. Forrester, M.D., D.C.H.	764	Coal Gas and the Brain. J. M. S. Pearce, M.D.	767	Surgery for Intracranial Tumours. C. de Voil, M.B., and S. N. Sinha, M.R.C.P. GLAS.	769
Congenital Heart Disease. A. M. Johnson, F.R.C.P.	764	Pink Disease. Constance C. Forsyth, F.R.C.P. ED., and D. C. L. Savage, M.R.C.P.	767	Fluphenazine Enanthate in the Maintenance Treatment of Schizophrenia. J. Lowther, D.P.M.	770
Motivation of Medical Scientists. J. C. Shee, F.R.C.P.	764	Pain in the Neck and Arm. A. J. Macleod, M.D.; J. H. Cyriax, M.D.	767	Terrible Tautologies. E. C. O. Jewesbury, F.R.C.P.	770
Haemoglobin E and α-Thalassaemia. P. Fessas, M.D.	764	Pulmonary Function in Asthma. C. M. Ogilvie, M.D.	768	Personal View. W. C. Watson, M.D.; M. P. Coplans, F.F.A. R.C.S.	770
Volvulus of the Sigmoid Colon. D. A. Andersen, F.R.C.S.	765	Corticosteroids and Rosacea. G. A. G. Peterkin, F.R.C.P. ED.	768	Hospital Attachments. G. N. Golden, F.R.C.S.	771
Neurosyphilis. K. W. G. Heathfield, M.D.	765	Twelve-lead Electrocardiogram. P. H. Rees, M.R.C.P. ED.	768	General Practice—A Worth-while Career. A. C. Inwald, M.B.; R. C. Humphreys, M.D.	771
Gastrectomy and Vagotomy. H. Taylor, F.R.C.S., and D. G. A. Eadie, F.R.C.S.	766	Swallowed Dental Reamer. G. D. Vinden, D.T.M. & H.	769	Australia Next? S. McPherson, M.B.	771
Aldosterone and Electrolyte Movements in the Colon. R. Shields, F.R.C.S.	766	Aminocaproic Acid and "Proteinuria." M. W. Cooksey, M.D.S., and M. S. Knapp, M.D.	769	Real Reasons for Emigrating. F. M. Shattock, M.B.	771
Haemophilus Epiglottitis. H. G. Pledger, M.D., F.F.A. R.C.S.; J. D. Hill, F.F.A. R.C.S.	766	Poisoning by Oil of Wintergreen. R. R. Gordon, F.R.C.P.	769	Prescription Charges. C. J. R. Hart, M.B., and I. H. Redhead, M.D.; K. B. Hallam, M.B.; J. E. Godrich, M.B.; B. W. Wyllie, M.B.	772
Deaths from Asthma. A. J. Akhtar, M.R.C.P. ED.; P. A. Walford, M.B.	766				

Vaccination Against Measles

SIR,—Your leading article on measles vaccine (17 February, p. 395) points out that the effectiveness of vaccination before nine months of age is likely to be inhibited by the presence of maternal antibody. This maternal antibody is a very potent force in protecting the infant against natural infection in the first nine months of life.

Has anyone considered whether the antibody acquired by the active immunization of the next generation of mothers will be carried over in sufficient quantity to protect their infants in the first nine months? Or must we face a complete break in this normal pro-

tection mechanism? The protection afforded to the school child by measles vaccine may only modify his natural attack, which, in turn, is just as infectious to his young unprotected sibling, who will be doubly at risk because of the difficulty in recognizing measles in the modified case. (This situation frequently occurs in pertussis at the present time.)

Can anyone tell me whether this problem has been worked out or is it just awaiting the next generation and their doctors?—I am, etc.,

R. M. FORRESTER.
Paediatric Department,
Royal Albert Edward Infirmary,
Wigan.

Congenital Heart Disease

SIR,—Your leading article (9 March, p. 595) is disturbing, for in my opinion it gives a dangerously misleading impression regarding the need for investigation in patients with the Eisenmenger syndrome. The syndrome can usually be diagnosed in the clinic or at the bedside, and it is often possible from the history and physical signs to differentiate atrial septal defect from ventricular septal defect or persistent ductus arteriosus in the presence of this syndrome. The helpful work of Rees and Jefferson,¹ pointing out the features in the plain chest radiograph by which these defects may be differentiated one from another, adds useful collateral evidence.

The differential anatomical diagnosis is, however, of academic interest only, for at the present time these patients are beyond the reach of surgery. Operation is contraindicated. To suggest that "cardiac catheterization and angiocardiology are therefore usually required for final diagnosis" (of the anatomy associated with the Eisenmenger syndrome) is to suggest that every such patient must be subjected to an investigation which is by no means without risk in this

condition and which offers these patients no therapeutic opportunity.

A firm diagnosis of the Eisenmenger syndrome, made by the cardiologist from the clinical findings together with plain x-ray appearances, is at the present time sufficient. If the presence of the Eisenmenger syndrome is in doubt, then, of course, special investigation must be undertaken.—I am, etc.,

A. M. JOHNSON.
Southampton Chest Hospital,
Southampton.

REFERENCE

- ¹ Rees, R. S. O., and Jefferson, K. E. *Clin. Radiol.*, 1967, **18**, 366.

Motivation of Medical Scientists

SIR,—You quote at some length on the topic of motivation of medical scientists from the preamble to the 48th Annual Report of the Walter and Eliza Hall Institute of Medical Research (20 January, p. 132). One of the factors examined, explained, defended, and approved was the desire to win the approbation of one's peers. This laudable competitiveness is of course not confined

to medical scientists, but is a fundamental driving force in virtually all fields of human endeavour, including professional, sporting, artistic, and social. It is simply another aspect of what is popularly known as "trying to keep up with the Joneses." This impulsion must have arisen very early in vertebrate evolution, and originally possibly served a sexual purpose as in the plumose display of the male bird before the female during the mating season.

It is perhaps an oblique commentary on the Welfare State that you should hold up as a new discovery what must be one of the mainsprings of human action. More than 300 years ago Milton expressed the urge very neatly:

"Fame is the spur that the clear spirit doth
raise
(That last infirmity of noble mind),
To scorn delights, and live laborious days."

—I am, etc.,

J. CHARLES SHEE.

Bulawayo,
Rhodesia.

Haemoglobin E and α -Thalassaemia

SIR,—In their interesting article on the combination of heterozygous haemoglobin E with α -chain deficiency of intermediate severity Dr. P. Wasi and others (7 October, p. 29) discuss why their patients have an average of 14% Hb E, 8.4% Hb_A, and practically no β^A_2 , the remainder being Hb A with minor quantities of F. This pattern differs from the expected one—namely, Hb β^A_2 in amounts comparable to those observed in α -thalassaemia intermedia (Hb H disease) and an A/E ratio of 2.5–3.0 to 1, as seen in simple AE heterozygotes. A surplus of β^E chains could be expected as well; in an earlier study on similar patients¹ no β^E could be detected.

The thalassaemia syndromes have demonstrated that failure to detect an expected

haemoglobin fraction in the haemolysate may be due to its early intracellular precipitation.² This applies to the surplus of α -chains in β -thalassaemias, including the β -thalassaemia/Hb E association³; it holds in part for the surplus of β -chains in α -thalassaemia intermedia.⁴ I suggest that in Hb E/ α -thalassaemia the expected excess of β^E chains precipitates intracellularly. Indeed, Dr. Wasi and his colleagues observed a small percentage of red cells with inclusion bodies, which they attributed to Hb β^E . It is not stated whether the inclusions were performed in vivo or had developed in vitro. It is more probable that the precipitates are composed mainly, though not of necessity exclusively, of β^E chains and that higher proportions of inclusions occur in red cell precursors. A low inclusion cell count in the circulation does not necessarily imply a minor degree of haemoglobin precipitation. On the contrary, it may be due to early and extensive precipitation, leading to rapid selective removal of inclusions and/or affected cells and to enrichment in cells containing Hb γ_4 , which is the most stable tetramer of this class. If β^E chains are less stable than β^A increasingly more β^A chains will be available to combine with the short supply of α -chains and the A/E ratio will rise. Perhaps one should expect a series of tetrameric molecules β_4^E , $\beta_3^E\beta_1^A$, $\beta_2^E\beta_2^A$, $\beta_1^E\beta_3^A$, $\beta_1^E\beta_2^A\beta_1^A$, and $\beta_1^E\beta_1^A\beta_2^A$, in order of increasing stability; lability of these molecules will lead to some loss of β^A chains as well, so that finally only minute quantities of free β^A will be left over.

The suggested preferential uptake of α -chains by β^A relative to β^E also may contribute to an increase of the A/E ratio.⁴ However, in the relevant hybridization experiment β^A chains outnumbered the β^E chains from the start by a factor of four. Were competition solely responsible for the high A/E ratio, one would expect a large surplus of β^E chains.

Finally, greater instability of β^E compared to β^A is compatible with the fact that α -chain deficiency combined with Hb E is more severe, clinically and haematologically, than classical Hb H disease, although the imbalance of chain synthesis is more obvious in the latter.—I am, etc.,

PHAEON FESSAS.

Haematology Section and Laboratory,
Alexandra Hospital,
Athens, Greece.

REFERENCES

- Tuchinda, S., Rucknagel, D. L., Minnich, V., Boonyaparakob, U., Balankura, K., and Suvatee, V., *American J. hum. Genet.*, 1964, 16, 311.
- Fessas, Ph., *Intern. Symposium on Comparative Haemoglobin Structure, Thessaloniki*, 1966, p. 51.
- Fessas, Ph., *Blood*, 1963, 21, 21.
- Fessas, Ph., and Yataghanas, X., *Blood*, 1968, in the press.
- Tuchinda, S., Beale, D., and Lehmann, H., *Human Genetik*, 1967, 3, 312.

Volvulus of the Sigmoid Colon

SIR,—The review of 425 cases from Mulago Hospital, Kampala, Uganda, by Mr. J. J. Shepherd (3 February, p. 280) during the period 1949–65 suggests a higher incidence in Uganda than in the part of India from which I reported 40 cases during a five-year period in 1956.¹ However, to establish this I suggest it would be valuable if the actual percentage of volvulus cases to all surgical inpatients could be given, so that a basis of comparison is established. The

rate at the Evangeline Booth Hospital during the years 1948–53 was about 1.5% of the surgical admissions for pelvic colon volvulus and 2.5% for all varieties of volvulus.

I recently had the opportunity in Norway of studying the rate of hospital incidence in surgical patients at the Ulleval Hospital, Oslo, for common abdominal surgical diseases from 1911 to 1960.² The "time trends" in the five diseases studied—namely, appendicitis, urolithiasis, cholecystitis and cholelithiasis, inguinal hernia, and volvulus—

rose gradually from 1940 to reach a peak in 1944 of about 0.51% and then subsided gradually over the next six years to return to a level of about 0.09%, as shown in the graph. The figures from six other hospitals in southern Norway which provided information about cases of volvulus showed a similar rise during the war, but this rise was not observed in two hospitals in northern Norway, which had a low level throughout.

These observations strengthen the suggestion made by Bruusgaard³ that a compulsory

were found to be similar in pattern, although with local variations, in 10 other Norwegian hospitals from 1921 to 1960, so far as records were available, and the pattern in appendicitis, urolithiasis, and cholelithiasis was also found to have the same "time trend" in the insured population incidence in the City of Oslo 1940–60. This seems to indicate that the hospital incidence figures can be regarded as a fairly reliable guide to the population incidence of the diseases studied when the incidence figures are obtained from a general hospital serving a wide area. The rates of incidence in the Oslo City Hospital showed a remarkably interesting relationship to both the first and the second world war years, which has already been pointed out.^{2–5} Volvulus was relatively common before and during the first world war, having an incidence of about 0.9% of the total surgical inpatients. It fell to under 0.1% between the two world wars, but

change of dietary habits during both world wars, from a mixed to a mainly coarser vegetarian one, had an aetiological relationship to the condition, possibly affecting a few of that proportion of the population with a congenitally long or otherwise abnormal pelvic colon. This supports the earlier observations made in India, and suggests the usefulness of further incidence studies in relation to our dietary habits and structure which could point towards remedial measures.—I am, etc.,

Institute of Urology,
University of London,
W.C.2.

D. A. ANDERSEN.

REFERENCES

- Andersen, D. A., *Brit. J. Surg.*, 1956, 44, 132.
- Anderson, D. A., *J. Oslo City Hosp.*, 1966, 16, 101.
- Bruusgaard, C., *Surgery*, 1947, 22, 466.
- Frimann-Dahl, J., 1960, *Röntgen Examinations in Acute Abdominal Conditions*, 2nd ed. Springfield, Ill.
- Andersen, D. A., *Brit. med. J.*, 1967, 1, 366.

Neurosyphilis

SIR,—I was interested to read the paper on the changing clinical picture of neurosyphilis by Dr. R. Joffe and colleagues (27 January, p. 211). Since June 1964 I have kept a spare copy of all letters filed according to the diagnosis. During this period up until January 1968 I have seen 30 cases of neurosyphilis. The various types of disease were as follows:

Tabes dorsalis	...	10 cases
Spinal meningovascular syphilis	...	13 "
Cerebral meningosyphilis with epilepsy	...	2 "
General paralysis of insane	...	3 "
Erbs syphilitic paraplegia	...	1 case
Syphilitic labyrinthitis	...	1 "

These figures are obtained from the neurological clinics of three major hospitals in the Greater London area and one in Bedfordshire, although I left Harold Wood Hospital in March 1966. Fourteen patients were attending Whipps Cross Hospital, 10 Luton and Dunstable Hospital, 4 Barnet General, and 2 Harold Wood Hospital.

The figures are comparable to those of the Newcastle series: 7 cases in a period of nine months—and the catchment population area is very similar (900,000). As in the Newcastle area, several presented difficult diagnostic problems, and one patient had a familial muscular dystrophy as well as signs of a spastic paraplegia and Argyll Robertson