Correspondence

Letters to the Editor should not exceed 500 words.

Choice of Injection Sites. L. P. Garrod, F.R.C.P.	836
Long-term Administration of the Pill. P. C. Steptoe, F.R.C.O.G.	836
Early Diagnosis of Large Bowel Cancer. E. J. Watson Williams, M.R.C.P., and M. C. G. Israels, F.R.C.P.	837
Immunosuppression and Antibiotics. B. Steiner, M.D.	837
Influenza Virus and Carcinogenesis. C. Sirtori, M.D.	837
Deaths from Asthma. Anne U. Tothill, M.B.; B. L. P. Dalton, M.B.	837
Corticosteroids in Asthma. M. J. Greenberg, M.R.C.P.	838
Intracardiac E.C.G. in Pacemaker Electrode Insertion. K. Chatterjee, M.R.C.P., and R. Sutton, M.R.C.P.	838
Dental Anaesthesia. J. G. Bourne, M.D., F.F.A.R.C.S.	838
Hirschsprung's Disease. V. Rayner, and others	839

Occult Hypovolaemia. A. R. Tappin, F.F.A.R.C.S.	839
Pelvic Examination. D. B. Brown, F.R.C.O.G.; S. C. Simmons, F.R.C.S.	839
Abortion and Huntington's Chorea. Jean M. W. Bolt, D.P.H.	840
Moment of Death. N. S. Barnett, M.B.	840
Post-vagotomy Diarrhoea. P. B. Kiely, F.R.C.S.I.	840
Hypertension in the Tropics. E. M. F. Luke, M.R.C.P.E.D., and W. A. Renner, M.D.	840
Malaria Originating in Britain. Lieutenant-General Sir Treffry O. Thompson, D.M.	840
Adams-Stokes Attacks and Corticosteroids. C. I. Roberts, M.R.C.P., and C. F. P. Wharton, B.M.	841
Royal Malady. L. Eales, M.D., and E. B. Dowdle, M.D.	841
Complication of Tuberculous Meningitis. M. I. Drury, F.R.C.P.I., and others	842

More Cases of Scabies. Constance M. Ridley, M.R.C.P.	842
Acrosclerosis. H. J. Kingsley, M.B.	842
Status of the Representative Body. R. P. Hendry, M.R.C.S.	842
Tax Relief and Retirement Annuities. N. V. P. Seal	843
Pensions. J. Kennedy Harper, F.F.A.R.C.S.	843
Work Load in General Practice. C. Taylor, M.B.	843
"B.M.A. News." M. Goldman, M.B.	843
Real Reasons for Emigrating. J. R. F. Ingall, F.R.C.S.	843
Australia Next? A. I. Lane, M.B.	844
General Practice—A Worth-while Career. G. J. M. Moore, M.B.	844
G.M.C. Disciplinary Committee. R. M. H. Lefever, M.B.	844
Vocational Training for General Practice. S. Wand, D.C.L., M.B.	844

Choice of Injection Sites

SIR,—The case report by Drs. P. W. Harvey and G. C. Purnell (23 March, p. 744) and your editorial comment thereon (23 March, p. 721) should finally convince everyone that the buttock is no place in which to inject adrenaline. But does the argument end there? Given that this is a relatively dirty area in the bacteriological sense, why inject anything there at all? Is my impression correct that over the years, in consequence of a change in nursing rather than strictly medical practice, the buttock has displaced the arm as the regular site for injecting medicaments of almost any kind? These include small volumes of drug solutions which cause no subsequent pain or reaction and can perfectly well be accommodated almost anywhere in the body. Is there in fact any reason for resorting to the buttock except as a capacious alternative site when other muscle areas are becoming sore from repeated injections during a long course of penicillin?

Another question arising is: Why are so many injections made into muscle? It is more vascular than connective tissue, and absorption from it is thus more rapid, which is an advantage when giving therapeutic serum. But if a speedy effect is essential, why not give this intravenously? What else is there to which this advantage applies? Not "soluble" penicillin; this is absorbed and excreted all too rapidly. The slower absorption from connective tissue could be a positive advantage. Not adrenaline, at least in Drs. Harvey and Purnell's patient: the preparation used was formulated to ensure slow absorption (and had it been injected subcutaneously he might well have escaped). Various drugs apt to cause painful local reactions, including some antibiotics (polymyxin, cephalothin, and parenteral forms of tetracyclines) are supposed to be better tolerated by muscle than by connective tissue. Is this always true?

Vaccines and toxoids are often injected quite deeply, whether with the express intention of depositing them in muscle or not. It is difficult and perhaps for special reasons undesirable to avoid intramuscular injection in infants, but is it not better avoided in older children and adults, in whom the local reaction is usually more severe? Is the subsequent dispersion of the injected material by muscular contractions considered to be an advantage? It presumably causes more pain and perhaps more general reaction. The area in which such materials are deposited has to be invaded by macrophages, and this inflammatory reaction goes on eventually to

fibrosis. Why inflict this permanent injury on a muscle? Or is the reaction less effective if quietly and harmlessly localized in connective tissue?

I have asked questions rather than venturing to dogmatize because these matters are only on the fringe of areas in which I can claim any special knowledge. If I may be allowed a final question: Assuming that there are any doubts or disagreements here, should not a few well-informed people get together and try to decide whereabouts in the body and into what tissue different kinds of injection should be made?—I am, etc.,

LAWRENCE P. GARROD.

Royal Postgraduate Medical
School,
London W.12.

Long-term Administration of the Pill

SIR,—Contraception for the nulliparous teenager, whether married or not, is a matter for the advising doctor and the individual patient. However, I would like to draw attention to the possible dangers of prescribing "the pill" for long periods to these patients.

It is generally agreed that women reach the peak of their fertility at the age of 25, after which there is a steady but inevitable reduction of fertility. I have been able to carry out a number of biopsies of the ovaries by laparoscopic techniques on patients who have been on one or other of "the pills" for three years or more. Some of these patients have developed a secondary amenorrhoea of at least 12 months' duration. Changes which take place in the ovaries are those of marked thickening of the capsule, with distortion of follicles, reduction of the primordial follicles, and hyalinization of some of the ovarian arterioles. In several of the patients there has been considerable reduction in the ovarian volume to less than half the normal size, and the primordial follicles have been found to be scanty in number.

Several reports have now drawn attention to a possible relation between infertility and the use of oral contraceptives.¹⁻³ There is a growing volume of information which suggests that the secondary amenorrhoea is causally related to the oral contraceptive. Many but not all of the patients with secondary amenorrhoea can be induced to ovulate again on clomiphene therapy. In the light of our present knowledge it appears to me to be dangerous to give patients who have not proved their fertility oral contraceptive pills for more than a very short period of time, such as might be given for the treatment of menstrual disturbances and dysmenorrhoea. The long-term use of these pills carries with it a definite danger of producing irreversible sterility.

It is hoped that doctors will exercise caution in prescribing "the pill" to the young population, and that any future new contraceptive therapy will be thoroughly investigated for its short- and long-term effects. "The pill" is attractive to the public and to certain sections of the medical profession, because it is simple to administer. This very attraction appears to dull the

critical faculties of both patients and some members of the profession. If the reader is unconvinced by this attitude, may I remind him of the thalidomide tragedy?—I am, etc.,

Oldham, Lancs.

P. C. STEPTOE.

REFERENCES

- ¹ Dodek, O. I., and Kotz, H. L., *Amer. J. Obstet. Gynec.*, 1967, **88**, 1065.
- ² Whitelaw, M. J., Nola, V. F., and Kalman, C. F., *J. Amer. med. Ass.*, 1966, **195**, 780.
- ³ Shearman, R. P., *Lancet*, 1968, **1**, 325.

Early Diagnosis of Large Bowel Cancer

SIR,—Your leading article (9 March, p. 592) emphasizes the value of a simple test for blood in faeces in screening for possible causes of gastrointestinal disease including carcinoma. We have recently surveyed 53 patients referred to us with unexplained hypochromic anaemia. Gastrointestinal blood loss was measured with ⁵¹Cr tagged red cells¹ and three consecutive specimens of faeces were tested with the "Hematest" tablet (Ames Company Ltd.). Correlation was good; 22 gave negative results by both methods. In 27 cases the faeces contained more than 3 ml. of blood per 100 g., and all gave positive "Hematest" results within 100 seconds. In four cases the tablet test was doubtful and the specimens contained 1–2 ml. of blood per 100 g. of faeces. We conclude that the "Hematest" tablet is sensitive enough for the detection of most cases of significant bleeding. As no special precautions with diet are necessary, it is ideal for general-practitioner use and should be employed in all cases of hypochromic anaemia. We think that previous views² that the tablets have too low a sensitivity may have prejudiced against their routine use.

Further investigation of the 27 patients with significant bleeding resulted in the following diagnoses:

Radiologically Demonstrated		Not Radiologically Demonstrated	
Hiatus hernia ...	4	Carcinoma of jejunum ...	2
Carcinoma of stomach ...	2	Benign ulcer of ileum ...	1
Benign tumour of stomach ...	1	Intussusception of ileum ...	1
Duodenal ulcer ...	1	Haemangioma of caecum ...	1
Carcinoma of caecum ...	1	Carcinoma of caecum ...	2
Diverticulitis of colon ...	2	Diverticulum of caecum ...	1

In eight patients a diagnosis was not made. Our results indicate that lesions of the small intestine and caecum are a relatively common cause of hypochromic anaemia, and support the plea that every case demonstrated to be due to gastrointestinal bleeding demands a determined search including exploratory laparotomy.³—We are, etc.,

E. J. WATSON WILLIAMS.
M. C. G. ISRAËLS

Department of Clinical
Haematology,
The Royal Infirmary,
Manchester 13.

REFERENCES

- ¹ Owen, C. A., Cooper, M., Grindley, J. H., and Bollman, J. L., *Surg. Forum*, 1954, **5**, 663.
- ² Ross, G., and Gray, C. H., *Brit. med. J.*, 1964, **1**, 1351.
- ³ Kay, A. W., *Brit. med. J.*, 1962, **1**, 1709.

Immunosuppression and Antibiotics

SIR,—I read with interest your leading article on "South African Heart Transplant" (20 January, p. 132). You referred to the fact that the patient was barrier-nursed in a special room, and great care was taken against introducing infection, but there was no information as to whether the doctors considered the pathogenicity of the so-called "opportunistic" bacteria—*E. coli*, *Proteus*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*—and *Pneumocystis carinii* and *Candida albicans*, which can produce even fatal disease in a patient who is on immunosuppressive therapy.

There is no information as to whether they were looking for Gram-negative bacteria in the sputum, in the nasopharyngeal smear, or in blood and faeces. The broad-spectrum antibiotics give a false sense of security, but our experience shows that the *Kl. pneumoniae* is very often resistant against chloramphenicol and often against streptomycin, neomycin, and polymyxin B also.

Our opinion is that the antibiotic protection of a patient undergoing heart transplantation should be based upon these facts too. The combination of oxacillin, cephaloridin, and polymyxin B has proved to be effective against Gram-negative bacteria; and gentamicin is effective against *Pseudomonas aeruginosa*.¹—I am, etc.,

Postgraduate Medical School,
II Pediatrics Department,
Budapest, Hungary.

B. STEINER.

REFERENCE

- ¹ *Brit. med. J.*, 1967, **2**, 522.

Influenza Virus and Carcinogenesis

SIR,—Regarding the cancerogenesis from influenza virus in the lungs of mice reported by Drs. R. J. C. Harris and G. Negroni (16 December, p. 637) I should like to report the results of an investigation of my own, in which I exposed mice of the Swiss strain to the inhalation of an aerosol of *Myxovirus influenzae B* Lee at the dose of 10⁶ egg infectious doses (E.I.D.) for 15 minutes.^{1–3}

With the electronmicroscope I observed specific damage in the alveolar cells containing lamellar inclusions. In effect, the reproduction of the virus took place at the level of cell membranes (Fig.) as well as in the membranes of the apparatus of Golgi and in those of the lamellar inclusions.

According to current opinion,⁴ these lamellar inclusions produce the tensioactive substance responsible for pulmonary elasticity. I submit that the lamellar inclusion damage wrought by the influenza virus could

Influenza viruses B Lee in an alveolar cell containing lamellar inclusions. Arrows indicate virus particles. (×35,000.)

result in carcinogenesis through two distinct mechanisms. One is by favouring the stagnation of inhaled carcinogens through reduced alveolar elasticity. (The contact time of carcinogens is often a decisive factor, as exemplified by the high frequency of vesical carcinoma in aniline workers as contrasted with the rarity of tumours of the renal pelvis where the passage of urine is rapid.)

Secondly, the loss of alveolar elasticity and reduced alveolar oxygenation may lead to sclerosis, which in turn may represent a carcinogenic factor. There is currently a tendency to find similarities between tumours from plastic substances and tumours from sclerosis;^{5,6} the latter may include tumours that develop at the sites of previous burns, frostbite, chronic osteomyelitis, pulmonary tuberculosis, lupus, and the like.—I am, etc.,

CARLO SIRTORI.

Department of Pathology,
National Cancer Institute,
Milan, Italy.

REFERENCES

- ¹ Sirtori, C., in *Proceedings of the Third European Regional Conference on Electron Microscopy*, edited by M. Titlbach, Prague, 1964, p. 329.
- ² Sirtori, C., *Gerontologia (Basel)*, 1964, **9**, 239.
- ³ Sirtori, C., *Panminerva med.*, 1966, **8**, 103.
- ⁴ Clements, J. A., Hustead, R. F., Johnson, R. P., and Gribetz, I., *J. appl. Physiol.*, 1961, **16**, 444.
- ⁵ Sirtori, C., *Minerva med.*, 1963, **54**, 3686.
- ⁶ Bryson, G., and Bischoff, F., *Progr. exp. Tumour Res. (Basel)*, 1967, **9**, 78.

Deaths from Asthma

SIR,—In their article on the increase in asthma deaths Dr. F. E. Speizer and others (10 February, p. 339) point out that catecholamines were the drugs used most often immediately before the terminal episode and that isoprenaline, usually in the form of an aerosol spray, was most frequently preferred.

Any drug given in the form of an aerosol would reach the smooth muscle of the bronchi in the full dose very rapidly. If the patient repeated the inhalation sooner than had been recommended he would rapidly receive an overdose. Isoprenaline is a drug which normally has the beta actions of adrenaline. It relaxes the smooth muscle of the arterioles and the bronchi; and it increases the rate and force of contraction of the heart. However, if it is given in repeated doses at frequent intervals its action changes. It no longer further stimulates the heart; it does not dilate the blood vessels but actually constricts them.¹ Isoprenaline inhibits the smooth muscle of the uterus, but in the presence of this drug the uterus starts to escape from the inhibition and a second dose of isoprenaline now causes contraction.² The uterus therefore escapes from the effect of isoprenaline, becomes resistant to its inhibitory effect, and the action of the drug is reversed. The uterine contraction induced by isoprenaline in these circumstances is readily antagonized by Hydergine (dihydroergocornine, dihydroergocristine, and dihydroergocryptine).³

As reversal of the action of isoprenaline occurs in several different types of smooth muscle, it might also occur in the bronchi. If this is so administration of excess isoprenaline might induce bronchospasm. Endogenous adrenaline is another possible contributory factor. Normally adrenaline relaxes arterioles in skeletal muscles, but this action is reversed by pretreatment with large doses of isoprenaline.¹ Adrenaline normally relaxes