in the true sense of the word to Mandrax (methaqualone and diphenhydramine)—that is, they have increased their dose up to fourteen a day and crave for it. Both these patients show the typical inadequate personalities which have previously been described,¹ and both say that they use the drug to escape the conflicts in their lives.

In talking to my fellow consultants, they have all become aware of the problem and we would say that in this area there are probably five patients who are addicted to Mandrax.—I am, etc.,

D. R. BENADY.

Shelton Hospital,
Shrewsbury.

REFERENCE

- ¹ Clein, L. J., and Benady, D. R., *Brit. med. J.*, 1962, 2, 456.

Agranulocytosis after Trimethoprim and Sulphamethoxazole

SIR,—The combination of trimethoprim and sulphamethoxazole (Septrin, Bactrim) has recently appeared on the market. This drug combination contains a sulphonamide, and side-effects due to this component may be expected. We present brief details of a case of agranulocytosis due to Septrin.

The patient was a woman aged 21 years who has had mild asthma since the age of 2. On account of a chest infection due to *Ps. pyocyanea* and *H. influenzae* which had persisted for a month and had only partially responded to a course of treatment with colistin sulphomethate sodium (Colomycin), she was treated with Septrin 400 mg. three times a day. After 23 days' treatment she developed pyrexia (39° C.) with shivering attacks and generalized aches and pains. Two days later, while she was still at home, an urticarial rash appeared on the face, arms, and neck and spread to the chest, lower arms, and legs. She vomited. Enlarged lymph nodes were palpable in the neck, and two days later in the groins. On day 36, after a total dose of 43.2 g., Septrin was stopped. The symptoms abated, but recurred four days later. She was admitted to the ward with fever (38° C.), malaise, and generalized aches and pains. Enlarged lymph nodes were palpable in the neck and groins. The fauces were ulcerated. The spleen was not palpable.

The following day, 42 days after starting Septrin and five days after stopping the drug, the white cell count showed 2,500 leucocytes per cu.mm. (neutrophils 1%, lymphocytes 96%, atypical mononuclear cells 3%). Agranulocytic angina was diagnosed. Bone marrow aspiration on day 43 showed marked depression of the myeloid series, with toxic changes of vacuolation and intensive granulation. A Paul-Bunnell test was negative. The neutrophil count improved slowly over the next five days, during which time she was treated with penicillin and carbenicillin. She made a full recovery.

Other drugs received in the month before the onset of agranulocytosis were prochlorperazine maleate, a proprietary cough linctus containing pholcodine and phenyltoloxamine, magnesium trisilicate mixture, and promethazine hydrochloride. All these drugs were prescribed after the onset of urticaria, pyrexia, and vomiting, symptoms which started after 23 days' treatment with Septrin and seem likely to be drug effects. Of these drugs, prochlorperazine and promethazine both belong to the phenothiazine group, which is a known cause of leucopenia: but they were prescribed for only a short period of four days; the patient had not been previously

treated with either drug; and agranulocytosis due to these drugs, like that due to sulphonamides, usually occurs only after many days' exposure.—We are, etc.,

D. I. K. EVANS.

RUTH TELL.

Royal Manchester Children's
Hospital,
Pendlebury, near Manchester.

REFERENCE

- ¹ Wintrobe, M. M., *Clinical Haematology*, 1967, 6th edition, p. 1218. Henry Kimpton, London.

Cerebral Malaria

SIR,—Mr. H. L. Peake does well to remind us of the virtues of quinine in life-threatening malaria (25 January, p. 256). However, the intramuscular injection of quinine, which is an acid and necrotizing solution, was, and probably still is, widely and unnecessarily used in East Africa. This was on the mistaken assumption that antimalarials are not quickly absorbed from the alimentary tract and that febrile convulsions in infants with or without parasitaemia was evidence of "cerebral malaria." There may on occasion be a case for intramuscular injections of quinine in competent hands, but intravenous quinine is safe if it is well diluted in an intravenous drip and administered slowly. As a result of the abuse of intramuscular quinine I have seen an epidemic of "injection polio," an outbreak of "injection tetanus" not to mention many large quinine abscesses.

It should be remembered that in severe "cerebral malaria" the mortality (and the intensity of the cerebral insult) is directly proportional to the parasite density.¹ In a really heavy infection I have seen the patient die from the damage to his cerebral capillaries days after his parasites have succumbed to an appropriate schizonticide. I may add that quinine has no advantage over the 4-aminoquinolines however severe the parasitaemia, except where resistance to these drugs

is feared. The prevention of falciparum deaths is dependent upon adequate malaria prophylaxis in non-immunes, vigilance, and the early and competent examination of properly stained blood films. Practitioners unaware of the lethal and insidious potentialities of the malignant tertian parasite should remember this when consulted by patients returning by jet from the tropics. The likelihood of dying of malaria these days is, I think, greater out of the tropics than in them.—I am, etc.,

P. W. HUTTON.

New Cross Hospital,
Wolverhampton.

REFERENCE

- ¹ Field, J. W., *Trans. roy. Soc. trop. Med. Hyg.*, 1949, 43, 33.

R.A.F. Medicine—First Fifty Years

SIR,—The Middle Article (5 October, p. 48) purports to record the creation of the Air Force Medical Service, but completely omits the historical fact that Trenchard specially selected Matthew Fell of the Royal Army Medical Corps to initiate the Service.

This fact is amply recorded in various obituary notices to Lieutenant-General Sir Matthew Fell, and I would draw your attention in particular to one in the *B.M.J.* signed by H. E. W., which states among other things "in 1919 after special selection by Trenchard he was seconded to the Air Ministry from the R.A.M.C." In the *Lancet* it says, "He was the Father of the R.A.F. Medical Services, and he expressed his great joy of living to witness the fruition of his plans, and the importance of the part played by that Service in World War Two."—I am, etc.,

W. K. MORRISON.

Edinburgh 9.

REFERENCES

- ¹ *Brit. med. J.*, 1959, 1, 370.
² *Lancet*, 1959, 1, 318.

Employment of Women Doctors

SIR,—For several years anxiety has been expressed in many quarters about the contribution which women doctors are making to the provision of medical care in Britain (*Supplement*, 15 February, p. 73).

The trend is for many married women to withdraw temporarily from medicine and return to practice after some time varying from two or three to fifteen years. These women need refresher and retraining courses and appropriate posts. It is very satisfactory that the postgraduate and the employing authorities, with the active encouragement of the late Ministry of Health and the present Ministry of Health and Social Security, together with the interest and encouragement of the Medical Women's Federation, have designed schemes whereby the returner can obtain the retraining she needs. Appropriate employment, not merely in locum or "stop gap" posts, is now easier to get. All who are interested, either themselves or on behalf of others, in the retraining schemes in general

practice or for hospital specialties, should communicate with the postgraduate dean of the medical faculty of their area. In the case of London and the South-east this is the British Postgraduate Federation.

The Medical Women's Federation, either at its central office or at its regional headquarters, will be delighted to be of help, for there will still be many individual problems to be ironed out. The community needs more doctors, both men and women, and it is important that women should realize that they can now be retrained, and appropriate advice is now available. Some examining authorities have also recognized that there may be women who can qualify to sit for the specialist qualifications only if they can be trained on a part-time basis, and they are prepared to use their discretion in such cases.

The Medical Women's Federation highly commends the retraining schemes and hopes that all medical schools will make them known to their undergraduates and young