The response was encouraging, and after 90 minutes she was breathing without assistance and showing signs of returning consciousness. The endotracheal tube was no longer tolerated and was removed. After a further 30 minutes, however (2½ hours after immersion), she suddenly became extremely cyanosed with signs of gross pulmonary oedema. She was transferred to the Royal Devon and Exeter Hospital.

On arrival at the hospital she was barely conscious, and had extreme central cyanosis and severe pulmonary oedema. She was hypothermic, her heart was fibrillating at a rate of 80 per minute, and her blood pressure could not be measured. Endotracheal intubation was again performed, the air passages cleared by suction, and oxygen was administered. She was given ampicillin, hydrocortisone, frusemide, and aminophylline intravenously and morphine intramuscularly, and transferred to the intensive care unit, where intermittent positive pressure ventilation was commenced. A Cape respirator was used (stroke volume 750 ml., pressure 20 mm. Hg., rate 20/minute). Examination of the blood at this time showed haemoglobin 120 g./100 ml., P.C.V. 55, W.B.C. 4,500, Na 152, K 4.0, Cl 110, urea 38. It was necessary to continue intermittent positive pressure ventilation for 54 hours, since, when on two occasions it was discontinued, the patient rapidly became cyanosed. Sedation was provided by the administration of morphine or phenoperidine. An endotracheal tube (nasal, Portex 9 mm.) was left for a further 112 hours to facilitate bronchial suction and efficient oxygen therapy. During this time auscultation revealed a steady decrease in pulmonary oedema, which was confirmed by radiological examination.

After eight days her condition allowed her transfer to the general medical ward. Her continued progress was interrupted by repeated melaena stool as a result of bleeding from a large hiatus hernia which required blood transfusion. She was finally discharged six weeks after her admission.

We are grateful to Dr. A. J. Daly for permission to report this case.

—We are, etc.,

W. BOND-TAYLOR.
M. G. HALFORD.
P. JENNINGS.
K. J. POWELL.
R. WHITTLESEY.

Royal Devon and Exeter Hospital,
Exeter, Devon.

Jaundice and Methyldopa

SIR,—Jaundice has been reported as a rare complication of methyldopa therapy, and abnormalities in liver function tests are occasionally found in patients taking the drug. Only three such cases have been reported^{1,2} and one case with abnormal liver function tests but without frank jaundice has also been reported.³

A 74-year-old man was first seen in 1961, when he was found to be hypertensive. He was a publican and took large amounts of alcohol. His liver function tests were within normal limits at that time. They were repeated two years later and were again normal. His blood pressure was controlled on reserpine and hydroflumethiazide.

He was not seen again until his admission to Hounslow Hospital on 29 October 1968. At the end of August 1968, for the first time, methyldopa 250 mg. per day had been prescribed. He was taking no other drugs with the methyldopa. Prior to that he was taking guanethidine. In mid-October 1968 he noticed that he was jaundiced and that his urine was dark. His stools were of normal colour. He continued to take the methyldopa until he was

admitted to hospital. Meanwhile the jaundice was becoming progressively deeper and his urine darker. He had had no symptoms prior to the onset of the jaundice; he was apyrexial and he remained symptom free during the whole period of his jaundice. He had not been in contact with any known case of hepatitis and had not received any transfusions or injections in the past.

On examination he was deeply jaundiced. His liver and spleen were not palpable and there was no abdominal tenderness. Blood pressure was 190/110. Haemoglobin 117%, W.B.C. 3,600/cu. mm. (differential count normal), E.S.R. less than 1 mm./hr., serum bilirubin 17.6 mg./100 ml., alkaline phosphatase 16.6 King-Armstrong units/100 ml., thymol turbidity 0.5 units; total protein 6.3 mg./100 ml.—albumin 3.8 mg./100 ml., globulin 2.5 mg./100 ml. Unfortunately the alanine transaminase was not estimated at that time. His urine showed increased urobilinogen and bilirubin was present.

The methyldopa was discontinued immediately and the serum bilirubin level dropped to 8.9 mg./100 ml. in seven days, to 5.1 mg./100 ml. in 13 days, and 4.5 mg./100 ml. in 16 days. The alkaline phosphatase was 9 King-Armstrong units/100 ml. in 13 days, and aspartate transaminase level at that time was 71 units/ml. The alanine transaminase one month after the onset of the jaundice was 79 units. Liver function tests returned to normal on 2 December 1968—that is, about four and a half weeks after discontinuance of the methyldopa.

In the absence of any constitutional disturbance or contact with known cases of hepatitis, injections, or blood transfusions and the rapid improvement on discontinuing the drug, no other explanation for his jaundice could be found other than a reaction to methyldopa. The fact that the patient developed jaundice on a very small dose of methyldopa (25 mg. daily), while other patients may take up to 3 g. daily for long periods and never develop jaundice, suggests a hypersensitivity reaction. One could suggest that, being a heavy drinker in the past, alcohol might have caused liver damage and this predisposed to the onset of the hepatocellular jaundice. However, his liver function tests previously were within normal limits.

We should like to thank Mr. S. A. Taylor, F.P.S., and Dr. D. C. P. Brown, of Merck Sharp & Dohme Ltd., for their kind assistance in preparing this report.

—We are, etc.,

R. WYBURN-MASON.
C. ANASTASSIADES.

Hounslow Hospital,
Hounslow, Middlesex.

REFERENCES

- Williams, E. R., and Khan, M. A., *J. Ther. clin. Res.*, 1967, **1**, (6) 5.
- Morin, Y., Turmel, L., and Fortier, J., *Amer. J. med. Sci.*, 1964, **248**, 633.
- Irvine, R. O. H., O'Brien, K. P., and North, J. D. K., *Lancet*, 1962, **1**, 300.

Haemophilus influenzae Type B Septicaemia

SIR,—We would like to report another case to emphasize the point made by Dr. R. J. Farrand (18 January, p. 150) that this condition may not be as rare as previously believed.

A 6-month-old girl was admitted in February 1969 under the joint care of Dr. E. M. Belton and Mr. A. Graham Apley, with a history of fretfulness for two days and reluctance to

move the left arm for one day. Attempts to move the arm resulted in screaming, but there were no other symptoms. The mother had been treated recently for a septic finger. On admission the child was pyrexial (101° F.; 38.4° C.), and the left arm was hot and swollen with acute tenderness above the elbow. The haemoglobin was 10.1 g./100 ml., E.S.R. 47 mm./hr., and white blood count 24,000/cu. mm. (45% neutrophils). X-ray examination showed no bony abnormality but an irregular swelling of the subcutaneous soft tissue. A diagnosis of osteomyelitis of the humerus was made, and blood cultures were taken which three days later yielded a growth of *H. influenzae* type b sensitive to ampicillin and erythromycin.

Treatment was started with intramuscular ampicillin and cloxacillin, but the latter was replaced by erythromycin when the haemophilus was isolated. By this time the condition was responding rapidly without the bone needing to be drilled, and the arm was being freely moved. Ten days after admission x-ray examination showed early periosteal new bone formation over the lower humerus. The child has now been discharged home. —We are, etc.,

R. J. EVANS.
A. J. FRANKLIN.

St. Peter's Hospital,
Chertsey, Surrey.

Through-knee Amputations

SIR,—May I comment very briefly on Mr. P. F. Early's letter (18 January, p. 184)? Given initiative and modern technology, a prefabricated domed end containing a hydro-elastic pad could be supplied for all through-knee stump sockets. It should occupy no more than 1 in. (2.5 cm.) below the stump, and give complete padded end-bearing.

With respect to Mr. Early, his other measurements are not relevant to through-knee amputation, where the axis of the joint must lie above the condyles and the mechanism must be laterally placed. This is not difficult to arrange with modern technology; all that is needed is initiative and prefabrication. Adequate module prefabrication would cut down the time of supply of artificial limbs very significantly.—I am, etc.,

Derby.

R. LUNT.

Herniorrhaphy

SIR,—Mr. O. K. Gibbon (18 January, p. 187) writes that the results of a widespread return to basic principles in the teaching and practice of inguinal hernia repair would show that there is no need for the introduction of yet more new operations, as he condemns the operation recommended by Mr. L. F. Tinckler (28 December, p. 832). It has been the return to basic principles, however, which has allowed more and more surgeons to take advantage of the preperitoneal approach in curing groin hernias.

The preperitoneal muscle-splitting approach has all the requisites for success listed by Mr. Gibbon, and Mr. Tinckler is right (in my opinion at least) in suggesting that the insertion of a prosthesis in the preperitoneal plane is a logical solution for some inguinal hernias. The approach is no innovation, having been used as long ago as 1943 for femoral hernias by the late Mr. Rupert Vaughan Hudson, of the Middlesex Hospital, London.¹ We followed 175 femoral hernias repaired with the Vaughan Hudson tech-

nique between three and 15 years before without discovering a recurrence, and the method for inguinal hernias has also been enthusiastically described.²

I believe that if Mr. Tinckler uses a muscle-splitting incision (instead of a low midline one)—we found that in our survey the McEvedy (pararectal) approach suffered from the incidence of incisional weakness, and the Henry (midline) operation often gave poor access in addition—Mr. Gibbon might be able to accept that the preperitoneal prosthesis (which is not commonly needed when repairing inguinal hernias by this method in any case) is a logical solution. May I recommend a muscle-splitting preperitoneal approach for operations on inguinal and femoral hernias to others who have not yet used it? Its advantages are well listed where the other operations are also, of course, fully described.²—I am, etc.,

W. M. KEYNES.

The Radcliffe Infirmary,
Oxford.

REFERENCES

- 1 Keynes, W. M., and Withycombe, J., in *Hernia*, ed. L. M. Nyhus and H. N. Harkins, 1964, pp. 309–314. Lippincott, Philadelphia.
- 2 *Hernia*, 1964, ed. L. M. Nyhus and H. N. Harkins. Lippincott, Philadelphia.

Natural and "Induced" Sleep

SIR,—Dr. J. I. Evans and his colleagues (2 November, p. 291) have reported their findings that the administration of a barbiturate as a hypnotic depressed rapid eye movement (R.E.M.) sleep. The work of Kales *et al.*¹ was cited as having shown that methaqualone also depressed this phase of sleep.

Dr. J. P. Duffy, Mr. J. W. Osselton, and I have carried out a study comparing, in 14 volunteers, continuous E.E.G. recordings of a single night's natural sleep with similar recordings of Tuinal (1 capsule) and Mandrax (1 tablet) induced sleep. (Tuinal capsules contain 100 mg. quinalbarbitone sodium and 100 mg. amylobarbitone sodium; Mandrax tablets contain 250 mg. methaqualone and 25 mg. diphenhydramine.)

Our results show that in Tuinal-induced sleep there is a highly significant reduction in R.E.M. phase ($p < 0.01$) as compared with natural sleep, while there is no statistically significant difference in the occurrence of R.E.M. between natural sleep and Mandrax-induced sleep.

It is intended that a detailed report of our work will be published in due course.—I am, etc.,

K. DAVISON.

Newcastle General Hospital,
Newcastle upon Tyne.

REFERENCE

- 1 Kales, A., *et al.*, presented at A.P.S.S. Meeting, March 1968.

Perinatal Mortality Survey

SIR,—Professor Douglas Hubble has written a most interesting review of the second report of the British Perinatal Morality Survey (22 February, p. 495).

In the last paragraph he refers to all those engaged on "this massive inquiry" but makes no mention of midwives. Since the questionnaires on the 17,000 births occurring during the week of the survey were all com-

pleted by midwives it is obvious that their contribution to this great enterprise was of vital importance. The second Report will be of great interest to them as well as to obstetricians, paediatricians, social workers, and teachers.—I am, etc.,

AUDREY WOOD,
General Secretary,
Royal College of Midwives.

London W.1.

Corticosteroid Therapy and Growth

SIR,—Your leading article entitled "Corticosteroid Therapy and Growth" (15 February, p. 393) does not mention the possibility that some steroid preparations may be more growth-inhibiting than others. The change from betamethasone to prednisolone in equivalent doses in three patients under my care has resulted in an increase in the rate of growth, and I have wondered whether other clinicians have noticed anything similar.—I am, etc.,

IAN G. WICKES.

St. Andrew's Hospital,
Billericay, Essex.

Gastrointestinal Haemorrhage and Aspirin

SIR,—Dr. A. J. Quick (1 February, p. 315) states that "aspirin appears to act by preventing the vascular contraction after mechanical injury of the micro-circulation." The following case appears to support his statement.

A 5-year-old boy had acute tonsillitis with high fever and generalized aches and pains. Oral penicillin for the tonsillitis and a 150 mg.-aspirin tablet for the pain were prescribed. A few hours later he had a haematemesis, losing about a half-pint (500 ml.) of blood. It was assumed that he was sensitive to aspirin.

One month later he was admitted to hospital for tonsillectomy. His record was clearly marked that he was sensitive to aspirin. Nevertheless the routine post-tonsillectomy aspirin gargle was given. Twenty minutes later he bled severely from the tonsillar bed and nearly died.

The potentially dangerous local action of an aspirin gargle is worthy of note. It is suggested that its use for this purpose be discontinued.—I am, etc.,

London E.10.

J. SAPERIA.

Tetracycline and Nystatin

SIR,—In commenting on the letter by Dr. K. M. Citron and his colleagues (1 March, p. 575) in answer to some of my criticisms (28 December, p. 829) of their paper (16 November, p. 411), I shall confine myself to the principal issues. In their original report these authors define the aim of their large multicentre trial as (a) to assess the frequency of side-effects of tetracycline given in a dose of 2 g. daily, (b) to determine whether side-effects could be prevented or reduced by the addition of nystatin (no reference made to the relevance of its dose), and (c) to correlate these clinical findings with the results of examination of the stools for *Candida* species.

One of the main and, if correct, most important conclusions after evaluation of their data was that there is not any association between *Candida* and gastrointestinal symptoms normally attributed to "chemotherapy." The absence of

quantitative methods, the large-scale inclusion of patients already suffering from gastrointestinal manifestations following earlier antibiotic therapy, and the failure to use nystatin on its own in varying doses invalidate in my opinion this conclusion. In their letter Dr. Citron and his colleagues now admit a more modest aim of their trial—that is, to determine whether the use of nystatin (in the form of Mysteclin) reduced the incidence of side-effects of tetracycline given for 10 days in acute exacerbations of chronic bronchitis.

One cannot help but regret that the Clinical Trials Subcommittee of the British Tuberculosis Association confined itself to such a limited study and then saw fit to draw sweeping general conclusions on issues which had been investigated either inadequately or not at all.

Whether, or to what extent, the manufacturers' claims in relation to Mysteclin are justified can be assessed only in relation to the conditions of a particular clinical trial, but I was under the impression that the relatively small dose of nystatin in Mysteclin (250,000 units) was expected to prevent the increase of *Candida* in the alimentary tract during tetracycline treatment and the concomitant gastrointestinal upset. It seems to me unreasonable to expect such a small dose of nystatin also to cure already existing candidosis, not only of the bowel but also of the mouth, genital tract, perineum, and natal cleft. I have in fact seen in several hundred patients fewer side-effects from oral antibiotics, including not only tetracycline but also cloxacillin and lincocin, by giving simultaneously 500,000 units, or, in some cases, 1,000,000 units of nystatin t.d.s.

The need for brevity prevented me from explaining in my earlier letter what I meant by "clearing" patients of an "excess" of *Candida* before admitting them to the kind of trial under discussion. I tried to convey that it is almost impossible to "clear" the alimentary tract of *Candida*, but it is nearly always possible to diminish the *Candida* population to such an extent that only repeated stool cultures show a scanty presence of *Candida* and gastrointestinal symptoms completely or almost completely disappear. This can only be achieved by treating such patients with amphotericin B lozenges, six to be sucked daily for one week after dentures have been removed, with nystatin 500,000 or 1,000,000 units t.d.s. for at least three weeks, and female patients also with nystatin pessaries, 1 nightly for three weeks. Any ano-genital skin lesions have to be treated at the same time with topical nystatin.

I, and evidently also Professor H. I. Winner (18 January, p. 186) feel that only after such preparations are patients who still have gastrointestinal side-effects from previous antibiotic therapy suitable for a "clean" trial comparing tetracycline with a combination of tetracycline and nystatin.

The relationship between antibiotic therapy and candidosis has been critically reviewed by Dr. Mildred Seelig in two papers^{1,2} listing over 300 useful references, including contributions concerning most of the immunological techniques used in efforts to shed light on the complex field of widely varying host reactions to *Candida*.—I am, etc.,

G. HOLTI.

Department of Dermatology,
Royal Victoria Infirmary,
Newcastle upon Tyne.

REFERENCES

- 1 Seelig, M. S., *Bact. Rev.*, 1966, 30, 442.
- 2 Seelig, M. S., *Amer. J. Med.*, 1966, 40, 887.