serotype *canicola* in the same way; and this aspect is under continuous review.—We are, etc.,

L. H. TURNER.

Leptospirosis Reference Laboratory (P.H.L.S.; W.H.O./F.A.O.) at London School of Hygiene and Tropical Medicine, London W.C.1.

A. F. MOHUN.

Clinical Pathology Unit,
Whittington Hospital,
London N.19.

REFERENCES

- 1 Heath, C. W., Alexander, A. D., and Galton, M. M., *New England Journal of Medicine*, 1965, 273, 857; 915.
- 2 Sturdza, N., Elian, M., and Tulpan, G., *Archives Roumaines de Pathologie Expérimentale et de Microbiologie*, 1960, 19, 571.
- 3 Sturdza, N., and Elian, M., *Archives Roumaines de Pathologie Expérimentale et de Microbiologie*, 1961, 20, 33.
- 4 Elian, M., and Nicoară, I., *Bulletin of the World Health Organization*, 1964, 31, 359.

Syringomyelia

SIR,—Your leading article on syringomyelia and cavities in the cord (27 December, p. 759) neglects, I think, some important points. The use of the term "syringomyelia" throughout medical literature is confused by a lack of a clear definition because of lack of understanding of the pathological processes concerned. Perhaps the inclusion of many conditions under the general heading of "syringomyelia" is justified in the present state of knowledge. There is, however, certainly a case for differentiating from the heterogeneous accumulation of diseases which have at various times been called "syringomyelia" at least one specific and clear-cut syndrome, which I have called "communicating syringomyelia".¹

The presence of a communication between the syrinx and the fourth ventricle has long been known and the great frequency and importance of this communication in pathogenesis have been stressed.²⁻⁷

This communication is invariably accompanied by an obstruction or partial obstruction in the subarachnoid pathways linking the intracranial cerebrospinal fluid compartments with the spinal subarachnoid space. It seems probable that intracranial pulsation is thereby directed down the communication and produces a transmural pressure gradient across the wall of the cord. This would account for the gradual and progressive destruction of cord tissue in this disease.

There is therefore a well-defined triad recognizable and worthy of redefinition. This consists of: (1) clinical evidence of intramedullary cord destruction with a cavity; (2) partial blockage of the cerebrospinal fluid pathways between the intracranial compartments and the spinal subarachnoid space; and (3) a communication between the fourth ventricle and the cord cavity.

I think that it is unwise to claim this syndrome as "true syringomyelia," since this would demand a very radical reclassification of books, articles, specimens, case-records, and, perhaps most important, the thoughts of men. Clearly this is responsible for a great deal of what has been called "syringomyelia" in the past. I suggest the term "communicating syringomyelia" as being clear and preserving the historical associations of the noun.

To define "syringomyelia" as Ellertsson⁸ attempts to do on the basis of x-ray findings seems to be unsatisfactory. All the cases which he defines as "syringomyelia" seem to be "communicating syringomyelia." It is the communication which allows the cysts to become slack when the patient is not performing movements of venous distension—for example, when relaxed on the x-ray table. It is the communication also which allows the pressure to build up again after the aspiration of fluid from the syrinx, thus giving rise to the phenomenon which was commented upon in your leader—that is, that aspiration of such cases is not beneficial. Not mentioned in your leading article was another publication by Ellertsson and Greitz⁹ in which they deliberately demonstrated the existence of the communication in the same series of patients by isotopic studies with injection of radioactive iodine-labelled albumin.

I would disagree with your last paragraph in which you state that the aetiology of syringomyelia remains obscure. The communicating variety has been demonstrated and treated in neurosurgical clinics all over the world, and though it may well be true that some cystic cavities have an obscure aetiology some of them have a mechanical and remediable cause. It is vitally important that persons having the care of such patients should refer them for neurosurgical investigation and treatment.—I am, etc.,

BERNARD WILLIAMS.

Combined Neurological Service,
Royal Infirmary,
Hull, Yorks.

REFERENCES

- 1 Williams, B., *Lancet*, 1969, 2, 189.
- 2 Gardner, W. J., and Angel, J., *Clinical Neurosurgery*, 1959, 6, 131.
- 3 Gardner, W. J., *Journal of Neurology, Neurosurgery and Psychiatry*, 1965, 28, 247.
- 4 Newton, E. J., *Annals of the Royal College of Surgeons of England*, 1969, 44, 194.
- 5 Appleby, A., Foster, J. B., Hankinson, J., and Hodgson, P., *Brain*, 1968, 91, 131.
- 6 Conway, L. W., *Journal of Neurosurgery*, 1967, 27, 501.
- 7 Foster, J. B., Hodgson, P., and Pearce, G. W., *Brain*, 1969, 92, 25.
- 8 Ellertsson, A. B., and Greitz, T., *Acta Neurologica Scandinavica*, 1969, 45, 418.

Wigs and Waste

SIR,—We also have the same problem as Dr. Anne E. McCandless (24 January, p. 235) in obtaining from the National Health Service, temporary wigs for children.

During the last twelve months we have treated at least eight children with either chemotherapy for leukaemia or radiotherapy for primary brain tumours or metastases who have subsequently lost their hair. Our aim is to get these children home as soon as possible and back to a normal life, possibly even back to school. This is rendered impossible if the child is bald. Some of these children may only have a few more months of symptom-free existence left, but one aims to make these as normal and as happy as possible. Not only does an N.H.S. wig cost between £35 and £40, but it also takes up to three months to obtain one. A perfectly satisfactory acrylic wig can be bought the same day for under £5, but the N.H.S. will not at present pay for this. The Department of Health and Social Security hope-

fully suggests that "free monies" may be available. Surely such a practical, economical, and time-saving alternative to the expensive wigs at present provided at the taxpayers' expense should be provided under the N.H.S.

This does not only apply to children having treatment. The delay in getting an N.H.S. wig for an adult often means that by the time the wig arrives the hair is already growing again and the wig is hardly used.—I am, etc.,

JANE V. BOND.

Westminster Hospital,
London S.W.1.

Proteins and Insulin Release

SIR,—The experiments of Dr. R. J. Jarrett and his colleagues reported recently in your journal (6 December, p. 598) show that exogenous secretin or pancreozymin enhances the increase in blood insulin in response to intravenous infusion of amino-acids in man, an effect which has already been amply demonstrated.^{1,2} Their observation that the concurrent administration of both hormones together with amino-acids has more than an additive effect is only evident in one of their two subjects (Fig. 2) and must await confirmation.

Changes in glucagon-like immunoreactivity in the previous studies suggested that stimulation of glucagon-release by pancreozymin is associated with prolonged enhancement of insulin secretion. Changes in the blood glucose suggested that maintenance of the glycaemic stimulus to insulin secretion by the glycogenolytic action of glucagon was involved. Similar observations in the dog have led to the same conclusions.^{3,4} It is therefore of interest that the insulin responses to pancreozymin with or without secretin in the subjects of Jarrett and his colleagues were not apparently related to changes in the blood glucose, but it is difficult to interpret the data in the absence of information about the blood amino-acid levels.

The physiological importance of an alimentary stimulus to insulin during the absorption of amino-acids from the intestine in man has been inferred from the effects of intravenous or enteric infusions of arginine.^{5,2} In these studies the blood levels of glucose and amino-acids were matched, but only the enteric infusions provoked maintained stimulation of insulin secretion, and it was suggested that pancreozymin may mediate this effect.

With respect to the role of gut hormones in the release of insulin after ingestion of glucose, evidence has been reported which conflicts with that cited by Jarrett and his colleagues. Enhancement of the insulin response to intravenous glucose has been provoked by acidification of the duodenum in two further studies in man, and reasons for the absence of this effect in experiments with achlorhydric subjects have been put forward.^{6,2} The validity of pancreatic exocrine responses as indexes of release of secretin or pancreozymin in all circumstances must be questioned. Studies with a radioimmunoassay for secretin have shown that glucose taken by mouth provokes an increase in blood levels of this hormone

sufficient to cause marked potentiation of insulin release.^{7,8} A radioimmunoassay for pancreaticozym has shown no response of this hormone after ingestion of 50 g. of glucose, but a marked response occurs after a protein meal or a very large oral load of glucose (200 g.),⁹ suggesting that pancreaticozym participates in the insulin-release mechanism in these situations. Since there is evidence that the ingestion of fats is capable of enhancing the rate of disposal of IV glucose,¹⁰ it is possible that insulin secretion is potentiated by the release of pancreaticozym that would be expected with this stimulus,¹¹ but direct evidence on this point is lacking.

The accumulated evidence suggests that secretin and possibly "gut" glucagon¹² may account for augmentation of the insulin response after a moderate-sized oral glucose load, and that pancreaticozym may take part after a protein meal or a very large glucose load. There remains the problem whether these hormones are responsible for all the features that distinguish responses to ingested nutrients from responses to intravenous nutrients. These features include the relatively low blood levels of metabolites attained after ingestion of nutrients, which in the case of glucose are not readily accounted for by hepatic uptake of the substrate or by potentiation of insulin secretion.^{2,8}—We are, etc.,

JOHN DUPRÉ.

Royal Victoria Hospital,
Montreal.

D. J. CHISHOLM.

Garvan Institute, Sydney,
and Royal Victoria Hospital,
Montreal.

REFERENCES

- 1 Dupré, J., Curtis, J. D., Waddell, R. W., and Beck, J. C., *Proceedings of the Royal Society of Medicine*, 1968, **61**, 815.
- 2 Dupré, J., Curtis, J. D., Unger, R. H., Waddell, R. W., and Beck, J. C., *Journal of Clinical Investigation*, 1969, **48**, 745.
- 3 Unger, R. H., Ketterer, H., Dupré, J., and Eisentraut, A. M., *Journal of Clinical Investigation*, 1967, **46**, 630.
- 4 Ohneda, A., Parada, E., Eisentraut, A. M., and Unger, R. H., *Journal of Clinical Investigation*, 1968, **47**, 2305.
- 5 Dupré, J., Curtis, J. D., Waddell, R. W., and Beck, J. C., *Lancet*, 1968, **2**, 28.
- 6 Chisholm, D. J., Young, J. D., and Lazarus, L., *Journal of Clinical Investigation*, 1969, **48**, 1453.
- 7 Young, J. D., Lazarus, L., Chisholm, D. J., and Atkinson, F. F. V., *Journal of Nuclear Medicine*, 1968, **9**, 641.
- 8 Kraegen, E. W., Chisholm, D. J., Young, J. D., and Lazarus, L., *Journal of Clinical Investigation*, in the press.
- 9 Young, J. D., Lazarus, L., Chisholm, D. J., and Atkinson, F. F. V., *Journal of Nuclear Medicine*, 1969, **10**, 743.
- 10 Tantibhedhyangkul, P., Hashim, S. A., and Van Itallie, T. B., *Diabetes*, 1967, **16**, 796.
- 11 Wang, C. C., and Grossman, M. I., *American Journal of Physiology*, 1951, **164**, 527.
- 12 Unger, R. H., Ohneda, A., Valverde, I., Eisentraut, A. M., and Exton, J., *Journal of Clinical Investigation*, 1968, **47**, 48.

Culdocentesis in the Tropics

SIR,—May I comment on your leading article (24 January, p. 187) and the report of Dr. Cynthia Lucas and Dr. A. M. Hassim from Zambia (24 January, p. 200) dealing with culdocentesis in the diagnosis of ectopic pregnancy? The routine use of culdoscopy or aspiration or both of the peritoneal cavity above the inguinal ligament has proved so effective in the diagnosis of doubtful acute and semi-acute abdomens that it is practised by the majority of doctors in southern Nigeria. It is likely that the widespread use of intraperitoneal infusions^{1,2} in the tropics has reduced the fear of puncturing this

membrane, since bowel or organ damage or the introduction of infection is remote, given adequate precautions and intelligent selection of cases. Laparoscopy is of no practical value to the busy doctor running a one-man hospital because it demands time, general anaesthesia, and considerable expertise, none of which may be available.—I am, etc.,

DOMINIC S. COLBERT.

University College Galway, Ireland,
and General Hospital, Abakaliki,
East Central State, Nigeria.

REFERENCES

- 1 Jelliffe, D. In *Medical Care in Developing Countries*, edited by M. King, chapter 15, section 10, Nairobi, Oxford University Press, 1966.
- 2 Walter, O. In *Medical Care in Developing Countries*, edited by M. King, chapter 23, section 17, Nairobi, Oxford University Press, 1966.

Proportional Representations

SIR,—The hospital junior doctors have been the first to suffer from the lack of confidence in their negotiating machinery initiated by the Representative Body when it adopted the proportional method of electing its members. Hospital doctors believed that they could not get adequate representation of their views by straight elections and so the present complicated method was evolved. Confidence in straight elections having been eroded, it is natural that hospital junior doctors should follow suit and demand proportional representation.

Whether it is possible to satisfy these demands is doubtful, but what about the claims of all the other sections of the medical profession? I used to represent the whole of my division but now I am forced to

appear as primarily the representative of general practitioners, and many members have no primary representative at all. Under either method, of course, representatives vote on matters affecting all members.

The advocates of sectional-interest voting at elections have had their trial run and it is already proving devious, and this will be reflected in the Representative Body.

If the latter is to keep its position as the "parliament" of the B.M.A. the electorate will have to give strong support to the reinstatement of straight elections.—I am, etc.,

J. M. W. SEDGWICK.

Ware,
Herts.

General Practitioners and the District Hospitals

SIR,—Reading your leading article on general practitioners and the district hospital (3 January, p. 2), I was, unlike Dr. S. Jenkins (24 January, p. 240), relieved at the realism expressed concerning the general-practitioner participation. As a general practitioner I cannot understand why G.P.s should wish to work in hospital when there is such a vast, interesting, expanding field open to us in general practice, taking up all time and effort that we care to give. Personally I have no more wish to open sebaceous cysts, tend other doctors' hypertensives or diabetics or rheumatoids, than I have to help our excellent local chemist with his dispensing—I have enough of my own. Young doctors emigrate, but I doubt if it is because G.P.s are not looking after their own patients in hospital. I suspect it is because either they have no idea how interesting general practice can be or they see the burdens of work imposed on our consulting colleagues in the peripheral hospitals, who haven't even a decent secretariat to help them.

It seems to me that we are well ahead of other countries, including the United States, in our organization of medical practice, developing a home/G.P.-based health ser-

vice, with the obsolete idea of bed rest for essential investigations disappearing. If only we could have a quick rapport with the hospital (and we are getting this), with patients referred to the hospital already investigated in return for a maximum wait for consultation of four to five days, and immediate inpatient treatment in big hospitals only for the ill who cannot be at home, we would have a first-rate national service, attracting young doctors to both hospital and general practice.—I am, etc.,

H. C. H. BIRD.

Coxheath,
Nr. Maidstone, Kent.

Isolation of B.M.A. Divisions

SIR,—The alterations to the constitution of the B.M.A. are becoming increasingly open to question. Obviously they should have been debated at a full S.R.M.—possibly lasting two days—and with no other subjects on the agenda to confuse the issue, rather than having been dealt with in the piecemeal fashion that they were.

That the Council agreed to decrease its numbers, possibly as a sop to the Representative Body, whose size was drastically curtailed, is now appearing to be less than wise, for the huge consequent increase in