Parent and Child

SIR,—In 1964 the then Ministry of Health¹ issued a warning to the family doctor that "there is evidence that separation before the age of 5 may have a serious effect on the emotional growth of children and may form the basis of neurotic reactions in later life." A recent instance shows that one doctor at least has, properly, found it right to ignore this advice.

As reported in the *Guardian* on 16 January, a mother wished, with the support of her doctor, to place her infant in the care of the children's department as she felt unable to care for the child. Her plea was rejected and the child died. The reason for rejecting the mother's plea was the "likely effect of parting the child from her mother." The coroner commented "I can't see the point of leaving a child with its mother merely to let it die of starvation."

This tragic instance has arisen because of the fallacious belief that separation is synonymous with deprivation. Not a single investigation has shown that separation *per se* leads to mental ill health in children. As demonstrated by this instance, deprivation can occur without separation; indeed the great majority of deprived children live at home. In a small number of instances and in the right circumstances separation is essential to overcome deprivation and to give the child an opportunity for emotional growth. Separation need not lead to deprivation if the placement of the child is carefully selected by skilled, trained professional workers. May the now Department of Health urgently reconsider its policy and advice.—I am, etc.,

J. G. HOWELLS.

The Institute of Family Psychiatry,
Ipswich and East Suffolk Hospital,
Ipswich.

REFERENCE

- 1 *The part of the family doctor in the Mental Health Service*, London, H.M.S.O., 1964.

Impotence from Toxic Chemicals?

SIR,—The premise "having been unable to find any alternative cause" is hardly a satisfactory starting point for the conclusion that impotence "was probably due to the toxic effect of the chemicals . . ." and in the absence of supporting evidence—clinical, laboratory, or statistical—it can be misleading.

From time to time many drugs and chemicals are accused of causing various symptoms (including impotence) for which no organic basis is discovered. Dr. M. L. E. Espir and others (14 February, p. 423) are correct to report this "cluster of cases." The facts are important, but I submit that the evidence is insufficient for even a tentative conclusion or probability.—I am, etc.,

A. L. LEIGH SILVER.

Fleet, Hants.

Electric Bell Alarm for Enuretics

SIR,—With reference to the article by Dr. W. I. Forsythe and Dr. A. Redmond, (24 January, p. 211), may I suggest the time is ripe for a consensus of opinion on this matter and a clinical assessment of the op-

timum mode of dealing with enuresis in children.

It is very difficult to understand how it comes about that the electric alarm, a most effective treatment which was first reported in 1938,^{1,2} is still not established as orthodoxy 32 years later. Equally difficult to understand is the situation which sees this treatment wholly accepted in some areas with visible success, whereas in other areas the treatment is hardly used at all.—I am, etc.,

HAROLD PALMER.

Hill End Hospital,
St. Albans, Herts.

REFERENCES

- 1 Mowrer, O. H., *American Journal of Psychology*, 1938, **51**, 163.
- 2 Mowrer, O. H., and Mowrer, W. M., *American Journal of Orthopsychiatry*, 1938, **8**, 436.

Atrial Function in Man

SIR,—Your leading article entitled "Atrial Function in Man" (24 January, p. 189) does not include mention of the importance of the atria in the control of heart rhythm. The sino-atrial node controls the heart by its higher intrinsic rate than the atria and ventricles. Disorder of the sino-atrial node with reduction of excitation of the atria is frequently followed by atrial fibrillation or flutter. In such cases reversion of the atrial fibrillation to sinus rhythm produces a bradycardia with later atrial escape beats. The patient and doctor feel better when atrial fibrillation returns.

No convincing evidence of specific atrial conducting fibres has been found. As your leader writer said, the atria act as reservoirs, particularly in the presence of heart failure, and their distension by blood might well impair the function of any specialized conducting fibres in their walls, particularly between the two atria.—I am, etc.,

S. TALBOT.

General Hospital,
Nottingham.

March Haemoglobinuria

SIR,—I was most interested to read Dr. A. J. Spicer's report (17 January, p. 155), especially as an attempt was made to reproduce the clinical situation invoked by karate exercises as reported previously by myself.¹ Although haemoglobinuria in the present case only occurred following running on a hard road in thin soled shoes as is usually the case, there were, in my opinion, very significant changes in the measured levels of plasma haemoglobin and plasma haptoglobin following the karate exercises.

In the previously reported case,¹ the patient had been performing his karate exercises daily for over eighteen months, and the force of his blows was such that when he performed his exercises under supervision the separate impacts could be clearly felt and heard on several floors of a ferro-concrete building. It would be unlikely that a novice could achieve this force of impact without suffering at best, considerable pain, or at worst, fractures, particularly in view of the duration and nature of the exercises as

previously reported. In the light of the present findings, it would appear that haemoglobinuria is not an essential feature, and that in fact many devotees of karate may well be haemolysing steadily, as they practise their exercises.

I would support Dr. A. J. Spicer's conclusions that haemolysis is due to intravascular trauma to the circulating erythrocytes. This trauma-induced haemolysis can obviously occur in any part of the body which is subject to an intense recurrent force of short duration.—I am, etc.,

JONATHAN A. STREETON.

Alfred Hospital,
Melbourne, Australia.

REFERENCE

- 1 Streeton, J. A., *Lancet*, 1967, **2**, 191.

Galen and the Arteries

SIR,—I should like to offer some remarks in support of Dr. C. E. Newman's review of Margaret Tallmadge's new translation of Galen's works (6 December, p. 609). It occurs to me that some readers might be surprised at the suggestion that parts of Galen's dissertation on the contents of the arteries could remain clouded with ambiguity and mystery, in some cases until the nineteenth century. In particular, Galen's supposed assumption that the arteries contained air was based on a mistaken comparison with Erasistratus's theory that the arteries are usually filled with air, but that this escapes when an artery is damaged, causing a vacuum which in turn draws the blood from the veins and through the capillaries into the arteries until it is finally emitted as the last of the escaping air is ejected.

But the reason for the comparison and the mistaken view of Galen's theory is to be found in the Stoic doctrine of "pneuma," by which Galen's medical theory was strongly influenced. "Pneuma" was the technical term originally used by Chrysippus (281-208 B.C.) to denote the "active principle," which comprised the contents and qualitative elements of all matter, as opposed to the "passive principle," which was unqualified and dormant matter. The former imparted life and purpose to the universe, ensuring that it fulfils its preordained development, while the latter was the purely passive and characterless recipient of the former. "Pneuma" is most frequently represented as being composed of two elements, air and fire, which give the body in which they reside the perceptible qualities it possesses in accordance with the ratios in which they occur. On the other hand, however, it was also occasionally used as a synonym for air itself, which, together with fire, was but an element in the "active principle." Owing to this ambiguity of meaning by which "pneuma" was sometimes used as a Stoic technical term and sometimes as a synonym for "breath," the translators and commentators were often in some apparent doubt about its significance in any particular context.

Galen's interpreters were not immune to the philosophical implications of the word, and tended to assume that where he referred to "life-giving spirit" in the arteries he really meant "air." The teleology of Galen's outlook, each part of the body sub-

serving its special purpose, produced further resemblances with Stoicism to lend support to a false interpretation.—I am, etc.

J. C. DAVIES.

Department of Classics,
The University,
Sheffield, Yorks.

Laryngitis and Aerosols

SIR,—A 39-year-old man was referred with a six-month history of progressive hoarseness. Biopsies of the right aryepiglottic fold and right ventricle of the larynx performed at the referring hospital showed a chronically ulcerated, hyperplastic squamous mucosa only. The inflammatory cell infiltration was predominantly plasma cell in type. Indirect laryngoscopy showed an oedematous larynx with a papilliferous lesion of the right aryepiglottic fold with extension into the medial wall of the right pyriform fossa. The larynx was fully mobile; there was no ulceration and no palpable lymph nodes. A repeat direct examination with multiple biopsies was again negative for carcinoma; the histological appearances being similar to the previous biopsy.

He was an asthmatic of seven years' duration, on low-dose steroids for four years. He originally used a pressurized aerosol of isoprenaline sulphate 4 mg./ml., but had changed to the stronger suspension containing 20 mg./ml. for the past 18 months. His asthma had worsened progressively over nine months to the extent that when seen he had been off work for five months, able only to manage one flight of stairs, or 200 yards on the flat as a maximum exercise at any one time, and unable to sleep without being woken three or four times each night by asthma. Coincident with the deterioration of his asthma, there had been an increasing use of the aerosol, so that a 400-dose aerosol was lasting only one week, equivalent to more than 50 doses per day. Patients have been warned that it is dangerous to exceed the recommended dosage of one to three doses, and that it should not be necessary for more than eight treatments a day.¹

One week prior to his readmission for a further direct laryngoscopy and biopsy, the patient changed back to the low dose isoprenaline aerosol which he had first obtained 18 months previously. There was a dramatic improvement in his voice over seven days and a considerable early improvement in his general condition. Subsequent follow-up on three occasions over two months confirmed that the improvement was maintained. He was back at work full time after two weeks and by two months his larynx and voice were normal; he was sleeping uninterrupted and had been swimming for the first time for seven years. Whereas he had previously become dependent on his aerosol he was now actively refraining from its use as much as possible. He had been changed at one month to an adrenaline aerosol and was using it only once or twice a week.

A presumptive diagnosis of carcinoma of the larynx had been made in this patient. However, because this seemed a strangely situated and small lesion to cause the severe degree of hoarseness he had, treatment was delayed until a positive biopsy could be obtained. A dramatic improvement in his laryngeal condition followed a reduction in the concentration and the frequency of use

of a pressurized aerosol containing isoprenaline.

There have been several reports of the dangers of excessive use of any sympathomimetic amine in bronchial asthma,^{2,4} and statistical evidence of a relationship between overdosage and mortality.^{5,6} The evidence for isoprenaline being the cause of this man's hoarseness must remain circumstantial. However, on clinical grounds alone it would seem a reasonable assumption.

I am grateful to Mr. H. J. Shaw for permission to report this case.

—I am, etc.,

G. E. STOLLARD.

Royal Marsden Hospital,
London S.W.3.

REFERENCES

- 1 Committee on Safety of Drugs, *Adverse Reactions Series*, 1967, No. 5.
- 2 Keighley, J. F., *Annals of Internal Medicine*, 1966, **65**, 985.
- 3 Pickvance, W., *British Medical Journal*, 1967, **1**, 756.
- 4 Greenberg, M. J., and Pines, A., *British Medical Journal*, 1967, **1**, 563.
- 5 Speizer, F. E., Doll, R., and Heaf, P., *British Medical Journal*, 1968, **1**, 335.
- 6 Inman, W. H. W., and Adelstein, A. M., *Lancet*, 1969, **2**, 279.

Pathogenesis of Pre-eclampsia

SIR,—It is true that research into pre-eclampsia has been bedevilled by lack of clinical definition of the large group of patients with hypertension and proteinuria in late pregnancy. Therefore it is of great interest to read Dr. Helen P. McEwen's claim that the presence of alpha₂ macroglobulin (alpha₂M) in the urine in organic disease of the kidney distinguishes chronic renal disease from pre-eclampsia (10 January, p. 111).

If this is so, it might be assumed that this characteristic urinary protein pattern of chronic renal disease is the result of a different degree of glomerular permeability to proteins, which should be identifiable by differential protein clearances. Sixteen patients were studied in this hospital; nine with a clinical diagnosis of severe pre-eclampsia, four with proved nephrotic syndrome, and three with an undetermined cause of proteinuria. The protein clearance was invariably unselective—that is, IgG/transferrin = 20-30%—and afforded no help in the differential diagnosis of these patients.

From these preliminary results we think it probable that with the exception of minimal change glomerulonephritis all proteinuria occurring in pregnancy will be unselective, and that the detectable presence of alpha₂M in the urine will be determined solely by the quantity of proteinuria. We have, for example, found alpha₂M in the urine of a severe pre-eclamptic with a protein loss of 13 g. per day and an absence of detectable alpha₂M in a nephrotic with a proteinuria of 4 g. per day. Nevertheless, we do believe that the presence of alpha₂M in the urine may have some prognostic value, as Dr. McEwen has recently claimed,¹ but considering the similarity of the protein clearance it would appear to us that this index of severity could be deduced more easily by accurate measurement of the proteinuria.

It is ironic that, while Dr. McEwen is rightly stressing the importance of detailed consideration of proteinuria in pregnancy, most obstetricians will entrust the

measurement of urinary protein to a nurse or medical student using Esbach's albuminometer. We have compared the results obtained from Esbach's test performed in the manner described by Varley² with the more accurate biuret method in 145 twenty-four-hour urine collections from 11 patients. Eight patients had severe pre-eclampsia and three had pregnancies complicated by the nephrotic syndrome. The proteinuria in all patients was unselective. The results are shown in the Figure below.

These data confirm the inaccuracy of the Esbach's technique at all levels of proteinuria. The use of this inaccurate method is particularly inappropriate in pre-eclampsia because if the hypertension has been successfully controlled by drugs the precise urinary protein loss may be the most important index of the severity of the pre-eclampsia available, and if it is true that appearance of alpha₂M macroglobulin in the urine in pre-eclampsia is a harbinger of fetal death¹ accurate measurement of urinary protein in pre-eclampsia must have a similar predictive value.—We are, etc.,

J. W. W. STUDD.

J. D. BLAINEY.

Birmingham Maternity Hospital,
Birmingham 15.

REFERENCES

- 1 McEwen, H. P., *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 1969, **76**, 809.
- 2 Varley, H., *Practical Clinical Biochemistry*, 4th edition, p. 143. 1967. London, Heinemann.

Phage Treatment of Severe Burns

SIR,—Many methods of treating infected severe burns have been advocated, including immunization against the common infecting organisms, chiefly hospital-resistant staphylococci and *Pseudomonas pyocyanea*, which inhibit healing and are antibiotic resistant. In the first world war, I used this method of treating infected septic