Correspondence

Correspondents are asked to be brief.

Mild Ulcerative Colitis I. E. Gillespie, F.R.C.S.	689	A Student's View R. D. Walker 692	Immunoglobulin Levels in C.S.F. B. M. Greenwood, M.R.C.P.	694
Bundle Branch Block B. B. Scott, M.R.C.S.; J. W. Todd, F.R.C.P.	689	Diarrhoea and Light Therapy in Neonates A. J. O'Donnell, D.C.H.; G. R. Griffith, F.R.C.P.ED., and others 692	Folate Deficiency, Megaloblastic Anaemia, and Endocarditis K. G. A. Clark, M.R.C.PATH.	694
Impotence in Farm Workers A. W. Pack, M.R.C.P.	690	Self-interest and the Common Good V. B. Purvis, D.O.	Holiday Typhoid and T.A.B. A. H. Griffith, M.D.	694
Post-thyroidectomy Recurrent Laryngeal Nerve Palsy Sir Victor Negus, F.R.C.S.; H. J. Shaw, F.R.C.S.	690	Vagotomy for Peptic Ulcer K. Jarvis, M.B., and others 692	Dr. Raymond Greene R. Greene, F.R.C.P.	694
Sterilization of Currettes R. J. Fallon, M.D.	690	Mass Radiography F. J. H. Walters, M.D.; J. G. Bancroft, M.B.	Schism in the Profession Sir George McRobert, F.R.C.P.	694
Unheard Voices Helen M. F. Firth, F.F.A.R.C.S., and others 690	690	Glossopharyngeal Neuralgia J. Penman, M.R.C.P.	Special Representative Meeting B. W. Hill, M.B.	695
Coagulation Failure in Pregnancy L. D. Courtney, M.R.C.O.G.	691	Phenformin and Stanazolol in Blood Fibrinolytic Activity G. R. Fearnley, F.R.C.P.	Green Paper and Postgraduate Medical Education D. C. Bowie, F.R.C.S.ED.	695
Postoperative Fibrinolysis E. T. O'Brien, L.R.C.P.I.	691	Silver Nitrate "Foreign Body" N. Hira, F.R.C.S., and F. W. Lewis, D.M.R.D.	Subconsultant Grade M. A. Gilbert, M.B.	695
Diazepam and Muscle Relaxants S. A. Feldman, F.F.A.R.C.S., and B. E. Crawley, F.F.A.R.C.S.	691	Post-vagotomy Diarrhoea? D. G. Weir, F.R.C.P.I.	Hospital Career Structure J. D. Hill, M.B.	695
Corticosteroids and Pituitary-Adrenal Function J. R. Daly, M.R.C.PATH.	691		Future Trends in Medical Staffing I. W. Young, F.R.C.S.	695

Mild Ulcerative Colitis

SIR,—The questions raised by Mr. A. K. Maitra in his letter (14 February, p. 431) on the management of mild ulcerative colitis are indeed difficult ones, and it is understandable that confusion may appear to exist. Since there are no final answers to some aspects in advising surgical as opposed to medical treatment, it is possible to interpret expert opinion in slightly different ways.

It does seem clearly established that patients with ulcerative colitis have a greater risk of the development of colonic or rectal carcinoma than others. However, it also seems clearly demonstrated that those with total involvement of the colon and/or severe relapses have a substantially greater risk of carcinoma than do those with mild distal non-ulcerative inflammatory changes. It would seem that all authors are agreed on this matter. The point made by Professor B. N. Brooke¹ was not that the greater proportion of patients developing cancer in the colon had mild colitis, but simply that this group should not be ignored when considering the possibility of the development of malignant change. The colonic cancer risk in those with mild distal colitis, though present, would seem to be much smaller than in those with the more extensive and/or more severe disease.

Since the usual operation for ulcerative colitis is total procto-colectomy with permanent ileostomy, there is naturally hesitation in advising this radical procedure unless the risk of development of carcinoma is substantially greater than the risks attendant on this operation. No absolutely accurate information on this particular point is available, although the studies of Watts and colleagues² suggest that for those with extensive colonic involvement a stage is

reached about three years after onset of the disease when the risk of tumour becomes greater than that of total colectomy. The results of the investigations of Dr. Morson and his colleagues³ into the possibility of selecting these patients with even mild colitis who might be more at risk of malignant change than the others, appears strongly convincing. All specimens removed for established carcinoma in ulcerative colitis patients had similar uniform histological appearances, and there was a high incidence of carcinoma and active mucosal changes in those who underwent colectomy as a result of the detection of the biopsy changes.

In summary, it would seem that the only absolute guarantee to any patient with any degree of ulcerative colitis is a total procto-colectomy. However, it seems genuinely doubtful whether the risks of such a radical procedure, small though they are, are significantly less than the risk of malignant disease in those who remain symptomfree for years at a time, and who have only mild distal colitis. The persistence of a very real fear of the development of malignant disease may in itself be every justification for coming down on the side of operation.—I am, etc.,

IAIN E. GILLESPIE.

Department of Surgery,
Western Infirmary,
Glasgow.

REFERENCES

- 1 Brooke, B. N., *Journal of the Royal College of Surgeons of Edinburgh*, 1969, 14, 274.
- 2 Watts, J. McK., de Dombal, F. T., Watkinson, G., and Goligher, J. C. *British Medical Journal*, 1966, 1, 1447.
- 3 Morson, B. C., and Pang, L. S. C., *Gut*, 1967, 8, 423.

Bundle Branch Block

SIR,—I read with interest your leading article "Bundle Branch Block" (21 February, p. 450). You state that in an otherwise healthy non-patient population the incidence of bundle branch block is low. Although you admit that the incidence is higher in the elderly, the high incidence in the following report may surprise you.

Electrocardiograms were done on 161 consecutive admissions to the geriatric unit at this hospital under the care of Dr. H. Droller from November 1969 to January 1970. They were divided into two groups. Group I (78 patients) included those with any of the following criteria: taking diuretics, digitalis, or hypotensive drugs before or after admission; symptoms or signs of heart failure; a history of angina or myocardial infarction. Group II consisted of those without any of these criteria (83 patients; 53 women, 30 men). The ages ranged from 65 to 94 years.

The reasons for admission to hospital of those in Group II were not directly related to disease of the cardiovascular system except in some patients who had recently had a cerebral ischaemic episode and others who had senile dementia. Many of these admissions were on social grounds. Thus this group perhaps approximates fairly closely with the general population aged over 65 years so far as cardiac status is concerned. Of this Group II, bundle branch block was found in 9 (17%) of the women (4 left and 5 right) and in 2 (7%) of the men (both right).

Furthermore, no important underlying organic basis for their conduction disturbance was found, contrary to your prediction. None were hypertensive, but two of the women were in atrial fibrillation (one with left bundle branch block, the other

right). Of the rest the E.C.G. was otherwise normal. It is intended to conduct a follow-up on these patients.—I am, etc.,

BRIAN B. SCOTT.

St. James's Hospital,
Leeds, Yorks.

SIR,—Your leading article (21 February, p. 450) states: "... there is a low incidence of right or left B.B.B. among an otherwise healthy, non-patient population. ... It is this situation which faces the practitioner with a dilemma." I can see no dilemma. The simple means of dealing with this situation is to withhold the information from the victim that he has bundle branch block. Then "the risk of undue cardiac invalidism" cannot arise.

An even better solution to this problem is to refrain from doing E.C.G.s in "an otherwise healthy non-patient population." Whatever the justification for some kinds of screening in certain people, I know of no evidence that E.C.G.s done on the "healthy non-patient population" benefits anyone.—I am, etc.,

JOHN W. TODD.

Farnham, Surrey.

Impotence in Farm Workers

SIR,—Dr. M. L. E. Espir and his colleagues found it difficult to explain the mechanism(s) by which a heterogeneous group of insecticides might have caused impotence in four agricultural workers (14 February p. 423). One possible explanation is that since the insecticides used included a chlorinated hydrocarbon, dieldrin, the metabolism of testosterone was increased and this led to a relative deficiency of this hormone.

It is well established that chlorinated hydrocarbon insecticides are potent inducers of liver microsomal enzymes in animals.¹ While interest has been focused chiefly on the ability of these enzymes to oxidise drugs and toxic chemicals to more polar metabolites which can be excreted, they are also involved in the hydroxylation of steroid hormones. Administration of chlordane or D.D.T. to rats for ten days approximately doubled the rate of hydroxylation of testosterone by rat liver.² Dieldrin, which is closely related structurally to chlordane, and is known to reduce the sleeping time of mice following fixed doses of hexobarbitone,³ might well have a similar action. By contrast, chlorthion, an organophosphate cholinesterase inhibitor, when given to rats for ten days, reduced the hydroxylation of testosterone by rat liver.²

The relevance of these experiments to man with respect to the hydroxylation of endogenous testosterone is uncertain, but it seems likely that certain liver hydroxylation reactions are increased by exposure to chlorinated hydrocarbon compounds.¹ This conclusion is supported by the observation that the half life of antipyrine was reduced to approximately 50% in a group of Swedish workers exposed to D.D.T., chlordane, and lindane, compared with the half life measured in 33 non-exposed control subjects.⁴ In addition, it is possible that insecticides of this type might interfere with the synthesis of various steroid hormones, an

action which accounts at least in part for the reduction in corticosteroids achieved therapeutically by the related compound *o,p'* D.D.D.⁵

Chlorinated hydrocarbon compounds are eliminated slowly from the body, and this would account for the duration of symptoms. The comments of informed clinical endocrinologists would be helpful.—I am, etc.,

ANTHONY W. PECK,

Clinical Research Division,
Wellcome Foundation Ltd.

Beckenham, Kent.

REFERENCES

- 1 Conney, A. H. (1967). *Pharmacological Reviews*, **19**, No. 3, 317.
- 2 Welch, R. M., Levin W., Conney, A. H. (1967), *J. Pharm. and Exp. Ther.* **155**, No. 1, 167.
- 3 Fouts, J. R. (1963), *Ann. New York Acad. Sci.* **104**, 875-880.
- 4 Kolmodin, B., Azarnoff, D. L., Sjöqvist F. (1969), *Clin. Pharm. & Therap.* **10**, No. 5, 638.
- 5 Calabresi, P., and Welch, A. D. (1965) in *Pharmacological Basis of Therapeutics*. Ed. Goodman, L. S. and Gilman, A. Macmillan Co., New York.

Post-thyroidectomy Recurrent Laryngeal Nerve Palsy

SIR,—I have read with considerable interest the very complete and exhaustive article by Mr. Victor Riddell on "Thyroidectomy: Prevention of Bilateral Recurrent Nerve Palsy."¹ I am impelled to add the testimony of a laryngologist of many years' standing to draw attention to the views expressed, which may not reach those members of the medical profession who do not receive the *British Journal of Surgery*. Exposure of the nerve on the first side is essential, together with testing of its integrity by electrical stimulation as is Mr. Riddell's practice; this is carried out before operation on the other side is contemplated.

I write to support Mr. Riddell in his decision to identify the nerve and to assure himself of its integrity before proceeding further.—I am, etc.,

VICTOR NEGUS.

Haslemere, Surrey.

REFERENCE

- 1 Riddell, V., *British Journal of Surgery*, 1970, **57**, 1.

SIR,—From a lifetime of wide experience of thyroid surgery, Mr. V. Riddell¹ gives emphatic attention to the identification and to an estimate of normal function of the recurrent laryngeal nerve on at least one side when carrying out the operation of total or subtotal thyroidectomy.

His main principle in thyroidectomy is an attempt to establish normal function of the nerve on the side first dissected before proceeding to the opposite side. Further, if there be real doubt regarding the nerve on the first side, dissection of the second side must be delayed until the function of the nerve and vocal cord on the first side is known. He emphasizes the tragedy of a bilateral recurrent nerve palsy after this type of surgery and the difficulties of really effective laryngeal surgery to restore normal respiration and voice. I am also glad to see that he lays stress on pre-operative laryngoscopy in addition to inspection of the larynx during surgery. Postoperative laryngoscopy is, I feel, also desirable.

As a laryngologist with an interest in the field of head and neck tumour surgery I would heartily endorse and support these principles. My only reservation would be that where total thyroidectomy is required for malignant disease, a more radical approach may of necessity put one or even both nerves at risk. In such cases I believe that results of functional tests of the nerves must not be allowed to delay a complete resection.—I am, etc.,

H. J. SHAW.

Royal Marsden Hospital,
London S.W.3.

REFERENCE

- 1 Riddell, V., *British Journal of Surgery*, 1970, **57**, 1.

Sterilization of Curettes

SIR,—Dr. J. M. McGarry (3 January, p. 49) states that the curettes were sterilized as recommended by the manufacturers—that is, "two minutes' immersion in a ½% solution of chlorhexidine in 70% alcohol prior to use". Immersion in alcoholic chlorhexidine is not a sterilizing procedure. This treatment will kill vegetative organisms, but will not kill bacterial spores. This point is most important when it is remembered that curettes may easily be contaminated with spore-bearing organisms during use. With inadequate sterilization techniques viable spores could remain on a curette and could be transferred during subsequent use into an area which may contain necrotic tissue and would be an ideal site for the growth of clostridia. Obviously such an instrument should either be made of materials which are heat-stable so that it can be sterilized by heat, or alternatively arrangements could be made to sterilize the instrument either by low temperature steam and formaldehyde or by ethylene oxide. In the present circumstances the use of a 2% buffered solution of glutaraldehyde could also be considered.

There is an increasing tendency for new apparatus to be made of materials which are not suitable for sterilization by reliable methods such as dry or moist heat. This is a most unfortunate trend and one which should be resisted, as far as possible, by those responsible for the purchase of new equipment.—I am, etc.

R. J. FALLON.

Department of Pathology,
Ruchill Hospital,
Glasgow N.W.

Unheard Voices

SIR,—Reading your leading article "Unheard Voices" (7 February, p. 316) we concluded that the *B.M.J.* had finally come to grips with a very real problem. Just how much we were misled was obvious when we studied "The Surgeon" (p. 358). In comparison with many, Mr. Stubton should know that one weekend on in four is paradise. What about those hospitals where there are only two consultants in the major specialties? Alternate weeks on call and, only if we are lucky, alternate weekends off. Holidays, sickness, and unexpected absence of any kind lead to continuous duty for indefinite periods.

We regard this as consultant exploitation, and if the *B.M.A.* genuinely intends to