MEDICAL PRACTICE

Gynaecology in General Practice

Fecundity

BARBARA LAW

British Medical Journal, 1971, 1, 443-445

The reproductive phase of a woman's life may extend for forty years or more. During this time the control of her fertility will influence her physical and mental health, and on this will directly depend the welfare of her husband and children.

Family Planning

Nowadays, most women, regardless of their age, parity, or marital status are prepared to seek family planning advice without prompting. Others, however, even when urgently in need of help are reticent to ask. The family doctor, with his knowledge of the home background, is the obvious person to provide the necessary advice, either by accepting full responsibility himself or by putting his patients in touch with one of the available services.

Practice in this field is very different from other branches of medicine. A healthy woman, free of symptoms of disease, may not be willing to accept dogmatic prescription of a contraceptive method, and always the selection of a method will be made more complicated by the need to consider the husband or partner. The use and effectiveness of any birth-control measure depends on its acceptability to the couple and their motivation to use it correctly and consistently. The ultimate choice therefore should only be reached after careful assessment of their needs and wishes, with reference to age, health, parity, race, religion, and social circumstances. Such consultation may prove very time-consuming, especially if the original choice is found to be medically contraindicated.

Supervision and reassessment of the suitability of the selected method at regular intervals are important; early

reassurance may be necessary, and after prolonged use warning against complacency and careless use may be indicated if unintentional failure is to be avoided.

Methods of Control

ORAL CONTRACEPTION

By the inhibition of ovulation almost total protection against pregnancy is possible provided "the pill" is prescribed and taken correctly.

Acceptability

The pill is highly acceptable, relatively cheap, capable of relieving dysmenorrhoea and menorrhagia, and achieving regular menstrual cycles and is consequently the most popular contemporary method among women. It is, however, not suitable for all. It increases the risk of thromboembolism and is responsible for numerous systemic effects, whose significance is not yet fully understood. The woman's present and past health and her obstetric and family history must all be scrutinized before deciding that she is suitable for oral contraception.

Contraindications

It is the method of choice for those demanding or needing maximum protection from pregnancy. Some of the medical conditions making this necessary, however, may themselves be adversely affected by the components of the pill—for example, diabetes, hypertension, and cardiac and psychiatric disorders. In such cases the possible consequences have to be weighed against the slight risk of pregnancy if less acceptable contraceptive measures are used.

In the presence of hepatic insufficiency, oestrogen-depen-

dent tumour, or a history of thromboembolism the pill is absolutely contraindicated. The prescription should also be avoided if other recognized predisposing factors to thrombosis are present such as obesity, hypercholesterolaemia, or excessive smoking. The risk is also increased where there is a strong family history of thrombosis, in the older age groups, and if the blood group is other than O. Wherever possible the pill should be discontinued at least six weeks before elective surgery.

Other Considerations

Pregnancy is possible only if pills are omitted or not absorbed, or if additional precautions are not taken during the first 14 tablet days in the first cycle. Similar care is necessary when changing to a preparation containing a lower dose of either component. If two successive pills are omitted precautions should be taken for the rest of the cycle.

If breakthrough bleeding occurs it is still necessary to adhere strictly to the 21/7 or 22/6 tablet-taking routine. When women are given the "everyday" (ED) system they must be warned that some tablets are placebos and that they must be taken in the correct order and not preserved as an economy measure.

It is not always possible to select initially the preparation which will ultimately prove the most acceptable. The lowest potency pill which will give good cycle control is obviously the best choice, and some women will be suited by any, and others by none. If numerous complaints are made physiological hypersensitivity or excessive anxiety should be suspected. The cost may be important, but the smallest dose pills are not necessarily the cheapest. Careful follow-up is important, and the appearance of excessive weight gain, rise in blood pressure, headaches, depression, and irregular bleeding indicate the need for investigation, change of preparation, or even abandonment of the method. Provided there are no symptoms or signs of systemic effects there is no time limit to prescription of the pill, though it should be stopped when the menopause is reached.

Subsequent fertility is usually unaffected, and there is no increased liability to miscarriage or multiple pregnancy. Prolonged amenorrhoea after oral contraception has occurred, but most cases have responded to clomiphene. The woman who has not had a pregnancy, especially if she has scanty irregular menstruation or had a late menarche, should be warned of this possibility and offered alternative methods.

The pill has revolutionized the control of fecundity but its long-term influences are still not known.

INTRAUTERINE DEVICE

The loop or coil is the method of choice for the unmotivated patient, giving the maximum protection with the minimum of effort (Fig. 1). When inserted by skilled and experienced doctors the failure rate is about 3 per 100 women-years, provided the device remains in situ. Extrusion, often unnoticed, occurs in about 10% of cases and is commonest in the early months.

FIG. 1

Insertion

Insertion is most easily carried out during a period (thereby excluding the presence of a pregnancy) and is usually painless in a multiparous woman. Only in special circumstances—for example, educationally subnormal girls—is it justifiable for a woman who has not had a full-term delivery, and an anaesthetic may then be required for insertion.

Contraindications

Caesarean section is not a contraindication to using the intrauterine device, but insertion is better delayed for three months. Congenital abnormalities of the uterus, cicatrization of the cervix, and submucous fibroids may prohibit the method. Pelvic infection in the preceding six months is an absolute contraindication.

Increased and prolonged menstrual loss is the commonest complication and is most frequent during the first few months. Anaemia must be watched for. Pain is unusual but may be a warning of imminent extrusion of the device. There is no increased incidence of pelvic infection in users of the intrauterine device, but if it occurs it is usually the result of reactivation of existing sepsis. Antibiotic treatment is usually effective, but in some cases removal of the device proves necessary. Perforation is rare (1 in 2,500) and probably occurs at the time of insertion. The device can be located by an x-ray film and does not cause harm unless a "closed" device is used, when intestinal obstruction is a remote possibility.

Probably 70% of women fitted with a device are happy with the method, and some are willing to tolerate gross degrees of menorrhagia and resist its removal as long as they can. Regular cervical smear testing is advisable, though there is no evidence of malignant effect. In the absence of symptoms or signs there is no time limit to leaving the device in situ. Unfortunately some women who might be suitable for this method refuse it, because of its poor reputation of efficiency or for fear of having "something inside." Much research is in progress to improve the shape and composition of devices to reduce the expulsion and pregnancy rate.

FEMALE BARRIER METHOD

This old-fashioned but completely harmless method has considerable value at present, but it requires the strongest and continued motivation for successful use, and is only suitable for the well-adjusted woman. The acceptability of this method depends to a great extent on the attitude of the

FIG. 2

doctor who offers it in the first place and on the way he presents it.

Diaphragms and caps require careful fitting, so that they present as complete a barrier as possible and yet cause no discomfort to either partner (Fig. 2). Regular checking of size is important, and additional use of chemical spermicide is imperative. Precise teaching of the technique is absolutely essential, so that complete understanding allows the maximum acceptability possible. The woman has to learn to use it in such a way that the natural spontaneity of the relationship is not spoiled.

Women of low intelligence can be taught the method, but overcrowding and lack of toilet facilities can make it too difficult to use. It is never 100% efficient, but having no systemic or local damaging effect and possibly even being protective to the cervix it can be used by the right type of woman with much greater reliability than is usually realized.

VAGINAL CHEMICAL BARRIERS

The large number of spermicidal products on the market is remarkable considering their modest efficiency when used without mechanical barriers. They are available as paste, cream or jelly, pessaries, foaming tablets, or aerosol foams, with different active principles and perfumes. The aerosol foams are the most acceptable and efficient and also the most expensive. Insertion must occur within 30 minutes of coitus and the failure rate is too great for those whose family is already complete, or where pregnancy is strongly contraindicated.

MALE BARRIER METHOD

This is a highly efficient method if used on every occasion together with a vaginal spermicide. Preliminary penile contact with the perineum must be avoided and great care exercised after intercourse to remove the condom from the vagina at the same time as the penis. As it is necessarily interrupting, some men find it difficult to accept and some women actively dislike it. Many men use it conscientiously for a while then tire of it, "take a chance," and pregnancy is the result. Being widely available and requiring no medical consultation, it is valuable as a temporary measure, and where the husband prefers to take responsibility or the wife has a dislike of semen it can be used happily and successfully as a permanent method.

RHYTHM METHOD

The practice of periodic abstinence, confining coitus to the days of the cycle following ovulation, is restricting and acceptable only to those couples whose strong religious convictions provide the necessary self-discipline and motivation. Accurate pinpointing of ovulation is difficult and can be judged only by daily recording of basal body temperatures—

but the method can be used with considerable success by conscientious women who have very regular cycles.

COITUS INTERRUPTUS

Coitus interruptus is widely practised but never recommended, and "withdrawal" or "being careful" is frequently reverted to after more reliable methods have been tried. It requires no forethought, purchasing cost, or storage, is always available, and allows a couple to preserve absolute privacy about their sexual activities—a very important consideration for those in the lower social groups. Possibly some psychosomatic conditions may result from the strain and anxiety experienced by one or other partner, but the method is probably much more successful in preventing pregnancy than is usually acknowledged.

STERILIZATION

Tubal ligation and vasectomy are the most effective contraceptive measures available. Both procedures must be considered irreversible and never carried out without extremely careful assessment of both husband and wife. Their age and health, the stability of the marriage, and the condition and number of existing children must all be considered.

This final step of sterilization is frequently requested nowadays and is becoming easier to obtain. When the woman has a medical condition which contraindicates pregnancy, or threatens to need a hysterectomy in the future, then the woman herself should be sterilized. Vasectomy is relatively simple and admission to hospital is not normally necessary, but sperm do not disappear from the semen for several months after the operation. Reassurance that hormone alterations do not follow these operations is often necessary.

Comments

Fecundity has been carefully protected by nature, and the efforts of man to control it frequently fail. Apparently the fundamental urge to reproduce is so powerful that some couples prove incapable of using any method of family planning successfully. Others manage to avoid knowledge of contraception and yet bitterly regret each pregnancy. Some women are particularly "good at having babies" and after temporary success with a method fail "accidentally on purpose."

Abortion, though wasteful and distasteful to many, often provides the solution to the problem. The difficulty of deciding to abort a woman whose emotions are strong and time is short inevitably means that sometimes the operation will be regretted later.

The perfect method of control of fecundity still eludes us despite active research, and so the practitioners must help those in need to use available methods as happily and successfully as possible.