Surgeon in Nepal

SIR,—Any of us who begrudge paying so much in taxation should re-read the Personal View on Nepal (13 March, p. 606). In 1960 the British Government established a British Military Hospital in Dharan, a town a day's trek south of the little village of Dankhutta, where the British-Nepal Medical Trust has set up a tuberculosis clinic. For over ten years now, the hospital and patients have fought a courageous battle against disaster and disease.¹ The Gurkhas are indeed truly grateful to the British taxpayer.

It was Chainpur, where Mark Howarth delivered the drugs, that a Himalayan black bear once held the whole village in terror, and where the local policeman had to be carried for five days through the mountains to Dharan, his face having been destroyed by the bear. Our biggest problems were the witch doctors and local folklore, for only after both had been tried unsuccessfully would the patients come to the hospital. Mark Howarth might be relieved to know that there is a certain black stone found only in the bed of the Arun, which, when rubbed on an open wound, will result in perfect union within 24 hours, or so my Nepalese friends assured me. The alternative is the peacock, whose flesh incidentally is delicious, being similar to pheasant. The feathers must first be burnt and then rubbed into the bleeding wound where they act as an excellent haemostatic agent.

Illegal poaching resulted in the admission of many patients like Dalbahadur (incidentally *bahadur* means warrior). One of the cruellest traps are explosives wrapped in a sweet bait, which are placed on the runs of the wild boar. The animal swallows the bait with the ghastly consequence. Sadly children have also sampled this sweet and treacherous killer.

The greatest good anyone can do for the health of the Nepalese is prophylactic medicine. The B.C.G. is the most important as there must be many hundreds of thousands of tuberculosis victims among the 11 million population of Nepal. The whole family sleep in the same room, the only room of the house. There are no windows, not only because of the intense cold in winter, but because it is imperative to retain the wood smoke from the central fire in the room to protect the timbers from being destroyed by insects. Without any ventilation, these buildings are an ideal media for "galloping" consumption. The saddest name the Nepalese have for tuberculosis is *Raj Banshi Rog*, royal family disease, or King's disease, as only a King, or at least a very rich person, can afford to be treated in Nepal. The rest have to resort to the witch doctors and their cure for tuberculosis—the meat of the jackal.—I am, etc.,

PETER PITT

Chase Farm Hospital,
Enfield, Middx

¹ Pitt, P., *Surgeon in Nepal*, London, John Murray, 1970.

Keeping on the "Medical Register"

SIR,—Until May 1970, the name of a registered doctor remained on the British Register till death unless the doctor's name was removed at his/her request, or because he/she failed to reply to a G.M.C. request for the current address, or for disciplinary reasons. Broadly speaking, to have one's

name removed from the Register implied serious professional misconduct, carelessness, or eccentricity.

I am writing to draw attention to the new situation which has arisen since May 1970, when the G.M.C. began to charge a fee to keep a doctor's name on the Register. The names of some doctors are now being removed from the Register because they have decided not to pay the fee. It is necessary for these persons to be fully aware of the consequences of being unregistered. Some have been qualified for several years and are widely known to be medically qualified. In spite of this, if their names do not appear in the Register, they would be unwise to describe themselves as doctor even if they hold a doctorate, as this might be interpreted as contravening section 31 of the Medical Act 1956 ("Penalty for pretending to be registered.") For instance, J.P.'s and others who may hold office, partly or wholly, by virtue of being thought to be doctors might find the absence of the title "doctor" to be embarrassing and disadvantageous.

Doctors not in regular medical work may be wondering about their position. They should appreciate that anyone doing occasional locums or sessions or other forms of intermittent medical work or regular medical work though small in amount must be registered. Such a doctor, to qualify for assistance from his defence society, must have been an ordinary paid-up member of the society at the time of the incident out of which the complaint, threatened claim, or litigation arose. Defence societies only offer a retired life membership without subscription to those who are genuinely retired from active practice, even if only temporarily. Retired life membership affords cover only for unpredictable emergency situations where the retired person acts, or is called upon to act, as a doctor.

Where an unregistered person is found to have practised medicine for gain as though he were registered, the G.M.C. when he applies for the restoration of his name to the Register will demand back-payment of the annual retention fee for each year or part of a year during which he so practised, in addition to the fee for restoration. Restoration to the Register does not confer retrospective registration, and does not confer retrospective rights to cover as an ordinary member of a defence society for any period when the person concerned was a retired life member, but if ordinary membership had been maintained by payment of subscriptions to the society during the relevant period the defence society would doubtless consider, on its merits, any application for assistance.

As is more generally known, an unregistered person may not sign certificates of death or incapacity or prescribe certain drugs. Nor should he/she sign any other documents, witness signatures, testify, or do anything else in the capacity of a registered medical practitioner. As for activities such as undertaking medical examinations, even on a voluntary basis, for such organizations as boy scouts, or conducting first aid classes, or anything else which might imply that the individual concerned was a registered medical practitioner, he would be wise to clarify his status for the information of the organizers of the body for whom he was undertaking the task. Unregistered practitioners might be required to serve on a

jury. They would have no electoral rights in a G.M.C. election.

Mailing lists for pharmaceutical firms, postgraduate authorities, Government departments, etc., are compiled from the Medical Register. Moreover, no unregistered person's name is knowingly included in the Medical Directory. Unregistered persons are, therefore, unlikely to receive informative literature, notices of medical meetings, postgraduate courses, etc. Their names will disappear from the medical scene.

Finally, it is possible that those of us who become unregistered in this way—for example, non-payment of the G.M.C. annual retention fee, and who therefore discontinue the use of the prefix "doctor" may well in ignorance be assumed by members of the public to have had their names erased from the Medical Register for disciplinary reasons. Before removing their names from the Register doctors would be well advised to study the disabilities which may follow such action.—I am, etc.,

JEAN LAWRIE
Honorary Secretary,
Medical Women's Federation

London W.C.1

Confidential Information

SIR,—The G.M.C.'s decision in the case of Dr. Robert Browne (*Supplement*, 20 March, p. 79) appears to draw a distinction between information which a doctor obtains directly from his patient and information which he receives unsolicited from other sources. Doctors apparently need no longer regard as confidential any information obtained from organizations such as the Family Planning Association or the Institute of Directors' Medical Centre as well as knowledge about patients which comes from such sources as hospital departments (of venereal disease, for example), solicitors, mental welfare officers, health visitors, and other members of the community health care team. The G.M.C.'s decision strikes at the very root of the doctor-patient relationship and introduces a very dangerous new ethical principle. We hope that the Association's Ethical Committee will make an immediate clear pronouncement deploring the G.M.C.'s decision and supporting the maintenance of the principle of confidentiality between patient and doctor.—We are, etc.,

JOHN WOODALL
O. WOOTON
E. TUCKMAN
J. D. PAULETT
YVONNE GRIFFITHS
M. E. ROBERTS
J. PEMBERTON (District Midwife)
D. C. WRIGHT (Mental Welfare Officer)
M. R. WHALEY (Health Visitor)
G. GREEN (District Nurse)

St. Paul's Cray,
Kent

Keratoderma Blennorrhagica in Reiter's Disease

SIR,—The case of keratoderma blennorrhagica in a woman with Reiter's disease (19 December, p. 723) illustrates the difficulty that may arise in differentiating this condition from arthropathic psoriasis.

The patient subsequently came under my care on 15 July 1970, three months after discharge from the Westminster Hospital. Prednisone had been discontinued two weeks previously, but relapse had occurred several

weeks before this. She was ill, with pyrexia of 101°F (38.5°C) and arthritis involving the hands, wrists, feet, the left knee, right elbow, and the hips. There were widespread skin lesions. Hyperkeratotic and pustular rash covered the toes, with marked nail involvement. Similar lesions were on the heels. There were psoriasiform patches on the palms and there were also nail changes on some fingers, the affected nails occurring where there was maximal joint involvement of the fingers. Discrete psoriasiform patches were present in the scalp and on the trunk and limbs. Subsequently, she developed guttate psoriatic lesions on the trunk and rupest psoriasiform plaques on the limbs. There was no involvement of the urethra, cervix, conjunctivae, or buccal mucous membrane.

Investigations showed E.S.R. (Westergren) 100-130 mm/1 hr., W.B.C. 9,700, (polymorphs 83%, lymphocytes 13%). Blood cultures, R.A. test, and L.E. tests, were negative. X-ray of os calcis and iliosacral joints did not show abnormality. Over a long period, her general condition gradually improved. A.C.T.H. injections appear to have had a good effect. She now has minimal joint involvement and minimal skin lesions.

Without knowledge of the transient urethritis, cervicitis, and buccal lesions early in the disease, I would have accepted this as a case of arthropathic psoriasis.—I am, etc.,

RODERICK HOWELL

Swansea, Glam.

Restless Legs and Latent Diabetes

SIR,—The leading article on "restless legs" (26 December, p. 758) and two original articles in the same issue need further comment with regard to the strikingly high incidence of this syndrome in cases of diabetes mellitus. Notable are the cases in which restless legs preceded the onset of diabetes. The actual number of diabetics who suffer from restless legs might be higher if one looks for the possibility of latent diabetes mellitus. In their interesting paper, Dr. N. K. Banerji and L. J. Hurwitz (p. 774) do not mention whether a glucose tolerance test was performed on the non-diabetics in their series. On the other hand, Dr. J. D. Spillane (p. 796) does not report his observations with regard to this metabolic abnormality; he states only that the neurological examinations were negative. However, as in cases of diabetic nephropathy or retinopathy, there does not necessarily have to be an intimate correlation between the restless legs syndrome and diabetes.

I have seen two patients with rather severe "restless legs," in whom a diabetic glucose tolerance test curve was found. One patient also had a moderate hypercalciuria and a mildly elevated urinary excretion of acid mucopolysaccharides.¹ All laboratory tests on the second patient were normal. The common abnormal physical findings in both was a reduction of vibration sense in the legs. This was regarded as a possible early sign of diabetic neuropathy. Both patients were treated symptomatically with a low carbohydrate diet, thiamine (200 mg/day), and a sedative (dichlorphenazone). Both showed a marked improvement. The first patient, who was followed for over a year, was able to decrease the amounts of both drugs gradually, without any significant reappearance of the previous complaints.

The possibility of restless legs heralding a (latent) diabetes mellitus and being an early expression of a diabetic neuropathy should be kept in mind.—I am, etc.,

I. MACHTEY

Hasharon Hospital,
Petah-Tiqva, Israel

¹ Machtey, I., *Harefuah*, 1969, 77, 468.

Uneasy Relations

SIR,—Dr. A. Paton's charmingly written Personal View (6 March, p. 554) reveals an attitude of mind which is common among those physicians who regard themselves as the only real doctors working in the hospital service.

Surgeons fail to meet his exacting requirements as they feel they must operate or prescribe the latest drugs. Radiologists and bacteriologists are well-meaning non-clinicians with insufficient knowledge of the patient. Anaesthetists are notorious expanders of empire.

To avoid the inevitable demarcation disputes surrounding patient care in a large hospital, Dr. Paton modestly proposes that all specialists should be subordinated to general physicians. If specialized medical knowledge is of such little value, would it not be more logical to subordinate us all to general practitioners or, better still, to medical social workers?—I am, etc.,

JAMES MCIVOR

Hillingdon Hospital,
Uxbridge, Middlesex

SIR,—I would like to add my humble dissent to the many that you must surely receive following Dr. A. Paton's Personal View (6 March, p. 554).

His reference to anaesthetists is an insult to my colleagues in the specialty and, far worse, it is untrue. Anaesthetists are medical practitioners, in addition to being "first class respiratory technicians"; they are quite capable of discussing prognosis with relatives and treating renal failure. The reason they do not normally practise this is probably on account of the ethics and etiquette that Dr. Paton sometimes finds lacking in medical practice.

The hazards of anaesthesia are no less, and often greater, than those of surgery. The pre and post operative intimacy of the anaesthetist with the patient must therefore be in the best interest of the patient, which Dr. Paton is so concerned about. His question "when do they have time to give anaesthetics?" ought really to be "when do they find time to see the patients they anaesthetize?"

Specialization has come to stay and the trend will continue in the years to come. If we can work together, with the welfare of the patient and his relatives foremost in our minds at all times, Dr. Paton will have nothing to fear.—I am, etc.,

J. L. HANDY

Department of Anaesthetics,
Freedom Fields,
Plymouth

SIR,—May an anaesthetist comment on part of the egregious Personal View propounded by Dr. A. Paton of Birmingham (6 March, p. 554)? If the seventh paragraph of my Personal View ran:

"Patients suffering from chronic bronchitis due to historical accident are frequently treated by general physicians, notorious for their deep grasp of theoretical considerations

and ineffectual floundering in the practical management of these diseases. I am sure they have high intellectual attainments but they need reminding that most of their time used to be spent in purging and bleeding dehydrated and anaemic patients. I doubt if they can really be expected to administer intravenous injections or perform lumbar punctures. And yet I hear that in addition to their treatment of respiratory disease, the giving of public health advice and medical administration generally, they are now presuming in many cases to prescribe psychotropic drugs. How do they find the time to instruct every other group of specialists, I wonder?" I could scarcely complain if my physician friends objected.

It should not be necessary to point that the publication of this sort of nonsense is likely to give offence. There are surely enough divisive tendencies in society without the gratuitous introduction to the profession of factitious interspecialty resentments. I would like my imaginary paragraph (which the saints forbend anybody take seriously) read in parallel with that of Dr. Paton to which I take exception: Can he say mine is unfair?—I am, etc.,

JOHN P. CONROY

St. Lawrence's Hospital,
Dublin 7

SIR,—Dr. Paton's Personal View (6 March, p. 554) convinces me that he is not a consultant physician but a general practitioner. I agree with everything that he says, but would add that I have paranoid delusions about consultant physicians taking over control of my patients.

I am glad that he used inverted commas when referring to "his" patients.—I am, etc.,

A. J. EARL

Barrow-upon-Soar
near Loughborough,
Leics.

Incontinence

SIR,—The letters (23 January, p. 229) commenting on Mr. K. E. D. Shuttleworth's article (19 December, p. 727) draw attention to the importance of physiotherapy in the treatment of stress incontinence.

Since 1965 the Sphincter Research Unit at Exeter, financed by the Medical Research Council, has been investigating the extension of this treatment by electronic stimulation. I would like to stress three points.

Firstly, I have been surprised at the number of cases referred from centres all over the country and elsewhere who have been "cured" simply by conscientious management.

Secondly, the very high proportion of patients who have been told that there was no treatment for incontinence and had suffered from this condition for years.

Thirdly, the surprising number who attend with a diagnosis of "incontinence" but without the primary cause being diagnosed.

I agree with your correspondents that much more publicity should be given to the possibility of cure in all these cases; to the medical and paramedical personnel, as well as to the public. Where possible cases should be referred to centres such as the one at Exeter or the Institute of Urology which are fully equipped to screen these cases, and proceed to treat them with whatever method is applicable, including the use of electronics. I would stress that the electronic treatments are an extension of