only unnecessary but may be positively dangerous, and should be carried out only for two reasons. One indication is to provide evidence where none other exists of a head injury having been sustained (the child who has been found unexplainably unconscious or where battering is suspected). The other is to confirm a clinical diagnosis of depressed fracture of more than trivial degree.

The really crucial part of the management of head injuries consists of careful observation in order to detect the onset of complications, haemorrhage, and meningitis. This is just as necessary for patients without a fracture as it is for those who have one, and this is the reason why I regard the taking of a skull x-ray as potentially dangerous. For if on the strength of a negative skull x-ray the patient is sent home and no warning given to the parents, they may have a false impression that all must be well and so be reluctant to associate any untoward symptoms that come on within the next 48 hours with the head injury.

What is needed is a period of careful observation by someone (the parents may be excellently suitable for this) who will notice the onset of untoward symptoms and, having been warned by the doctor to take them seriously in the unlikely event of their occurring, will not hesitate to bring the child back to the family doctor or the casualty department for re-appraisal. The decision whether to admit a child to hospital for observation following a head injury depends not on the presence or absence of a skull fracture but on the quality of the observation which is available in the home, the time of day, and, perhaps, the degree of disturbance of consciousness with which the injury is accompanied.

The lawyers ask for a skull x-ray simply because they have been wrongly led by the medical profession to believe that it is an essential part of the proper management of all head injuries. They would as readily ask for the burning of a candle at the bedside or the letting of a pint of blood if the profession had led them to suppose these things to be equally necessary.

Does the medical profession really believe that this wasteful and often frightening ritual non-investigation is always necessary? If so, let it say so and give convincing reasons for saying so. If not, let it speak out against it so that casualty officers and hospital junior staff can come off this absurd medicolegal hook.—I am, etc.,

JOHN BURKINSHAW

London S.E.26

Dahlak Blindness

SIR,—In your correspondence columns you published recently (25 December 1971, p. 811) a description of a corneal disease termed Dahlak blindness. This disease is said to be similar to Labrador keratopathy and Bietti's dystrophy. While surveying eye diseases among the Nama people of South-west Africa during July 1971, I found a similar corneal dystrophy to be present in these people. Fifty per cent of the Nama over 35 years of age show some form of corneal dystrophy. Like Dahlak dystrophy, and like Freedman's and Bietti's dystrophy, climate seems to play a part.

This dystrophy will be reported on in detail.—I am, etc.,

JEFFREY FREEDMAN

Department of Ophthalmology,
University of Witwatersrand,
Johannesburg, S. Africa

Childhood Hypoglycaemia

SIR,—Your leading article on "Childhood Hypoglycaemia" (1 January, p. 5) drew attention to the importance of this condition as a possible cause of brain damage. Recently two groups of workers^{1,4} have reported combined physiological, biochemical, and neuropathological studies of profound hypoglycaemia in primates. All animals developing brain damage had experienced blood glucose levels below 20 mg/100 ml for 2-10 hours. However, hypoglycaemia of this severity could last 3-5 hours without producing brain damage. Initially, the brain damage takes the form of "ischaemic cell change" affecting neurones in the neocortex, hippocampus, and cerebellum.⁵ Thus the lesions cytologically and in their pattern of selective vulnerability resemble those seen after cerebral hypoxia or oligoemia. The effects of arterial hypotension or cerebral hypoxia may be additive to those of hypoglycaemia in producing such damage.^{3,4}

These experiments were performed in adult animals and all cases subsequently showing brain damage had experienced periods of profound C.N.S. depression (usually with an isoelectric E.E.G.). Further experiments in newborn animals will be required to resolve the problem of brain damage occurring after symptomless episodes of hypoglycaemia.

Your article referred to the importance of convulsions as a symptom of hypoglycaemia. Less well recognized is the possibility of hypoglycaemia as a secondary feature of prolonged seizures. In children with status epilepticus blood glucose may be normal or elevated at the time of admission to hospital, but may subsequently fall to critical levels. Experimental studies in adolescent primates^{6,7} indicate that hypoglycaemia in status epilepticus may be severe enough to be a contributory cause to the permanent neurological disability commonly seen after prolonged seizures.⁸ It is advisable that children with prolonged seizures, if they are not receiving intravenous glucose, should have serial blood glucose determinations continuing well beyond the end of clinical seizure activity.—I am, etc.,

B. S. MELDRUM

M.R.C. Neuropsychiatry Unit,
Carshalton, Surrey

- 1 Khan, K. J., and Myers, R. E., in *Brain Hypoxia (Clinics in Developmental Medicine, 39/40)*, p. 185. London. Heinemann, 1971.
- 2 Myers, R. E., and Khan, K. J., in *Brain Hypoxia (Clinics in Developmental Medicine, 39/40)*, p. 195. London. Heinemann, 1971.
- 3 Meldrum, B. S., Horton, R. W., and Brierley, J. B., in *Brain Hypoxia (Clinics in Developmental Medicine, 39/40)*, p. 207. London. Heinemann, 1971.
- 4 Brierley, J. B., Brown, A. W., and Meldrum, B. S., in *Brain Hypoxia (Clinics in Developmental Medicine, 39/40)*, p. 225. London. Heinemann, 1971.
- 5 Brierley, J. B., Horton, R. W., and Meldrum, B. S., *Brain Research*, 1971, **25**, 483.
- 6 Brierley, J. B., Horton, R. W., and Meldrum, B. S., *Journal of Physiology*, in press.
- 7 Meldrum, B. S., Horton, R. W., and Brierley, J. B., *Electroencephalography and Clinical Neurophysiology*, in press.
- 8 Aicardi, J., and Chevrie, J. J., *Epilepsia*, 1970, **11**, 187.

Creatinine Clearance and Glomerular Filtration Rate

SIR,—Drs. W. M. Bennett and G. A. Porter (9 October 1971, p. 84) reported that the endogenous creatinine clearance is an accurate measurement of glomerular filtration rate and that the ratio of creatinine clearance to inulin clearance is almost unity. They postulated that the discrepancy between their results and our findings¹ was attributable to the normal values of inulin clearance reported by us being low and to the possibility that our patients were not in a steady state after preclearance water loading.

In 1966² we reported that normal values of inulin clearance were between 105 and 135 ml/min in our renal laboratory. These are similar to the values reported by Homer Smith.³ The values of creatinine clearance and inulin clearance in 37 normal men and women reported by us in 1969¹ were lower than those reported by us in 1966² and by Homer Smith³ because the age range of subjects was extended to between 20 and 68 years. The lower mean value for glomerular filtration rate reflects the lower values known to occur in normal older subjects.^{4,5} Our patients were in steady states after preclearance water loading as demonstrated by three 20-minute consecutive inulin and endogenous creatinine clearances, which were within 8%.

The fact that endogenous creatinine clearance overestimates actual glomerular filtration rate is reported not only by us but also by numerous other investigators, including Berlyne and colleagues.⁶ Reubi,⁷ in reviewing the literature, also concluded that in large groups of normal subjects the average creatinine/inulin clearance ratio does not differ significantly from unity, but in a given case the ratio is unpredictable and the creatinine clearance cannot be considered a precise measure of the glomerular filtration rate. There is great variability of the creatinine/inulin clearance ratio in individual cases. Our data¹ showed that the creatinine/inulin clearance ratio rises with decline of inulin clearance. These results are in agreement with those obtained by Reubi,⁷ who compared endogenous creatinine clearance with glomerular filtration rate measured by the clearance of thiosulphate.

A careful analysis of Bennett and Porter's data reveals a wide range of endogenous creatinine clearance at a given value of inulin clearance despite a linear correlation between inulin clearance and creatinine clearance. According to the authors' Figure 1, for an inulin clearance of about 100 ml/min endogenous creatinine clearances ranged from 75 to 130 ml/min, and at an inulin clearance of about 40 ml/min endogenous creatinine clearance ranged from 15 to 50 ml/min. Therefore we conclude, as we did in 1969,¹ that a single determination of endogenous creatinine clearance is unpredictable and cannot be considered a precise measure of the glomerular filtration rate, because of a wide range of endogenous creatinine clearance at a given value of inulin clearance. However, in serial determinations of endogenous creatinine clearance an excellent correlation was found between the rate of decline or increase in inulin and creatinine clearance. Therefore, endogenous creatinine clearance is very useful in following the direction of change of glomerular filtration rate during the course of renal disease.

We were also surprised to find in Bennett and Porter's data that clearances repeated one hour after an initial study during constant plasma inulin concentration and urine flow showed marked discrepancy between initial clearance and subsequent clearance in some patients. For example, one normal subject's inulin clearance went up from 105 ml/min to 133 ml/min. The other patient's inulin clearance went down from 101 ml/min to 76 ml/min. Glomerular filtration in man was found to be relatively constant throughout the 24-hour period when patients were confined to bed rest,⁸ and glomerular filtration rate in normal man is found to be remarkably constant on repeated measurement from day to day and from month to month over years.⁹

If we believe the data reported by Bennett and Porter with hour-to-hour variations physicians will be hard put to follow the change of glomerular filtration rate of patients with renal disease by either inulin clearance or endogenous creatinine clearance.—We are, etc.,

KWAN E. KIM

GADDO ONESTI

CHARLES SWARTZ

Hahnemann Medical College and Hospital,
Philadelphia, Pa., U.S.A.

- 1 Kim, K. E., Onesti, G., Ramirez, O., Brest, A. N., and Swartz, C., *British Medical Journal*, 1969, 4, 11.
- 2 Onesti, G., Kim, K. E., Swartz, C., and Brest, A. N., *Postgraduate Medicine*, 1966, 40, 327.
- 3 Smith, H. W., *Principles of Renal Physiology*, p. 32. New York, Oxford University Press, 1956.
- 4 Shock, N. W., *Geriatrics*, 1946, 1, 232.
- 5 McCance, R. A., in *Renal Disease*, ed. D. A. K. Black, p. 166. Oxford, Blackwell, 1962.
- 6 Berlyne, G. M., Varley, H., Nilwarangkur, S., and Hoerni, M., *Lancet*, 1964, 2, 874.
- 7 Reubi, F. C., *Clearance Tests in Clinical Medicine*, p. 13. Springfield, Charles C. Thomas, 1963.
- 8 Sirota, J. H., Baldwin, D. S., and Villarreal, H., *Journal of Clinical Investigation*, 1950, 29, 187.
- 9 Pitts, R. F., *Physiology of the Kidney and Body Fluids*, p. 68. Chicago, Yearbook Medical Publishers, 1968.

Myasthenia and Systemic Lupus Erythematosus

SIR,—Dr. Peter Isaacs (6 November 1971, p. 339), in his report of a patient with myasthenia gravis and systemic lupus erythematosus, argues that thymectomy would best be avoided in patients with both these diseases. I do not feel that this conclusion has either logical or empirical validity. It is based upon a report¹ of two myasthenic patients who developed lupus erythematosus several years after thymectomy, which had been followed by a remission of the myasthenia. Since, however, these two diseases have usually occurred together in non-thymectomized patients, one is not justified in concluding that thymectomy is unwise in the patient who already has both diseases, or in depriving the patient of the possibility of benefit to the myasthenia from thymectomy.

The interrelationship between myasthenia gravis and lupus is complex.² As would be expected from the known female preponderance in both diseases, almost all the reported patients are women. Either condition can occur first, and the interval between them has ranged from one month to 20 years. The effect of the development of lupus on the severity of the myasthenia has been quite variable. In several cases there were remissions of the myasthenia prior to or with the development of systemic lupus erythema-

tosis. There are other cases where the myasthenia was unaffected or worse as systemic lupus erythematosus evolved. In another case³ an excellent remission of the myasthenia occurred following thymectomy, despite antecedent lupus.—I am, etc.,

SHELDON M. WOLF

Southern California Permanente Medical Group,
Los Angeles,
California, U.S.A.

- 1 Alarcón-Segovia, D., Galbraith, R. F., Maldonado, J. E., and Howard, F. M., jun., *Lancet*, 1963, 2, 662.
- 2 Wolf, S. M., and Barrows, H. S., *Archives of Neurology*, 1966, 14, 254.
- 3 Mäkelä, P. E., Ruosteenoja, R., Wager, O., Wallgren, G. R., and Jokinen, E. J., *Acta Medica Scandinavica*, 1964, 175, 777.

Raised Elbows and Coronaries

SIR,—I have been interested for some years as a general practitioner to note how frequently it is possible to diagnose coronary artery disease in the preinfarction stage or when there is a small infarction. Fibrositis or gastric upset may well be explanations given to the patient—particularly the young patient—when family history, stress, blood pressure, cholesterol, and cardiographs are not abnormal. It is my experience that when findings are normal but suspicion remains, or when there is no other positive explanation, then if the patient is an "arm-mover" he or she has coronary artery disease, often proved later.

I define an "arm-mover" as one moving the elbows excessively or keeping them raised during conversation, relaxation, or sleep. I would be interested to hear of others finding this a significant diagnostic sign, as I am in no doubt as to its value.—I am, etc.,

ALAN N. RENNIE

Busby,
Glasgow

Caroli's Disease

SIR,—It is misleading, we feel, to dissociate Caroli's syndrome completely from choledochal cyst, as suggested by Dr. J. G. McNulty (4 December 1971, p. 623), for there is a spectrum of developmental abnormalities affecting the intrahepatic and extrahepatic biliary tree. A classical choledochal cyst may be associated with non-obstructive cystic or saccular dilatation of the left or right hepatic ducts and sometimes diverticula may arise from these ducts.¹ Cases of non-obstructive dilatation of the smaller intrahepatic ducts (Caroli's syndrome) may be associated with dilatation of the common bile duct² or even a choledochal cyst.³ Though the latter is usually an isolated finding, the presence of additional proximal biliary tree dilatation is of some clinical importance, for, as in Dr. Kelly's case (13 November 1971, p. 407), recurrent bouts of cholangitis are likely to follow removal or drainage of the choledochal cyst.

In Caroli's syndrome the hepatic parenchyma is normal, apart from some fibrosis and inflammation immediately adjacent to the dilated bile ducts, and portal hypertension has not been recorded. However, an identical pattern of dilatation of the intrahepatic biliary tree also occurs in some cases of congenital hepatic fibrosis,⁴ and in these cases portal hypertension with bleeding oesophageal varices is common but not in-

variable.⁵ We have recently seen two patients with congenital hepatic fibrosis in whom gross non-obstructive dilatation of the intrahepatic biliary tree with intractable cholangitis dominated the clinical picture and led to death. One of these cases also had unexplained dilatation of the common bile duct, again emphasizing the association of developmental abnormalities of the intrahepatic and extrahepatic biliary tree.—We are, etc.,

IAIN M. MURRAY-LYON

ROGER WILLIAMS

Liver Unit,
King's College Hospital,
London S.E.5

- 1 Tsuchida, Y., and Ishida, M., *Surgery*, 1971, 69, 776.
- 2 Caroli, J., Couinaud, C., Soupault, R., Porcher, P., and Etévé, J., *Semaine des Hôpitaux de Paris*, 1958, 34, 496.
- 3 Gots, R. E., and Zuidema, G. D., *American Journal of Surgery*, 1970, 119, 726.
- 4 Grumbach, R., Bourillon, J., and Auvert, J.-P., *Archives d'Anatomie Pathologique*, 1954, 30, 74.
- 5 Erlinger, S., Sakellaris, D. A., Maillard, J.-N., and Benhamou, J.-P., *Presse Médicale*, 1969, 77, 1189.

Leicester Royal Infirmary

SIR,—Leicestershire doctors will be delighted to read the excellent review of Mr. Frizelle and Miss Martin's book *The Leicester Royal Infirmary* (15 January, p. 187). There are, however, two small errors. C. J. Bond was a very famous Leicester surgeon, but the matron referred to in the review was a Miss Burt (not Bond), and her memory is still revered in both Leicester and at Guy's Hospital. Sir George McRobert's Aberdonian upbringing, together with his delight to discover that the founder of the Leicester Royal Infirmary was educated at Aberdeen, must have carried him away, so that he allocated Dr. William Watts to the Scottish clan of Watt. Personally I would like to think that we belong to the same clan, but south of the border we add a significant S.—I am, etc.,

CUTHBERT WATTS

Ibstock, Leics

Representation on the Joint Consultants Committee

SIR,—The report (*Supplement*, 1 January, p. 1) of the decision by the Central Committee for Hospital Medical Services acting on the advice of its executive to oppose the admission of the Regional Hospitals' Consultants and Specialists Association to the Joint Consultants Committee in no way underlines the importance of this decision, and I trust that you will allow me the courtesy of your columns to do so.

It may not be known that the R.H.C.S.A. has been applying for admission to the Joint Consultants Committee for over 20 years, and the recent attempt to do so again was entirely in accordance with the B.M.A. policy determined by the resolutions of successive Annual Representative Meetings in 1970 and 1971 (*Supplement*, 4 July 1970, p. 15; and 7 August 1971, p. 90). These called for an increase in regional consultant representation and a democratic system of election, but both have been disdainfully ignored by the Joint Consultants Committee.

The recent invitation by the chairman of the Joint Consultants Committee for Mr. R. E. Jowett and myself to journey to Edinburgh and give evidence in support of our application for membership of the J.C.C.