colleagues they should estimate the requirements for the next 12 months, and based on this assessment and the anticipated fractionation yields collect the necessary raw material within their region for the patients for whom they are directly responsible. This done, the ball is firmly in the court of the central fractionation units to make the necessary capacity available to prepare and deliver high quality products to the regional transfusion centres in the quantity requested. This principle should hold for all plasma fractionation products and would thus throw the burden of responsibility for patient care where it belongs—to the regional transfusion centres.—We are, etc.,

J. D. CASH
R. A. CUMMING

South-east Scotland Regional
Blood Transfusion Centre,
Royal Infirmary,
Edinburgh

The Red Cell

SIR,—Dr. G. W. G. Bird's article on "The Red Cell" (29 January, p. 293) prompts me to record some of my own observations. I have found¹ that the geometry of the avian egg is that of conic sections. This discovery was made possible by the assumption that the contents of the egg are paracrystalline and that the shape is derived from the figures related to liquid crystals.²

The following shapes are found in connexion with red blood corpuscles: biconcave discs, signet rings, crescents, cups, flattened elliptical discs in the camel family, and spheres when the cells are placed in hypotonic solution and pointed spines in the so-called "thorn apples" found when they are placed in hypertonic saline. Furthermore, the cells may range from 6-9 microns in any one individual.

These shapes are readily derived from conic sections. The biconcave disc is a cyclide figure, derived from confocal conic structure, that comprises a family of curves, so covering variation in size of the red cell. The signet ring is a cyclide with the hole or dimple off-centred and the crescent is a broken anchor ring; the cup a distorted anchor ring where the dimple in one side is exaggerated and in the other is everted. The red cell of the camel is a distorted ellipsoid figure, also found in the red cells of ovipara.

When placed in hypotonic saline, the red cell absorbs water and swells up into the form of a sphere that can be regarded as a limiting figure of the cyclide or ellipsoid. Placed in hypertonic saline, however, it loses water, ceases to be in the mesophase, assumes crystalline structure, and displays prismatic radiation. These findings will be of interest to haematologists, as red cells of signet ring shape are found in disease, notably malaria. Crescents are found in sickle-cell anaemia and spherocytosis is found in the disease of that name.

These observations provide strong evidence for the paracrystalline nature of haemoglobin in the red blood corpuscle. The outer "membrane," which might be described better as the sheet of the figure, is an interface between the mesophase in the cell and that of the plasma. The importance of the discovery, however, is in application to genetics. Sickle-cell anaemia, for example, is of genetic origin, likewise spherocytosis. The cyclides are derived from the ellipse and the

hyperbola and these have parametric descriptions, which, in turn, suggests a mathematical approach to the subject of genetics. It may well be found to apply to the theory of blood grouping.—I am, etc.,

J. E. MALCOLM

Royal Air Force Hospital,
Ely, Cambs

¹ Malcolm, J. E., Arris and Gale Lecture, Royal College of Surgeons of England, 20 January 1972, to be published.

² Hartshorne, N. H., and Stuart, A., *Crystals and the Polarising Microscope*, 4th edn. London, Edward Arnold, 1970.

Bacteriological Aspects of Mycobacterial Infections

SIR,—Professor D. A. Mitchison (12 February, p. 424) lists mycobacteria and, although writing in a section under the general heading of tuberculosis, includes *M. marinum* (*M. balnei*) and *M. ulcerans*, which cause ulceration of skin. *M. leprae* is not mentioned, perhaps because it cannot be cultured on artificial media. It may, however, be found in sputum of patients with advanced lepromatous leprosy involving the larynx and trachea. Although the diagnosis of leprosy will not be in doubt, as pulmonary tuberculosis not infrequently complicates neglected lepromatous leprosy, confusion between the mycobacteria may arise.—I am, etc.,

FREDERICK J. WRIGHT

Department of Tropical Medicine,
Lister Institute,
Edinburgh 8

Tropical Splenomegaly, Sickle-cell Trait, and P. falciparum Infection

SIR,—The letter of Dr. D. I. K. Evans and others (22 January, p. 250) entitled "Tropical Splenomegaly, Sickle-cell Trait, and P. falciparum Infection" needs to be commented upon from this part of the world, where enlarged spleens, sickle-cell traits, and *P. falciparum* infections are an everyday occurrence. In spite of the fact that the authors stated "This case has all the features of tropical splenomegaly syndrome" the evidence they presented could easily be interpreted to mean that the little Nigerian boy was not suffering from the tropical splenomegaly syndrome (T.S.S.), which, surely, has well known and accepted clinical and diagnostic features. Some of these features are:

A large spleen, which various investigators quoted in the authors' own letter specified to be at least 10 cm below the costal margin.^{1,2} The boy described had a spleen of 7.5 cm, a size which literally hundreds of West African boys would have without being considered here to be in the T.S.S. range.

Abdominal discomfort as a result of the size of spleen, with dragging pain, debility, and sometimes fever and anaemia.^{3,4} There were no febrile episodes in the case reported, and the abdominal pain could as well be explained by ascariasis, which the patient had.

There is a tendency to an increase in IgM, with IgG levels not significantly different from normal.² The authors found the very opposite, stating "Hyperglobulinaemia is present, but IgM is not so often raised as IgG", and in support of this statement they cited three authors,^{1,2,5} none of whom,

in fact, support their statement. It was surprising that Dr. Evans and colleagues omitted to refer to the work of Loewenthal *et al.*,⁶ who measured IgM values in 17 of 19 cases of suspected T.S.S. and found all greatly increased.

It usually takes at least 3-6 months' continuous drug therapy before shrinkage of the spleen occurs.^{5,6} The Nigerian boy's spleen shrank to one-third its original size after six weeks of antimalarial therapy.

In view of the above (and there was no liver biopsy, which, I agree, would not have been justified merely to prove T.S.S. in this boy), may I suggest, Sir, that there was no evidence whatsoever that this boy had the tropical splenomegaly syndrome. I submit that the cause of the splenomegaly which responded so well and fast to antimalarial therapy was chronic falciparum malaria infection. The child came from an endemic area and had by then developed a considerable amount of immunity. As immunity develops, parasitaemia decreases, but malaria parasites can still be found in small numbers. He also had multiple helminthic infestation, a likely cause for abdominal pain and even splenomegaly in children. Chronic malarial infection and helminthic infestation would account for the high IgG.

Finally, as tropical splenomegaly is not proved, the problem of sickle-cell trait in this case is irrelevant, and so are the authors' recommendations. Indeed, workers here have long ceased to call every AS pattern sickle-cell trait. The authors do not tell how they excluded sickle-cell β -thalassaemia, in which 8-year-olds often have a large spleen, an AS pattern, and *P. falciparum* infection.—I am, etc.,

MARIANNE JANOSI

Department of Medicine and Therapeutics,
University of Ghana Medical School,
Accra

¹ Watson-Williams, E. J., and Allan, N. C., *British Medical Journal*, 1968, 4, 793.

² Saeco, A. S., *British Medical Journal*, 1970, 3, 378.

³ *British Medical Journal*, 1970, 3, 360.

⁴ Marsden, P. D., and Hamilton, P. J. S., *British Medical Journal*, 1969, 1, 99.

⁵ Stuijver, P. C., Ziegler, J. L., Wood, J. B., Morrow, R. H., and Hutt, M. S. R., *British Medical Journal*, 1971, 1, 426.

⁶ Loewenthal, M. N., O'Riordan, E. C., and Hutt, M. S. R., *British Medical Journal*, 1971, 1, 429.

Prevention of Deep Vein Thrombosis

SIR,—We are delighted that Mr. N. H. Hills and others (15 January, p. 131) have confirmed our finding (13 November, p. 394) that intermittent calf compression is an effective prophylactic against deep vein thrombosis and would congratulate them on presenting such a thorough analysis of their trial.

Our own trial of the efficacy of intermittent calf compression was preceded by a rigorous haemodynamic study, which was conducted both in greyhounds¹ and in patients² in order to find the most effective method of applying the compression. This investigation, as well as our previous one into the effects of passive flexion (10 July 1971, p. 78), revealed, despite Mr. Hills's reservations to the contrary, that when changing the flow in one leg the flow in the other remains largely unchanged. The techniques of statistical analysis used by ourselves and by Mr. Hills and his colleagues are, therefore, equally acceptable.

It is, however, surprising to find that our