dates. In reply, Sir Keith suggested that I was basing my judgement of the future on my experience of the past, and that he believed that Government pressure on local authorities would be successful in persuading them to find the vast sums of money required. Does anyone else believe this?—I am, etc.,

W. A. HEATON-WARD

Stoke Park Hospital Group,
Stapleton,
Bristol

¹ Department of Health and Social Security. *Better Services for the Mentally Handicapped*. Cmd. 4683. London, H.M.S.O., 1971.

Benign Forms of Multiple Sclerosis

SIR,—May I draw your attention to an error in the leading article on "Benign Forms of Multiple Sclerosis" (12 February, p. 392) where it states: "In 1959 McAlpine and N. Compston surveyed 241 cases of multiple sclerosis . . ." The paper which your reviewer must have had in mind was published in 1961.¹ A further point—these 241 cases were seen within three years of onset—not a truly prospective study as implied but a "quasi-prospective" one (a term I use in the new edition of *Multiple Sclerosis: A Reappraisal*). I would be glad if this error could be put right.—I am, etc.,

DOUGLAS McALPINE

Marnhull, Dorset

¹ McAlpine, D., *Brain*, 1961, 84, part II, 186.

**Dr. McAlpine is quite right and we apologize for the error.—ED., *B.M.J.*

Haemophilus Epiglottitis

SIR,—We read with great interest the letter from Dr. Patricia E. Mortimer and others (26 February, p. 567) regarding their views on the management of children suffering from haemophilus epiglottitis. Since children with acute supraglottic oedema are prone to sudden obstruction of the airway we feel that early laryngoscopy should be carried out when the presence of acute epiglottitis is suspected. If this disorder is the cause of the obstruction, endotracheal intubation should produce relief provided that the tube has been correctly placed. The disease is usually of limited duration, and if efficient therapy is instituted tracheostomy would not appear to be warranted.

Dr. Mortimer and her associates consider that fulminating toxæmia was the main factor initiating cardiac asystole in their cases. However, they did not indicate whether determination of PaO_2 , PaCO_2 and pH were made. It may well be that the instances of cardiac asystole they describe resulted not from toxæmia but from a combination of hypoxæmia and metabolic acidosis. This would be in keeping with the clinical history of two hours' mild respiratory obstruction preceding sudden clinical deterioration.

It should be emphasized that all instances of respiratory difficulty in the child require continuous observation together with ready availability of blood gas determinations.—We are, etc.,

G. W. BLACK
S. R. KELTY
S. H. S. LOVE

Royal Belfast Hospital for Sick Children,
Belfast

Head Injuries in Children

SIR,—I would like to congratulate Dr. J. H. Burkinshaw on his excellent letter (5 February, p. 378) concerning the value of x-ray examination in the assessment of head injury in children. I hope his letter will be widely read and noticed, and that it will produce a suitable response.

I have long been influenced by Dr. Burkinshaw in this matter and aware of his previous publications. I am in no doubt that he is correct in what he says. Of course the assessment of the severity of head injuries in children is a difficult clinical matter, but I am sure that the right decision is made by careful history taking and clinical observation. The use of an x-ray is an adjunct which is by no means always helpful. Unfortunately in my present position I have not felt able always to follow the practice he advocates, or to recommend it to my junior medical staff. This is because, as Dr. Burkinshaw points out, one is continually aware of the current medico-legal view. The law still places unwarranted importance on the x-ray examination. One of the difficulties in establishing Dr. Burkinshaw's view in the profession has been the lack of senior influential doctors in accident and emergency departments. Most of these cases are seen by junior staff who are unsupervised.

I hope that now that the specialty of accident and emergency work has been recognized it will be possible for the more senior doctors observing and recording cases themselves to improve the standard of assessment of all head injuries.—I am, etc.,

DAVID CARO

St. James's Hospital,
London S.W.12

A Practical Electrocardiograph

SIR,—As a relatively junior doctor who has attended many emergency resuscitative procedures in my capacity as medical and anaesthetic house officer in various hospitals, I am concerned about the standard and basic design of the equipment supplied to deal with these emergencies. Over the past 30 such emergencies, it has taken an average of around 10 minutes to produce a recognizable trace on an E.C.G. This time margin obviously has many variables, but one of the most time-consuming factors is the manipulation of large, heavy, and complicated equipment to the scene, and even worse, connecting this to the patient with separate straps, electrodes, coloured leads, and quantities of a proprietary jelly. I would like, therefore, to make the following suggestions for a practical E.C.G.

The E.C.G. oscilloscope and defibrillator must be combined. Only very recently have two British manufacturers even managed to produce a portable battery-operated D.C. defibrillator, while in America they have had a combined instrument weighing only 35 lb for over two years now. The latter instrument also ingeniously uses the defibrillator paddles as monitoring electrodes. I would also like to see significant theoretical advance incorporated in the design of these instruments, for example, the double square pulse method of defibrillation as described by Kugelberg.¹

It must be small, light, and portable, with a fast warm up high input impedance, and common mode rejection and with three power

supplies—mains, external 12 volt (car battery), and optional rechargeable battery pack, the latter two, for use by general practitioners, etc. outside hospital. Modern lightweight plastics, electronic integrated circuits, and recent advances in battery design make these criteria relatively simple to attain.

Above all, the monitor-defibrillator should be simple with the minimum of controls. In an emergency situation, there is no need for switched leads, gain, calibration controls, rate meters, internal defibrillation, and synchronized D.C. shock and all the other little-used electronic sophistication which only makes the equipment heavier and more complex to operate. One is dealing with asystole or ventricular fibrillation in the great majority of cases. Therefore only a single lead II is required and as the higher frequency components of the trace are not important in this situation, a restricted bandwidth of 3 db. down at 20 Hz would reduce noise, myographic, and mains frequency interference. An optional plug-in pen recorder with switched leads for a permanent trace would be a useful accessory.

An improved electrode system must be adopted. It is time the ancient practice of rubber straps, plate electrodes, yards of wire, and messy electrode jelly was scrapped.² The suction "multipoint" type of electrode, as described by Lewes,³ has a definite advantage in speed of application, as have the "paddle electrodes" mentioned earlier. However, a quicker and more efficient idea would be three spring-loaded "pincer" electrodes with needle points which would pierce the stratum corneum and hold the electrode firmly in place, require no electrode jelly, be very quick in application, and give a low value of skin resistance. The discomfort experienced on application and the risks of cross infection are negligible in comparison to the emergency situation which exists. However, the electrodes could be made detachable for sterilization.—I am, etc.,

JOHN B. DONALD

Department of Anaesthetics,
Western General Hospital,
Edinburgh

¹ Kugelberg, J. E. W. B., *Medical and Biological Engineering*, 1968, 6, 167.

² Turner, R. W. D., *Lancet*, 1966, 1, 1158.

³ Lewes, D., *World Medical Electronics*, 1966, 4, 240.

Sustained Hyperglycaemia in Pregnancy

SIR,—In their article on the effect of sustained maternal hyperglycaemia on the fetus in normal and diabetic pregnancies (19 February, p. 466) Dr. N. W. Oakley and others have provided evidence of great value both to obstetricians and to physicians with a special interest in the management of pregnancy in diabetics.

They formulated a hypothesis that hyperglycaemia in pregnant diabetics produced excessive secretion of insulin in the fetus, and that the resulting hypoglycaemia was responsible for the known tendency of such babies to die suddenly in utero before or during labour. To test this hypothesis, they gave diabetic and normal mothers in whom labour was to be induced a large intravenous glucose load, and confirmed that the fetuses of the diabetic mothers did produce excessive insulin in response to this. Fetal hypoglycaemia did not occur, though one fetus became asphyxiated and acidotic. The outcome of the pregnancies is not stated.