Year	1962	1963	1964	1965	1966	1967	1968	1969	1970
Anencephaly rate	1.82	1.78	1.73	1.66	1.65	1.56	1.47	1.50	1.51

for 1970, and this followed a low potato blight year, 1969.

Dr. J. H. Renwick's table of the figures for 1962-9 (20 January, p. 172) (taken from mid-February to mid-February and including live-born anencephalics), which are already corrected for regression of -0.06, show this same constancy. His diagram, which shows no zero, magnifies the small annual variations in the incidence of anencephaly in comparison with those for blight.

In view of this uniform rate of decline in anencephaly it is unlikely that the marked annual variation in the percentage of blighted potatoes, or indeed any other agent which shows marked annual variation, could be making a substantial contribution to the incidence of anencephaly. If there were one main environmental agent responsible for anencephaly it must be something that has been steadily declining in strength over the decade.—I am, etc.,

C. O. CARTER

M.R.C. Clinical Genetics Unit,
Institute of Child Health,
London W.C.1

SIR,—Scottish statistics on anencephaly have been extensively quoted in epidemiological studies in recent years, most recently by Dr. J. H. Renwick,¹ who correlated the stillbirth rate from anencephaly in Scotland with the presumed teratogenic insult from blight in the susceptible areas in the west of Scotland. He estimated that the incidence of blight was moderately severe in 1950 and 1953 and found that the stillbirth rate from anencephaly was higher in 1951 and 1954 than in other years. The table below gives the stillbirth rate from anencephaly in Scotland from 1947 to 1956 arranged according to the parity of the mother. It shows that the overall rate per 1,000 total births increased from 2.4 in 1950 to 2.7 in 1951. It was higher in 1st and 5th+ pregnancies but lower in 2nd, 3rd, and 4th pregnancies. In 1954, while the overall rate was higher than in 1953, the rise occurred in 2nd and 3rd pregnancies and a slight fall occurred in 1st and 5th+ pregnancies. These trends were repeated inside each social class. Analysis of the same data by the age of the mother, using the technique of the five-years moving average, shows that the rate rose in the 15-19 age group from 1947 onwards, in the 20-24 age group from 1952, and in the 25-29 age group from 1958.

In his more recent letter (20 January, p. 172) Dr. Renwick makes a similar comparison for England and Wales in the years 1961-8 and draws similar conclusions although the rates are lower and the variations much smaller. If a teratogen acting during pregnancy, such as potato blight, is so important as to be responsible for 95%

of the deaths from anencephaly, one would expect that it would act on women of all ages and parities alike in the years in which its concentration had reached dangerous levels. The fact that the stillbirth rate from anencephaly was rising from 1950 onwards in Scotland even in the years when blight in the previous year was negligible also raises doubt as to the probability of a causal relationship.—I am, etc.,

DUGALD BAIRD

M.R.C. Medical Sociology Unit,
Centre for Social Studies,
Aberdeen

¹ Renwick, J. H., *British Journal of Preventive and Social Medicine*, 1972, 26, 67.

Gastric Ulcer after Highly Selective Vagotomy

SIR,—Mr. A. Hall's report (30 December, p. 789) on gastric ulcer after highly selective vagotomy is important, and a reminder of the unknown outcome of this operation despite the encouraging early reports from Leeds and Copenhagen. I have no reason to doubt the rationale of this procedure and no cause for complaint in observing the progress of almost 100 patients treated in my unit. Nevertheless I do not think this operation should be generally adopted at this stage.

Mr. Hall reports postoperative secretion studies, which are of interest but would have been even more valuable if compared with preoperative tests. A proper assessment of the operation demands good facilities for measuring gastric function as well as a reliable follow-up. Unless this can be done we are in danger of discarding operations with known results in favour of a fashionable technique whose true value may be missed by inadequate appraisal. Not long ago a distinguished Continental surgeon described our gastric surgeons as "individualists, each of whom had his own branch of the vagus which he divided or preserved according to his ideas." We could improve our image by exercising some patience until the way is clear.—I am, etc.,

C. G. CLARK

Surgical Unit,
University College Hospital Medical School,
London W.C.1

SIR,—I was interested to read Mr. R. Hall's letter (30 December, p. 789) regarding a case of benign lesser curve gastric ulcer following proximal gastric vagotomy (highly selective vagotomy) without drainage done apparently for basal duodenal and pyloric channel disease.

Three years ago I began using proximal

gastric vagotomy for duodenal ulceration, preserving the prepyloric vagal supply to 5-6 cm. One patient developed later a benign lesser curve gastric ulcer, and gastric retention was shown using the food/barium meal. A gastroenterostomy was done later to drain the antrum and to cure the ulcer. Amdrup (personal communication) had a similar case, also with retention.

Preservation of the vagal supply to 5-6 cm from the pylorus invites gastric retention and benign lesser curve gastric ulcer. I think that Amdrup in Denmark and Hedenstedt in Sweden, as well as Johnston and myself in this country, are retaining much more innervated stomach and all choosing the same point of preservation. This is in fact the point that Holle in Munich has used for many years. It is easily chosen above the point where the main anterior nerve of Latarjet crosses the lesser gastric curve. About 8-9 cm of stomach remains innervated and food/barium studies show that there is no retention post-operatively in the absence of organic stenosis in the pyloric canal or in the duodenum.

Damage to the nerves of Latarjet may too, of course, cause gastric retention and benign lesser curve gastric ulcer, and great care is needed at operation to protect these nerves.—I am, etc.,

HAROLD BURGE

West London Hospital,
London W.6

Radiography of Potentially Pregnant Females

SIR,—It is to be hoped that a full discussion of this subject will be prompted by the letter from Drs. G. M. Ardran and F. H. Kemp (18 November, p. 422). Ultimate agreement on the responsibilities of clinician, patient, and radiologist within an accepted scheme for avoiding this hazard would be an ideal conclusion.

The code of practice issued by the Department of Health¹ states that radiological examinations involving the lower abdomen should, if practicable, be carried out within the first 10 days following the first day of the menstrual period. The practical difficulties of a strict application of a "10-day rule" are considerable. The clinical staff of the Aberdeen hospitals discussed this matter with the radiologists in 1967. While accepting their responsibility for ensuring that patients who might be pregnant were not referred for x-ray unless absolutely necessary, the clinicians could not give a guarantee to include invariably the date of the last menstrual period on the request form. The radiologists undertook to ascertain the date of the last menstrual period in cases of booked appointments occurring some time after the original request; they felt that a rigid application of a "10-day rule" for outpatients was virtually impracticable. The system which has been in force for the past five years is, in effect, a "28-day rule." It applies to outpatient appointments for barium examinations, intravenous pyelography, and micturating cystography previously booked on all female patients (married or unmarried) aged between 15 and 50 years. This simply means that the date of the last menstrual period is obtained by a nurse and/or radiographer and transmitted to the radiologist when the patient arrives at the x-ray department.

Pregnancy	Stillbirth Rate from Anencephaly per 1,000 total births (Scotland)									
	1947	1948	1949	1950	1951	1952	1953	1954	1955	1956
1st	2.1	2.7	2.2	1.8	3.1	3.7	3.6	3.3	2.6	3.1
2nd	1.4	1.6	1.5	2.2	1.8	1.6	1.7	2.8	2.1	2.3
3rd	2.2	2.1	2.6	2.8	1.8	1.9	2.4	2.8	2.9	2.9
4th	2.5	3.3	2.3	3.4	2.6	3.6	3.1	3.1	3.5	3.2
5th	4.2	4.1	2.3	3.8	4.4	3.5	2.8	2.6	4.5	3.3
All	2.2	2.5	1.1	2.4	2.7	2.9	2.8	3.0	2.9	2.9