Should the patient appear to have missed one menstrual period a fresh appointment is given 7-14 days ahead; the patient is provided with a pre-paid form addressed to the radiologist, whom she notifies in due course whether or not menstruation has occurred. In the latter event the x-ray examination is cancelled and the general practitioner or consultant so informed. The actual number of cases in which rebooking is required has been very small. This system is not ideal in that it does not prevent the possible irradiation of a 10- to 18-day fetus; it does, however, safeguard an older fetus and offers minimal delay and inconvenience for all concerned.

It would be of great value in this discussion to have comments on the following:

(1) What are the views of the clinicians and radiologists on the strict application of the "10-day rule"? Is it possible to apply it to non-emergency inpatients without plunging hospital wards and radiology departmental administration into chaos?

(2) Ideally all fertile female patients who are to receive irradiation to the lower abdomen should be given the opportunity of considering a possible pregnancy. Are clinicians willing to accept a method such as that suggested by Drs. Ardran and Kemp which depends on a prior consultation between clinician and patient in every case?

(3) Failing this, is a system as described above and operated solely by radiologists acceptable?—I am, etc.,

LEWIS A. GILLANDERS

Royal Infirmary,
Aberdeen

¹ Department of Health and Social Security, *Code of practice for protection of persons against ionising radiations arising from medical and dental use*. London, H.M.S.O., 1972.

Vitamins in Illness

SIR,—In "Any Questions" (2 September, p. 582, and 30 December, p. 786) the question whether vitamins should be given in acute illness, or at other times, is discussed.

At the meeting of the Nutrition Society on 7 December 1972¹ a paper was read showing that elderly people in a large hospital had an average vitamin C level of 12.5 $\mu\text{g}/10^8$ W.B.C., confirming earlier work.^{2,3} Last winter a Medical Research Council team reported that over one-third of elderly people, in and out of hospital, in South Wales had vitamin levels of 0.2 mg per 100 ml plasma, or 15 $\mu\text{g}/10^8$ W.B.C., or below this level (personal communication).

Recent work in the United States^{4,5} showed that haemorrhages around hair follicles begin after 26 days of complete deprivation of vitamin C in young men previously given large amounts of vitamin C. Haemorrhages on the under surface of the tongue began a few days later. The perifollicular skin haemorrhages are regarded by the American authors as "almost pathognomonic of scurvy." They began at plasma levels of vitamin C at or above 0.2 mg/100 ml, correspondingly to 15 $\mu\text{g}/10^8$ W.B.C. These skin haemorrhages occur often in 20% of elderly hospital patients in Britain. Sublingual haemorrhages are even more frequent (personal findings). Blood levels of vitamin C at or below 15 $\mu\text{g}/10^8$ W.B.C. are very common in elderly people in Britain. Low levels of vitamin B₁ are frequently found with low vitamin C levels.³ According to the Depart-

ment of Health,⁶ elderly people eat much the same kind of food as the general population, though in smaller quantities.

Tetracycline, and possibly other antibiotics, cause a fall of vitamin C in the blood, often to below 15 $\mu\text{g}/10^8$ W.B.C. The levels of some of the B group of vitamins in the blood also fall when tetracycline is given.⁷ Some hospitals give supplements of vitamin C and the B group of vitamins to all patients. The evidence now available suggests that this should be a universal practice, so that in illness, at least, all people have the recommended daily allowance of vitamin C and the B group of vitamins. Recent Canadian⁸ and Scottish surveys⁹ suggest that much larger amounts of vitamin C, 1 g. daily, may reduce the days lost by illness by 30% and reduce the incidence and duration of the common cold, as Linus Pauling has suggested.¹⁰—I am, etc.,

GEOFFREY TAYLOR

Ilminster, Somerset

¹ Andrews, J., Atkinson, S. J., Ridge, B. D., and Wyn-Jones, C., *Abstract of Communications: Nutrition Society Meeting*, December, 1972.

² Kataria, M. S., Rao, D. B., and Curtis, R. C., *Gerontologia Clinica*, 1965, 7, 189.

³ Brocklehurst, J. C., et al., *Gerontologia Clinica*, 1968, 10, 309.

⁴ Hodges, R. E., Baker, E. M., Hood, J., Sauberlich, H. E., and March, S. C., *American Journal of Clinical Nutrition*, 1969, 22, 535.

⁵ Hodges, R. E., Hod, J., Canham, J. E., Sauberlich, H. E., and Baker, E. M., *American Journal of Clinical Nutrition*, 1971, 24, 432.

⁶ Stephen, J. M. L., *Health Trends*, 1972, 4, 76.

⁷ Windsor, A. C. M., Hobbs, C. B., Treby, D. A., and Cowper, R. A., *British Medical Journal*, 1972, 1, 214.

⁸ Anderson, T. W., Reid, D. B. W., and Beaton, G. H., *Canadian Medical Association Journal*, 1972, 107, 503.

⁹ Charleston, S. S., and Clegg, K. M., *Lancet*, 1972, 1, 1401.

¹⁰ Pauling, L. C., *Vitamin C and the Common Cold*. San Francisco, W. H. Freeman, 1970.

Staffing of Accident and Emergency Departments

SIR,—I would like to support the suggestion made by Dr. P. A. M. Weston (13 January, p. 114) that a period of training in an accident and emergency department would be a valuable prerequisite to the entry to general practice. Approaches to the Royal College of General Practitioners and the B.M.A. in the past on the possibility of making this an essential (as it is with the Royal College of Surgeons) have been turned down because of inadequate organization and lack of instruction in many departments.

An essential feature of making such a condition to entry to general practice would be a system of inspection and registration of accident and emergency departments recognized for this purpose. This, if properly carried out, would provide the strongest possible stimulus to improvements in the facilities and management in these departments as a means of ensuring job satisfaction and proper staffing. Another benefit might well be the closure of many existing casualty departments where there was no justification for pursuing the required standard.—I am, etc.,

J. C. SCOTT

Accident Service
United Oxford Hospitals

Shortage of Radiologists

SIR,—Owing to an impossible work load on radiologists in the Plymouth clinical area there have had to be reductions in the ser-

vice provided. This has included the closing of the radiodiagnostic department to general practitioners, though chest x-rays can still be obtained through the chest clinic. Because of this a meeting of general practitioners passed the following resolution: "The meeting regrets and protests at the withdrawal of general practitioner radiological facilities in the Plymouth area and considers this an unnecessary hindrance to the delivery of normal general practitioner services. The meeting proposes that a system of rationing of radiological services be introduced immediately, whereby individual general practitioners may request a limited number of x-rays each week."

In the light of the above and the fact that there are 70 unfilled radiologist posts in the country I have been asked to write to you urging that the Association use every means available to bring this need to the notice of the Department of Health and Social Security with a view to rectifying the situation as soon as possible.—I am, etc.,

J. P. GRIER

Honorary Secretary,
Plymouth Division, B.M.A.

Hartley, Plymouth

Working of the Mental Health Act

SIR,—My attention has been drawn to the article by your legal correspondent (30 December, p. 800) dealing with the case of V.E. (Mental Health Patient).¹ I write as chairman of the tribunal stating the case in this particular matter. There is one matter of interest to your readers which became apparent during the course of the hearing but appears only as a side issue in the three judgements.

Section 4 of the Mental Health Act 1959 lays down a statutory definition of "mental disorder" which I understand is rather different from the usual clinical definition adopted by psychiatrists. It is necessary to classify the form of mental disorder into one of various categories, *all of which are mutually exclusive*. Thus a patient must be stated to be suffering from any one of the classifications set out in section 4 of the Act—namely, "mental illness," "arrested or incomplete development of mind," "psychopathic disorder," and "any other disorder or disability of mind." "Arrested or incomplete development of mind" is subdivided into "severe subnormality" and "subnormality."

For the purpose of section 25, a patient who is suffering from any form of mental disorder of a nature or degree which warrants the detention of the patient in hospital under observation may be detained for a limited period for observation. Under section 26 only those patients who are classified as suffering from four specific types of mental disorder—namely, "mental illness," "severe subnormality," "subnormality," or "psychopathic disorder"—may be detained for treatment. Of these forms of mental disorder, only "mental illness" is not defined by the Act; however, for the reasons given above, if the form of disorder comes within the definition of "psychopathic disorder" the patient cannot be classified as suffering also from "mental illness," unless the patient is at one and the same time suffering from two quite separate and distinguishable forms of mental disorder.

The statutory definition of "psychopathic disorder" (namely, "a persistent disorder or disability of mind . . . which results in . . .

seriously irresponsible conduct on the part of the patient and requires or is susceptible to medical treatment") was clearly applicable in the case of this patient. It followed therefore that as the conduct was caused by brain damage, the Act required the classification to be that of "psychopathic disorder" and not "mental illness," with the consequences which are set out in your correspondent's article.

I seriously doubt whether any mental disorder which is a disability of the mind caused by brain damage can be classified as "mental illness" for the purposes of the Act. This would not prevent a classification of "mental illness" if such a patient suffered from some secondary disorder also (such as dementia), amounting to mental illness.

The position of a patient who suffers from mental illness which results in "seriously irresponsible conduct," etc., is unclear, having regard to the watertight compartments laid down in section 4. It is difficult to see how a classification of "mental illness" in such a case could be faulted.—I am, etc.,

JAMES A. COOKE

Chairman,
Mental Health Review Tribunal for the
South-west Metropolitan Regional Hospital
Board Area

London W.C.1

¹ In re V. E. (Mental Health Patient) *The Times*,
7 July 1972 (1972) 3 W.L.R. 669.
² *All England Law Reports*, 1972, 3, 373.

Deaths Associated with General Dental Anaesthesia

SIR,—Dr. M. P. Coplans and Mr. I. Curson (13 January, p. 109) report that evidence at a coroner's court is seldom adequate for determining the precise cause of death in these cases; if the investigator is to make a valid diagnosis, they suggest, it has to be supplemented through personal communication with those present in the surgery when the collapse occurred.

With this I entirely agree. Applying this principle I have, over the years, investigated a considerable number of these fatalities and very rarely indeed has the correct explanation emerged from the coroner's court. Often the inquest is delayed several weeks, by which time the evidence of those most closely involved in the shattering catastrophe tends to be influenced by hindsight and other considerations. Nor is the expert witness always as informed and impartial as the coroner may suppose. And since nearly all the victims of this sudden unexpected collapse are young and healthy, necropsy is essentially negative. The pathologist will report oedema of the lungs and congestion of the viscera, which are common findings, but to explain the death he has recourse to such suggestions as asphyxia (even though there had been no cyanosis or other sign of suffocation under the anaesthetic, only pallor), idiosyncrasy to one of the agents used (nitrous oxide, trichlorethylene, halothane, and methohexitone have all been blamed), "anaesthetic shock," vagal inhibition associated with thymic hypertrophy, or even (in 1970) "status thymolymphaticus."

Truth and science would be better served were a panel of experts, at least two in number and of the highest academic standing, commissioned the moment a death associated with general dental anaesthesia was notified to hold an immediate on-the-

spot inquiry and duly report to the coroner.—I am, etc.,

J. G. BOURNE

Salisbury, Wilts

Anthrax

SIR,—Dr. Robert Lamb's article on anthrax (20 January, p. 157) mentions the disinfection of imported hair and wool, which has recently been discontinued.

The Government Wool Disinfecting Station in Liverpool was opened in 1921 but was closed some two years ago. The reasons given by the Department of Employment for its closure were that considerable quantities of imported hair and wool were coming in through other ports, and that the manufacturers of these raw materials were now sufficiently equipped to cleanse the consignments of hair and wool as part of the manufacturing process. During its existence the station dealt with many hundreds of consignments of imported hair and wool. These were tested bacteriologically before and after being treated by the Duckering process. Very many of the consignments were heavily infected with anthrax spores on receipt. However, the process was so effective that I cannot recall a positive result after the material was treated.

As these infected consignments of wool and hair are now being widely distributed to various parts of the country I would think it worth while for the Department of Employment to prepare a list of manufacturing processes in which these potentially infected materials are used so that doctors seeing patients working in these processes may be alerted to the anthrax hazard. If it is important, as our tropical disease colleagues exhort us, to ask the patient "Where have you been?", I would suggest that for any patient visiting his doctor with an inflammatory skin lesion it is equally important for the doctor to ask "What materials do you handle at work?"—I am, etc.,

ANDREW D. SEMPLE

Port Health Authority,
Liverpool

Suicide Rate

SIR,—The suicide statistics of the past decade are remarkable for the steady decline in the suicide rate, beginning in 1964. That decline is present in all age and sex groups (see table); the young have not been excluded as Dr. D. Lester (9 December, p.

Suicide Rates in England and Wales per 100,000 by Age and Sex, 1963 and 1970^a

Age (years)	Men			Women		
	Rate		% Change	Rate		% Change
	1963	1970		1963	1970	
15-24	6.9	6	-13	3.1	2.6	-16
25-44	14	10	-29	9.6	5.9	-39
45-64	26	16	-38	17	12	-29
65-74	34	21	-38	20	15	-25
75+	39	24	-38	18	9.7	-46
All ages	14	9.5	-32	9.9	6.6	-33

612) states, provided we look at the most recent statistics.¹ It is true that the suicide rate for the 15-24 age group has not gone down as much as the older age groups' suicide rates, but it would be hasty to infer,

as Dr. Lester would, that difficulty in treating psychological disorder in the young is the explanation. Trends in admission rates for the younger age groups in fact run counter to those for the older part of the population. Between 1964 and 1970 first admission rates for the 15-19 age group increased² while rates for the other age groups decreased. Secular trends in suicide statistics are often easier to observe than explain.—I am, etc.,

B. M. BARRACLOUGH

M.R.C. Clinical Psychiatry Unit,
Graylingwell Hospital,
Chichester, Sussex

¹ Registrar General, *Statistical Reviews for England and Wales*, 1963 and 1970, London, H.M.S.O., 1965, 1972.

² Department of Health and Social Security, *Psychiatric Hospitals and Units in England and Wales*, Statistical and Research Report Series No. 4, 1970, London, H.M.S.O., 1972.

Radiological Pseudotumour in Chronic Ulcerative Colitis

SIR,—In two cases of ulcerative colitis radiological evidence indicating stricture due to tumour formation was obtained. In neither case was a tumour or stricture found to be present at operation; nevertheless, histological examination at these sites showed cancer to be present.

Case 1.—A woman aged 34 known to have had ulcerative colitis for 11 years was having six to eight bowel actions a day at the time of her admission to hospital. The stools were fluid, containing blood, pus, and mucus. She had colicky pain in the right iliac fossa. On examination no abdominal tumour could be felt. A barium enema showed extensive ulcerative colitis of long standing with a constant stenotic lesion at the hepatic flexure (fig. 1). This lesion

FIG. 1.—Barium enema showing changes of chronic ulcerative colitis with constant stricture at the hepatic flexure. Case 1.

was still present at a second barium enema undertaken a month later. Panproctocolectomy was therefore performed. At operation neither stenosis nor tumour could be found, though the whole colon was markedly thickened, with increased vascularity. Examination of the colon after removal revealed no evidence of cancer either as a stricture or a tumour. Histological examination showed a moderately well-differentiated adenocarcinoma to be present at the site of the radiological tumour. No penetration of the bowel wall had occurred and no lymph nodes were involved. She remains well after 17 years' follow-up.

Case 2.—A 28-year-old man developed septic