and is available from Down Brothers, Mayer and Phelps Ltd., Church Path, Mitcham, Surrey.—I am, etc.,

LAURENCE TINCKLER

Maelor General Hospital,
Wrexham

Treatment of Status Asthmaticus

SIR,—A noteworthy point on which I differ from the opinion expressed in your commendable leading article (9 December, p. 563) is in the statement that "physiotherapy has no place in the treatment of status asthmaticus." It has been our custom during the past three years in this hospital (which is part of Helsinki University Central Hospital), in addition to other routine treatment in cases of status asthmaticus, to facilitate the patient's breathing by manual pressure on the lower part of his chest during

This position is the most beneficial in status asthmaticus for movement of the diaphragm if the patient is still conscious or even semi-conscious.

every expiratory phase (see fig.). Measurements (performed with a Spirostat flow meter) show that this method expels 100-200 (or even 300) ml more air from the lungs than would happen spontaneously. In my opinion this respiratory help has often been decisive. It was used in 1972 in the management of 153 patients with status asthmaticus, none of whom died.

We have also diverged from the choice of drugs recommended in your article. For instance, we administer methylprednisolone intravenously instead of the hydrocortisone preparation suggested by you. This prevents retention of fluid and hypokalemia, while the antiallergic effect is at least equally good. I would like to point out, too, that ordinary adrenaline is often a more efficacious drug for asthma than beta-2-sympathomimetic preparations.—I am, etc.,

KLAUS A. J. JÄRVINEN

Hospital for Allergic Diseases
Helsinki, Finland

Viral Infection and Renal Transplant Rejection

SIR,—Dr. J. D. Briggs and others (2 December, p. 520) reported acute rejection of transplanted kidneys in three patients infected with influenza A virus and suggested that this might be due to a generalized stimulation of the host's immune response. I should like to suggest another mechanism

by which the virus may have caused rejection.

Influenza A virus has long been known to agglutinate erythrocytes.^{1,2} After a short time the erythrocytes separate and cannot be agglutinated a second time. This agglutination is due to the virus attaching itself to receptor sites on two different red cells and linking them together. The cell receptor sites are glycoproteins containing sialic acid, probably in a terminal position³ which is removed by a potent neuraminidase possessed by the virus, thus allowing the cells to separate once again.^{4,5} Neuraminidase-sensitive receptors have been found on other cells.⁶⁻⁸

When sialic acid is contained in an antigen it is very rarely antigenic itself. Indeed, it seems that sialic acid may serve to conceal more potent antigens and prevent them from stimulating an immune response, for treatment of many types of cell with neuraminidase has been shown to render them more highly antigenic.⁹⁻¹³ There is a possibility, therefore, that influenza virus may cause acute rejection of a renal transplant by removing terminal sialic acid groups from glycoprotein molecules in the renal cells and rendering them highly antigenic. If this hypothesis is correct there should be no danger of causing acute rejection by vaccinating against influenza so long as the virus is inactivated in such a way as to render it incapable of digesting sialic acid from renal cells.—I am, etc.,

W. T. MORRIS

Royal Postgraduate Medical School,
London W.12

- 1 Hirst, G. K., *Science* 1941, **94**, 22.
- 2 Hirst, G. K., *Journal of Experimental Medicine*, 1942, **76**, 195.
- 3 Winkler, R. J., *International Review of Cytology*, 1970, **29**, 77.
- 4 Cohen, A., in *Mechanisms of Virus Infection*, W. Smith, London, Academic Press, 1963.
- 5 Gottschalk, A., *Physiological Reviews* 1957, **37**, 66.
- 6 O'Connell, C. J., and Milgrom, F., *Proceedings of the Society for Experimental Biology and Medicine*, 1966, **122**, 1255.
- 7 Peleg, B. A. and Ianculescu, M., *Nature*, 1966, **211**, 1211.
- 8 Pepper, D. S., and Jamieson, G. A., *Nature*, 1968, **219**, 1252.
- 9 Springer, G. F., *Bacteriological Reviews*, 1963, **27**, 191.
- 10 Currie, G. A., and Bagshawe, K. D., *British Journal of Cancer*, 1968, **22**, 843.
- 11 Kemp, R. B., *Nature*, 1968, **218**, 1255.
- 12 Bagshawe, K. D., and Currie, G. A., *Nature*, 1968, **218**, 1254.

Immunological Responses in Pregnancy

SIR,—The letter from Dr. P. Brain and others (25 November, p. 488) drew my attention to an earlier article by Dr. Ronald Finn and others (15 July, p. 150) reporting reduced T-cell activity during pregnancy. They found that the number of circulating T lymphocytes capable of responding to stimulation with phytohaemagglutinin was reduced in pregnant women by about 10-20% and that the incidence of negative tuberculin reactions was increased by about 64%.

Quite apart from the fact that some other authors have failed to find a deficiency in T-cells during pregnancy,^{1,2} the depression of tuberculin reaction could perhaps be attributed to hormonal changes in the course of pregnancy. In view of the most marked increase of oestrogen production it should be mentioned that although oestrogens have been shown to depress humoral antibody responses³ and cause involution of the thymus⁴ there is no conclusive evidence that

they affect peripheral immunologically competent cells,⁵ and their effects on various cell-mediated immune reactions are rather controversial.⁶ The possibility should be considered, however, that reduced tuberculin reactions during pregnancy could be the result of non-specific actions of oestrogens. There is evidence that these hormones alter the physicochemical properties of the ground substances of connective tissue⁷ and it has been reported that they restrict the spreading of materials and fluids in the skin.⁸ In fact, the reduced tuberculin reaction of oestrogen-treated animals has been attributed to this mechanism rather than to depression of immunological reactivity.⁹—I am, etc.,

V. S. ŠLJIVIĆ

St. Mary's Hospital Medical School,
London W.2

- 1 Brain, P., Marston, R. H., and Gordon, J., *British Medical Journal*, 1972, **4**, 488.
- 2 Watkins, S. M., *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 1972, **79**, 990.
- 3 Warr, G. W., and Šljivić, V. S., *Experimentia* 1972, **28**, 1356.
- 4 Dougherty, T. F., *Physiological Reviews*, 1952, **32**, 379.
- 5 Šljivić, V. S., and Warr, G. W., *British Journal of Experimental Pathology*, 1973. In press.
- 6 Šljivić, V. S., and Warr, G. W., *Periodicum Biologorum* (Zagreb), 1973. In press.
- 7 Hytten, F. E., and Leitch, I., *The Physiology of Human Pregnancy*, 2nd edn. Oxford, Blackwell, 1971.
- 8 Sprunt, D. H., McDearman, S., and Raper, J., *Journal of Experimental Medicine*, 1938, **67**, 159.
- 9 Lurie, M. B., *Annals of the New York Academy of Sciences*, 1950, **52**, 1074.

Role of Pacing after Myocardial Infarction

SIR,—The mortality figures quoted by Dr. Helen H. Tucker and others in their study (6 January, p. 10) seem to provide some statistical evidence that electrical pacing of patients who develop complete heart block following a myocardial infarction may, in most instances, not be necessary. In view of their findings a controlled study of the problem might appear useful, but provided the necessary skills and facilities are available, few would dispute the advantages of pacing over an isoprenaline infusion in a patient who, at that very time, is at greatest risk of developing a dangerous arrhythmia. It would be of interest to know the reasons why the authors decided to adopt the non-pacing policy in their patients.

The role of pacing in preventing ventricular arrhythmias is certainly not confined to patients with heart block. Its value in the control of recurrent ventricular tachycardia and fibrillation not responding to drug therapy is not emphasized in most reports. Although these refractory dangerous arrhythmias often denote a poor prognosis because of underlying severe myocardial damage, a very gratifying response can sometimes be achieved with a combination of pacing and antiarrhythmic drugs, as two cases recently treated at this hospital demonstrate.

The first patient, a man of 56, suffered within a period of 24 hours from eight episodes of ventricular tachycardia, going on rapidly to ventricular fibrillation, two weeks after an extensive myocardial infarct. In spite of combined administration of all antiarrhythmic drugs available, the dangerous arrhythmias continued unabated until the intrinsic heart rate was over-paced to a rate of 120-130/min so that all ventricular ectopic beats could be suppressed. The pacemaker catheter was removed a week later, sinus rhythm has continued since discharge from hospital, and normal life has been resumed.

The second patient, a hypertensive man of 33, had suffered a myocardial infarct at the age of 31. A permanent internal pacemaker with a fixed rate of 70 impulses/min was inserted a year later because of permanent complete heart block. Subsequently he was readmitted following episodes of ventricular tachycardia and fibrillation and within a period of eight hours required 21 defibrillations. In spite of full drug therapy the episodes were occurring with increasing frequency until the catheter wire was exposed and connected to an external pacemaker box so that the impulses could be increased to 110/min. The only episode of ventricular arrhythmia which occurred from then on was when one of the connexions became loose, and this was quickly rectified. Full recovery has occurred and the rate of the pacemaker has now been cut down to 70/min as he continues with complete heart block.

This form of therapy should seriously be considered whenever these dangerous arrhythmias do not respond to drug administration.—I am, etc.,

M. E. BENAIM

Royal Victorian Infirmary,
Newcastle upon Tyne

Persisters

SIR,—In your leading article (13 January, p. 63) on "Infectious Disease" you refer to the discovery of the persister state by virtue of which bacteria become phenotypically resistant to certain antimicrobials, notably penicillin. The reference given was to a paper by Crofton published in 1969.¹ For the sake of historical accuracy I wish to point out that persisters were first described by the late Professor J. W. Bigger in a paper entitled "Treatment of Staphylococcal Infections with Penicillin" published in 1944.² Bigger says, "Despite the well-marked bactericidal power of penicillin, attempts to sterilize broth containing *Staphylococcus pyogenes* often fail because, among the millions of cocci present, there are a few which are insensitive to penicillin. These I call 'persisters' . . . Persisters are believed to be insensitive to penicillin because they are in a dormant, non-dividing phase . . ."

This paper of Bigger's, in which he also presents conclusive evidence that the action of penicillin is a bactericidal one, is a minor classic. So much so that, having gone back to it again after having read your leader, I have recommended it, and will continue to recommend it, to students as a model of research procedure and presentation.—I am, etc.,

F. S. STEWART

Department of Bacteriology,
Trinity College,
Dublin

¹ Crofton, J., *British Medical Journal*, 1969, 2, 209.
² Bigger, J. W., *Lancet*, 1944, 2, 497.

Safety-belt Snag

SIR,—I was interested in the case reported by Dr. S. S. Sanders (13 January, p. 114) because I am at present treating an active 65-year-old man who sustained severe comminuted fractures of both bones of his right leg in a similar fashion—catching his rear foot in a loose-hanging safety belt while getting out of a car.

Obviously this serious hazard should be given greater publicity, and perhaps the owners of cars without spring-loaded seat belts would thereby be encouraged to hang up the belts when not in use. It must be in

the interests of insurers and insured to see that something is done about this.—I am, etc.,

C. J. MCCORMACK

Portlaoise, Ireland

SIR,—I wholly endorse the views expressed by Dr. S. S. Sanders on car safety belts (13 January, p. 114). I recently treated an old lady who had suffered a fractured neck of a femur after tripping over a safety belt which hung out of her car. The commonly seen belts are either too short to be quickly fitted when safely stowed away or too long to be safely stowed away when set at the correct length for use, and I agree that spring-loaded belts should supercede the usual fittings found in the majority of vehicles.—I am, etc.,

M. R. WARREN

Milnthorpe,
Westmorland

SIR,—I read with interest the letter from Dr. S. S. Sanders (13 January, p. 114) regarding the dangers of car safety belts. While I agree with him that these straps should be worn, as they prevent injury, there is no doubt they have also caused a number of serious accidents. My wife recently broke her ankle by catching her foot in such straps, which were lying on the floor, as she was leaving a car. This stimulated my interest in the subject, and since then several examples of this have come to my notice. They range from broken hips to sprained ankles. Such accidents might be expected to be caused only by loose straps and not by the variety that are supposed to retract automatically. Unfortunately, this is not so, as some straps which are meant to retract automatically do not do so. The safety straps used by service pilots retracted on a wire cable activated by a strong spring, but it appears that car seat belts will sometimes not retract automatically as the spring controlling the inertia reel is not strong enough.

I shall be glad to receive details of accidents caused by supposedly self-retracting safety straps, as there appears to be a strong case for modifying their design.—I am, etc.,

NEIL S. PAINTER

Manor House Hospital,
London N.W.11

Case for Free Contraceptives

SIR,—No one wants unwanted pregnancies. On this all agree, but you make two assumptions in your leading article (20 January, p. 130) which we would challenge.

The first assumption, implied in your article, is that the first cause of unwanted pregnancy is the omission of contraception rather than the commission of the irresponsible and selfish act(s) of sexual intercourse.

The second assumption is that doctors will be willing to prescribe contraceptives to all comers, married and unmarried, and that we can do nothing to mitigate the permissive society our patients, especially the young, have to live in. These two facets are related because our prescribing will be read as, at least, failure to show disapproval of the sexual licence undermining marriage and the family.

Universal, "free" contraception may reduce the incidence of abortion operations three months later and of unwanted babies nine months later, but irregular sexual activity will produce inexorably, as all doctors see daily, a harvest of broken hearts, broken homes, and homeless or one-parent children—not to mention venereal disease. This is not theorizing about morals but an immediate practical issue. It is most regrettable that you, speaking for doctors as a whole, simply accept that irresponsible, permissive behaviour will go unchecked.—We are, etc.,

JEAN R. C. BURTON-BROWN
S. HAMILL
T. D. SAYER
N. H. TUCKER
G. J. FIELD

J. LOVELESS
ELEANOR A. THOMPSON
J. K. N. YEARSLEY

Margate

Broadstairs

Ramsgate

J. D. BEALE
R. STOWE
D. A. J. NEDEN

Controlling Inflation

SIR,—I was disgusted by your unpatriotic leading article "Controlling Inflation—Second Stage" (27 January, p. 1971). What does it matter how much consultants get paid so long as this country is saved from inflation?—I am, etc.,

E. M. WRIGHT

Coombe Bissett,
Salisbury, Wilts

Serum Lithium Estimations

SIR,—The paper by Dr. B. von Hartitzsch and others on permanent neurological sequelae despite haemodialysis for lithium intoxication (30 December, p. 757) brings a reminder of the dangers of the valuable drug lithium carbonate. In this connexion I should like to raise a disquieting matter: the inaccuracy of serum lithium estimations.

Here are three examples from my recent experience. A perceptive patient of mine noticed that the results were higher when he went to his family doctor's pathologist than when he went to mine, so he visited both on the same day. The first pathologist reported a serum lithium of 0.6 mEq/l. and the second pathologist on a sample taken 10 minutes later reported a level of 1.0 mEq/l. Another patient of mine, a conscientious fellow who could be relied upon to take his 1,200 mg of lithium carbonate per day if anybody could, was coasting away over the months at a serum lithium level of between 0.4 and 0.6 mEq/l. when suddenly, according to the teaching hospital which does the estimations, a level of 2.0 mEq/l. was reached. He was without any side effect when I saw him next a fortnight later, never had had any, and was still taking his 1,200 mg daily. This time I tried my first patient's trick and asked another teaching hospital to make a simultaneous estimation. The results were 0.2 and 0.46 mEq/l. respectively. At another hospital I started an inpatient on 1,600 mg of lithium carbonate per day. On the sixth day, as she began to show side effects the drug was withdrawn for a few days but recommenced at 800 mg per day.